Name	Siddhesh Sonar		
UID no.	2021700063		
Experiment No.	1		

AIM:	To implement the various functions e.g. linear, non-linear, quadratic, exponential etc.					
Program						
PROBLEM STATEMENT :	For this experiment, we have to implement at least 10 functions from the list. The input (i.e. n) to all the above functions varies from 0 to 100 with increment of 10. Then add the function n! in the list and execute the same for n from 0 to 20 with increment of 2.					
ALGORITHM/ THEORY:	In this experiment part 1-A we must use 10 functions from the given list write a simple C code for it and then use the output from the code to draw a graph for the 10 functions + the factorial function and draw our conclusion regarding the functions.					
PROGRAM:	<pre>#include <stdio.h> #include <math.h> unsigned long long fact(int n) { if (n == 0) { return 1; } else { // for (int i = n - 1; i > 0; i) // { //</math.h></stdio.h></pre>					

```
return n * fact(n - 1);
float func1(int n)
       //(3/2)^n
        return pow(1.5, n);
int func2(int n)
       // n^3
       return pow(n, 3);
float func3(int n)
       //(lg^2)^*n
       return pow(log2(n), 2);
float func4(int n)
       // sqrt(log(n))
       return sqrt(log2(n));
float func5(int n)
       // n log n
        return n * log2(n);
float func6(int n)
       // ln ln n
       return log(log(n));
float func7(int n)
       // log n
        return log2(n);
```


```
float func8(int n)
        // 2^n
        return pow(2, n);
float func9(int n)
        // ln n
        return log(n);
int func10(int n)
        // n+5
        return n + 10;
void print1(int start, int end, int inc, int (*func)())
        for (int i = start; i \le end; i = i + inc)
        printf("%d = \%d\n", i, func(i));
void print2(int start, int end, int inc, float (*func)())
        for (int i = start; i \le end; i = i + inc)
        printf("%d = \%f\n", i, func(i));
void print3(int start, int end, int inc, unsigned long long (*func)())
        for (int i = start; i \le end; i = i + inc)
        printf("%d = \%lld\n", i, func(i));
int (*intf)(int);
float (*floatf)(int);
unsigned long long (*longf)(int);
int main()
```

```
printf("\ln(3/2)^n\ln^n);
floatf = func1;
print2(0, 100, 10, floatf);
printf("\n^3 \n');
intf = func2;
print1(0, 100, 10, intf);
printf("\ln(\lg^2)*n\ln'n");
floatf = func3;
print2(0, 100, 10, floatf);
printf("\nsqrt(\log(n))\n'n");
floatf = func4;
print2(0, 100, 10, floatf);
printf("\nn log n \setminus n');
floatf = func5;
print2(0, 100, 10, floatf);
printf("\nln ln n\n');
floatf = func6;
print2(0, 100, 10, floatf);
printf("\nlog n\n\n");
floatf = func7;
print2(0, 100, 10, floatf);
printf("\n2^n\n'");
floatf = func8;
print2(0, 100, 10, floatf);
printf("\nln n \setminus n');
\overline{\text{floatf}} = \overline{\text{func9}};
print2(0, 100, 10, floatf);
printf("\n+5\n'");
intf = func 10;
print1(0, 100, 10, intf);
printf("\n!\n");
longf = fact;
print3(0, 20, 2, longf);
return 0;
```

RESULT:	3/2)^n
RESULI.	(3/2) II
	0 = 1.000000
	10 = 57.665039
	20 = 3325.256836
	30 = 191751.062500
	40 = 11057332.000000
	50 = 637621504.000000 $60 = 36768468992.000000$
	70 = 2120255143936.000000
	80 = 122264599134208.000000
	90 = 7050392827330560.000000
	100 = 406561191922499584.000000
	n^3
	0 = 0
	$\frac{0-0}{10=1000}$
	20 = 8000
	30 = 27000
	40 = 64000
	50 = 125000
	60 = 216000
	$\frac{70 = 343000}{80 = 512000}$
	90 = 729000
	100 = 1000000
	(lg^2)*n
	$\frac{0 = \inf}{10 = 11.035206}$
	20 = 18.679062
	30 = 24.077576
	40 = 28.322918
	50 = 31.853113
	60 = 34.891357
	$\frac{70 = 37.568111}{80 = 39.966774}$
	90 = 42.144157
	100 = 44.140823
	$\operatorname{sqrt}(\log(n))$
	$\frac{0 = -\text{nan}}{10 = 1.822616}$
	10 - 1.022010

20 = 2.07892530 = 2.21515040 = 2.30693150 = 2.37568060 = 2.43041070 = 2.47573980 = 2.51434490 = 2.547911100 = 2.577568n log n 0 = -nan10 = 33.21928020 = 86.43856030 = 147.20671140 = 212.87712150 = 282.19281060 = 354.41342270 = 429.04980580 = 505.75424290 = 584.266785100 = 664.385620ln ln n 0 = -nan10 = 0.83403220 = 1.09718930 = 1.22412840 = 1.30532350 = 1.36405560 = 1.40960770 = 1.44656580 = 1.47751190 = 1.504035100 = 1.527180log n 0 = -inf10 = 3.32192820 = 4.32192830 = 4.90689040 = 5.32192850 = 5.643856


```
60 = 5.906890
70 = 6.129283
80 = 6.321928
90 = 6.491853
100 = 6.643856
2^n
0 = 1.000000
10 = 1024.000000
20 = 1048576.000000
30 = 1073741824.000000
40 = 1099511627776.000000
50 = 1125899906842624.000000
60 = 1152921504606846976.000000
70 = 1180591620717411303424.000000
80 = 1208925819614629174706176.000000
90 = 1237940039285380274899124224.000000
100 = 1267650600228229401496703205376.000000
ln n
0 = -inf
10 = 2.302585
20 = 2.995732
30 = 3.401197
40 = 3.688879
50 = 3.912023
60 = 4.094345
70 = 4.248495
80 = 4.382027
90 = 4.499810
100 = 4.605170
n+5
0 = 10
10 = 20
20 = 30
30 = 40
40 = 50
50 = 60
60 = 70
70 = 80
80 = 90
90 = 100
```



GRAPH:


	(lg^2)*n	sqrt(log(n))	n+5	ln ln n	log n	ln n	
T .	inf	-nan		-nan	-inf	-inf	
10	11.035206	1.822616	20	0.834032	3.321928	2.302585	
20	18.679062	2.078925	30	1.097189	4.321928	2.995732	
30	24.077576	2.21515	40	1.224128	4.90689	3.401197	
40	28.322918	2.306931	50	1.305323	5.321928	3.688879	
50	31.853113	2.37568	60	1.364055	5.643856	3.912023	
60	34.891357	2.43041	70	1.409607	5.90689	4.094345	
70	37.568111	2.475739	80	1.446565	6.129283	4.248495	
80	39.966774	2.514344	90	1.477511	6.321928	4.382027	
90	42.144157	2.547911	100	1.504035	6.491853	4.49981	
100	44.140823	2.577568	110	1.52718	6.643856	4.60517	

The above are columns B, C, D, E, F, G and H respectively



	n^3	n log n	n!	(3/2)^n	2^n	
0	0	-nan	1	1	1	
10	1000	33.21928	2	57.665039	1024	
20	8000	86.43856	24	3325.256836	1048576	
30	27000	147.206711	720	191751.0625	1073741824	
40	64000	212.877121	40320	11057332	1.09951E+12	
50	125000	282.19281	3628800	637621504	1.1259E+15	
60	216000	354.413422	479001600	36768468992	1.15292E+18	
70	343000	429.049805	87178291200	2.12026E+12	1.18059E+21	
80	512000	505.754242	2.09228E+13	1.22265E+14	1.20893E+24	
90	729000	584.266785	6.40237E+15	7.05039E+15	1.23794E+27	
100	1000000	664.38562	2.4329E+18	4.06561E+17	1.26765E+30	

The above are columns I, J, K, L, M and N respectively



CONCLUSION:	We observer that due to the large exponential values in some of the functions some of
	the graph become very huge and we also learned how to create graphs using excel