

Existence and Localization of a Limit Cycle in a Class of Benchmark Biomolecular Oscillators

Sidhanta Mohanty, Shaunak Sen

Abstract—Oscillatory behaviour is important in multiple biological contexts. However, the inherent nonlinearity and high dimensionality of mathematical models in biology makes proving the existence and the localization of limit cycle oscillations challenging. Here, we provided an elementary proof for the existence and a method for rigorously localizing the oscillatory solutions in a class of benchmark biomolecular oscillators. To construct the proof, we used a geometric approach based on Brouwer’s Fixed Point theorem. We constructed a toroidal-like manifold within a positively invariant set by removing the hypervolume containing the fixed point and the trajectories converging to it along its stable manifold. We showed that the vector field describing the system dynamics maps a cross section of the toroidal-like manifold onto itself. The existence of a limit cycle solution in this manifold was guaranteed by Brouwer’s Fixed Point theorem. For different sets of initial conditions in these cross-sections, we used an interval-based Reachability Analysis to localize the oscillatory behaviour. These results add a simple and elegant approach to demonstrating the existence of limit cycles in biomolecular systems as well as a method for rigorous localization of the region of existence.

Index Terms—Limit Cycle, Cyclic Gene Regulatory Network, Brouwer’s Fixed Point Theorem, Reachability Analysis.

I. INTRODUCTION

The Poincaré-Bendixson theorem provides a way to demonstrate the existence of limit cycle oscillations in two-dimensional systems [1]. The basic idea is to construct a positively invariant set without steady states so that the trajectories converge to the limit cycle. An explicit solution of the limit cycle is not needed. This theorem is not applicable in dimensions greater than two because of the possibility of more complex attractors and chaotic behaviour. Limit cycles in systems of dimension greater than two, however, are typically found in biological systems, where they underlie crucial control processes such as circadian rhythms and the cell cycle [2]. For example, the famous Goodwin oscillator has a negative feedback loop wrapping a chain of intermediate biomolecules so that the overall length of the chain is greater than or equal to three [3]–[5]. Another canonical example is a ring oscillator with an odd number (≥ 3) of genes that sequentially repress each other [6]. The experimental demonstration of the repressilator, a ring oscillator with three genes, heralded a new era for synthetic

biology [7]. Therefore, methods to establish the existence and rigorous enclosure of limit cycles in higher-dimensional systems are needed.

While various approaches have been used to prove the existence of limit cycles in biological systems of dimensions greater than or equal to three, the region of existence is relatively large and conservative. Like the Poincaré-Bendixson theorem, these approaches seek to establish the existence of a limit cycle without an explicit analytical solution. One approach is to analyze biological oscillators in a Boolean framework using kinetic logic models [6], [8]. A geometric approach based on Brouwer’s fixed point theorem was used to prove the existence of a limit cycle in the three-dimensional Goodwin oscillator [9], which we recently adapted for the repressilator [10]. This approach was subsequently generalized to an n -dimensional Goodwin oscillator [11]. This approach provides concrete insight into the limit cycle behaviour, but has been referred to as ‘elaborate’ [12] and ‘laborious’ [13]. An n -dimensional version of the Poincaré-Bendixson Theorem was developed by for monotone cyclic feedback systems [14]. The above ideas were used to address the existence and stability in ring oscillators and their variants [12], [15]. An explicit characterization of biological parameter sets for the existence of oscillations for the ring oscillators was also reported [16]–[18]. Recent work has expanded on the notion of 2-cooperativity to prove the existence of limit cycles with wide applicability in biology [19]–[21]. Typical estimates of the region of existence of the limit cycle obtained using the above approaches are conservative in nature, and rigorous tighter enclosures are desirable.

In this work, we localized the region of existence of the limit cycle in a benchmark biomolecular oscillators using the Brouwer’s Fixed Point theorem and interval-based Reachability Analysis. We provided an alternative, elementary proof for the existence of a limit cycle in cyclic gene regulatory network with odd number of genes. For the existence proof, we constructed a positively invariant set for oscillator dynamics by analyzing the direction of the vector field at the outer facets of a hypercube. We constructed a torus-like structure by removing a hypervolume associated with the unstable steady state and the trajectories approaching it along its stable manifold. We showed that a cross-section of this torus-like region maps onto itself, thus proving the existence of a limit cycle. For different sets of initial conditions on these cross-sections, we rigorously evolved the network dynamics using Interval Analysis to

Sidhanta Mohanty¹ and Shaunak Sen² are with the Department of Electrical Engineering, Indian Institute of Technology, New Delhi, India. {¹sidhanta.mohanty,²shaunak.sen}@ee.iitd.ac.in

localize the limit cycle.

II. EXISTENCE OF LIMIT CYCLE

In this section, an elementary proof for the existence of a limit cycle in m -gene cyclic regulatory network based on Brouwer's Fixed Point theorem is presented.

Let $x_1, x_2, \dots, x_m$ represent the proteins of the m -gene cyclic regulatory network. The differential equations describing the dynamics of the network are given in (1). A symmetric model with only the protein dynamics, ignoring mRNA dynamics, is chosen for simplicity.

$$\frac{dx_i}{dt} = F(x_{i-1}) - \gamma x_i, \text{ for } i = 1, 2, 3, \dots, m. \quad (1)$$

Here x_i is the concentration of the i -th protein with $0 \leq x_i \leq \frac{\alpha}{\gamma}$. α, γ, k , and n are positive constants, $F(p) = \frac{\alpha}{1 + (\frac{p}{k})^n}$ is the Hill function, n is the Hill coefficient, and m is an odd number. As the system is cyclic in nature, $x_0 = x_m$.

The steady state of the cyclic regulatory network in (1) is unique. The uniqueness can be demonstrated graphically [2]. Denote the steady state as $(x_{10}, x_{20}, \dots, x_{m0})$. So, $x_{10} = \underbrace{\frac{1}{\gamma} F\left(\frac{1}{\gamma} F \dots \left(\frac{1}{\gamma} F(x_{10})\right) \dots\right)}_{m\text{-times}}$. As the gene regulatory

network model is symmetric, the steady state is of the form $(x_0, x_0, \dots, x_0)$. The stability of this steady state can be assessed through the eigenvalues of the Jacobian of the linearization of the system in (1) at the steady state $(x_0, x_0, \dots, x_0)$. The eigenvalues of the Jacobian are: $\lambda_{1,\dots,m} = -\gamma + \rho \left[\cos\left(\frac{2p+1}{m}\pi\right) + j \sin\left(\frac{2p+1}{m}\pi\right) \right]$, where $p = 0, 1, \dots, m-1$ and $\rho = -F'(x_0)$. An eigenvalue is real and negative, $\lambda_1 = -(\gamma + \rho)$. The remaining eigenvalues are complex in nature. For certain values of p , the cosine component in the real part of the eigenvalues become negative. This implies that the corresponding eigenvalues will certainly have negative real parts. We used the instability of the steady state to prove the following sufficient condition for the existence of a limit cycle.

Theorem 2.1: There exists a limit cycle solution to the cyclic gene regulatory network in (1) if $\gamma < \max_{p \in \mathcal{P}_+} \left\{ \rho \cos\left(\frac{2p+1}{m}\pi\right) \right\}$, where $\mathcal{P}_+ = \left\{ p \in [0, m-1] : \cos\left(\frac{2p+1}{m}\pi\right) > 0 \right\}$.

Proof: For certain values of p , the cosine component in the real part of the eigenvalues becomes negative. This implies that the corresponding eigenvalues will certainly have a negative real part. This contributes to the possible stability of the steady state. However, the instability of the steady state implies that nearby trajectories are repelled from the steady state and may converge to a closed invariant set, such as a limit cycle. To ensure that the steady state is unstable, at least one of the complex conjugate pair of eigenvalues should have a positive real part. For the complex eigenvalues to have a positive real part, it is needed for $-\gamma + \rho \cos\left(\frac{2p+1}{m}\pi\right) > 0 \quad \forall p \in \mathcal{P}_+$, where,

$\mathcal{P}_+ = \left\{ p \in [0, m-1] : \cos\left(\frac{2p+1}{m}\pi\right) > 0 \right\}$. So, to ensure atleast one of the complex conjugate pair of eigenvalues have positive real part: $\gamma < \max_{p \in \mathcal{P}_+} \left\{ \rho \cos\left(\frac{2p+1}{m}\pi\right) \right\}$.

Upon establishing the instability of the steady state, a geometric approach is used to prove the existence of a limit cycle within the hypercube $0 \leq x_1, x_2, \dots, x_m \leq \frac{\alpha}{\gamma}$.

Construction of an Invariant Box: To demonstrate the existence of limit cycle in the m -gene cyclic regulatory network described by (1), we showed that the hypercube is a positively invariant set. This is done by computing the inner product (Ξ) between the vector field characterized by the system dynamics within the hypercube ($\vec{G}$) and the normal to the outer facets of the hypercube ($\hat{n}$): $\Xi = \langle \vec{G}, \hat{n} \rangle$. We considered the unit normal vectors for the different facets of the hypercube directed inward. The value of Ξ determines the direction of the flow of the vector field through the facets: (i) $\Xi > 0$ indicates the vector field flows through the facet in the direction of $\hat{n}$, (ii) $\Xi = 0$ indicates the vector field does not flow through the facet, and (iii) $\Xi < 0$ indicates the vector field flows through the facet in the direction opposite of $\hat{n}$.

The vector field described by the system dynamics in (1) is: $\vec{G}(x_1, x_2, x_3, \dots, x_m) = [F(x_m) - \gamma x_1] \hat{x}_1 + [F(x_1) - \gamma x_2] \hat{x}_2 + [F(x_2) - \gamma x_3] \hat{x}_3 + \dots + [F(x_{m-1}) - \gamma x_m] \hat{x}_m$, where $\hat{x}_i$ (for $i = 1, 2, \dots, m$) are the unit vectors in the direction of x_i axis. We calculated the inner product (Ξ) of the vector field with unit normal vector to the different facets. The normal unit vector is always considered to be directed into the hypercube. The outer facets of the hypercube are $x_1 = 0, x_1 = \frac{\alpha}{\gamma}, x_2 = 0, x_2 = \frac{\alpha}{\gamma}, \dots, x_m = 0$, and $x_m = \frac{\alpha}{\gamma}$. We computed the inner product between the vector field characterized by the system dynamics within the hypercube and the normal to the outer facets of the hypercube.

TABLE I: Inner product between the vector field and the normal to the outer facets of the hypercube.

Facet	Unit normal Vector, $\hat{n}$	Inner Product, $\Xi = \langle \vec{G}, \hat{n} \rangle$	Value of Ξ
$x_i = 0$	$\hat{x}_i$	$\frac{\alpha}{1 + \left(\frac{x_{i-1}}{k}\right)^n}$	+ve
$x_i = \frac{\alpha}{\gamma}$	$-\hat{x}_i$	$\alpha - \frac{\alpha}{1 + \left(\frac{x_{i-1}}{k}\right)^n}$	+ve
for $i = 1, 2, 3, \dots, m$. As it is a cyclic network, $x_0 = x_m$.			

The inner product between the vector field characterized by the system dynamics within the hypercube and the normal to the outer facets of the hypercube is given in Table I. For protein concentrations between 0 and $\frac{\alpha}{\gamma}$, the value of Ξ is always positive. This implies that the vector field flows through the facets into the hypercube. Thus, the vector field flowing through all the outer facets is always directed into the hypercube, making the hypercube a positively invariant box.

Construction of an m - dimensional Toroidal-like Manifold: Trajectories approach the unique steady state along the path defined by the eigenvector of the negative eigenvalue. We calculated the eigenvector for the real negative eigenvalue. Let the eigenvector be $[\hat{x}_1 \ \hat{x}_2 \ \hat{x}_3 \ \dots \ \hat{x}_m]^T$. So,

$$\begin{bmatrix} -\gamma & 0 & 0 & \dots & -\rho \\ -\rho & -\gamma & 0 & \dots & 0 \\ 0 & -\rho & -\gamma & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & -\gamma \end{bmatrix} \begin{bmatrix} \hat{x}_1 \\ \hat{x}_2 \\ \hat{x}_3 \\ \vdots \\ \hat{x}_m \end{bmatrix} = -(\gamma + \rho) \begin{bmatrix} \hat{x}_1 \\ \hat{x}_2 \\ \hat{x}_3 \\ \vdots \\ \hat{x}_m \end{bmatrix},$$

$$\implies \hat{x}_m = \hat{x}_1, \hat{x}_1 = \hat{x}_2, \dots, \hat{x}_{m-1} = \hat{x}_m,$$

$$\implies [\hat{x}_1 \ \hat{x}_2 \ \dots \ \hat{x}_m]^T = [1 \ 1 \ \dots \ 1]^T. \quad (2)$$

In the direction of the eigenvector $[\hat{x}_1 \ \hat{x}_2 \ \hat{x}_3 \ \dots \ \hat{x}_m]^T$, we cut out a hypervolume from the given hypercube such that the steady state and the trajectories approaching it along this singular path are eliminated. This hypervolume may be visualized as a cyclinder with a small radius $\varepsilon > 0$ along the eigenvector.

For some values of p , there is a possibility that some of these complex conjugate eigenvalues may have negative real part. So, we calculated the eigenvectors associated with such complex eigenvalues and removed the corresponding hypervolume along it. Let the eigenvector associated with such complex eigenvalues be $[\tilde{x}_1 \ \tilde{x}_2 \ \tilde{x}_3 \ \dots \ \tilde{x}_m]^T$. So,

$$\begin{bmatrix} -\gamma & 0 & 0 & \dots & -\rho \\ -\rho & -\gamma & 0 & \dots & 0 \\ 0 & -\rho & -\gamma & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & -\gamma \end{bmatrix} \begin{bmatrix} \tilde{x}_1 \\ \tilde{x}_2 \\ \tilde{x}_3 \\ \vdots \\ \tilde{x}_m \end{bmatrix} = -(\gamma + \rho f(p, m)) \begin{bmatrix} \tilde{x}_1 \\ \tilde{x}_2 \\ \tilde{x}_3 \\ \vdots \\ \tilde{x}_m \end{bmatrix},$$

where $f(p, m) = \left[\cos\left(\frac{2p+1}{m}\pi\right) + j \sin\left(\frac{2p+1}{m}\pi\right) \right]$ and

$$\begin{aligned} \tilde{x}_m &= \left[\cos\left(\frac{2p+1}{m}\pi\right) + j \sin\left(\frac{2p+1}{m}\pi\right) \right] \tilde{x}_1, \\ \tilde{x}_1 &= \left[\cos\left(\frac{2p+1}{m}\pi\right) + j \sin\left(\frac{2p+1}{m}\pi\right) \right] \tilde{x}_2, \\ &\vdots \\ \tilde{x}_{m-1} &= \left[\cos\left(\frac{2p+1}{m}\pi\right) + j \sin\left(\frac{2p+1}{m}\pi\right) \right] \tilde{x}_m. \end{aligned} \quad (3)$$

Using De Moivre's theorem on (3), we get the eigenvector

$$\begin{bmatrix} \tilde{x}_1 \\ \tilde{x}_2 \\ \tilde{x}_3 \\ \vdots \\ \tilde{x}_m \end{bmatrix} = \begin{bmatrix} 1 \\ \cos\left(\frac{(2p+1)(m-1)}{m}\pi\right) + j \sin\left(\frac{(2p+1)(m-1)}{m}\pi\right) \\ -\cos\left(\frac{(2p+1)(m-2)}{m}\pi\right) - j \sin\left(\frac{(2p+1)(m-2)}{m}\pi\right) \\ \vdots \\ -\cos\left(\frac{2p+1}{m}\pi\right) - j \sin\left(\frac{2p+1}{m}\pi\right) \end{bmatrix}.$$

Although complex eigenvectors are not elements of the real space, their real and imaginary components reside within it. We divided the above eigenvector into its real and imaginary

parts. These are the eigenvectors associated with a pair of complex conjugate eigenvalues,

$$v_{real} = \begin{bmatrix} 1 \\ \cos\left(\frac{(2p+1)(m-1)}{m}\pi\right) \\ -\cos\left(\frac{(2p+1)(m-2)}{m}\pi\right) \\ \vdots \\ -\cos\left(\frac{2p+1}{m}\pi\right) \end{bmatrix}, \quad v_{im} = \begin{bmatrix} 0 \\ \sin\left(\frac{(2p+1)(m-1)}{m}\pi\right) \\ -\sin\left(\frac{(2p+1)(m-2)}{m}\pi\right) \\ \vdots \\ -\sin\left(\frac{2p+1}{m}\pi\right) \end{bmatrix}.$$

Because some of these complex conjugate eigenvalues could have a negative real part, we removed the hypervolumes associated with the linear combination of v_{real} and v_{im} .

So, we obtained an m -dimensional toroidal-like manifold without the steady state. All the trajectories that enter this manifold stay inside because the hypercube is a positively invariant set and there are not trajectories that approach the steady state.

Continuous Map of a Cross-Section Onto Itself: We divided the hypercube into 2^m smaller hyperrectangles using the hyperplanes $x_i = x_0$ (for $i = 1, 2, 3, \dots, m$). Further, we computed the flow that traverses every facet within the smaller hyperrectangles.

TABLE II: Inner product between the vector field and the normal to the hyperplanes $x_i = x_0$ (for $i = 1, 2, 3, \dots, m$).

Hyper-plane	$\hat{n}$	$\Xi = \langle \vec{G}, \hat{n} \rangle$	Value of Ξ
$x_i = x_0$	$\hat{x}_i$	$\frac{\alpha}{1 + \left(\frac{x_{i-1}}{k}\right)^n} - \gamma x_0$	$\begin{cases} +ve, \text{ for } x_{i-1} < x_0, \\ -ve, \text{ for } x_{i-1} > x_0. \end{cases}$
for $i = 1, 2, 3, \dots, m$. As it is a cyclic network, $x_0 = x_m$.			

The flow of the vector field through the outer facets of the hypercube obtained in Table I, combined with the flow through the hyperplanes $x_i = x_0$ (for $i = 1, 2, 3, \dots, m$) obtained in Table II, is used to determine the effective flow from one smaller hyperrectangle to its adjacent ones. We considered one of the smaller hyperrectangles, $0 \leq x_1 \leq x_0$, $0 \leq x_2 \leq x_0$, ..., $0 \leq x_m \leq x_0$, as Box 1. From Table I, we have the flow of the vector field from the facets $x_i = 0$ (for $i = 1, 2, 3, \dots, m$) into Box 1. From Table II, we have the flow of the vector field from the hyperplanes $x_i = x_0$ (for $i = 1, 2, 3, \dots, m$) out of Box 1 and into the adjacent boxes. Similarly, we calculated the effective flow of the vector field for the rest of the boxes.

Upon inspection of the effective flow from one hyperrectangle to another, we observed that the vector field formed a closed path within certain hyperrectangles. This showed that a cross section of these smaller hyperrectangles mapped onto itself. By Brouwer's fixed point theorem, there exists a fixed point, which is a limit cycle for the system under consideration. The above construction also provides a rigorous bound on the region within the toroidal-like manifold where the limit cycle exists. ■

To summarize, we obtained the region in the hypercube where the limit cycle exists. This is done by constructing a toroidal-like manifold within a positively invariant set obtained by eliminating a hypervolume containing the steady state and the trajectories approaching it. Finally, we used Brouwer's Fixed Point theorem on a cross section of this manifold, which map onto itself, to show the existence of limit cycle.

III. CASE STUDY

The differential equations describing the dynamics of a 5-gene cyclic regulatory network are given in (4).

$$\frac{dx_i}{dt} = F(x_{i-1}) - \gamma x_i, \text{ for } i = 1, 2, 3, \dots, 5. \quad (4)$$

Here x_i is the concentration of the i -th protein with $0 \leq x_i \leq \frac{\alpha}{\gamma}$. α, γ, k , and n are positive constants. $F(p) = \frac{\alpha}{1 + (\frac{p}{k})^n}$ is the Hill function and n is the Hill coefficient. The eigenvalues of the Jacobian after the linearization of the system in (4) at the steady state $(x_0, x_0, x_0, x_0, x_0)$ is $\lambda_{1,2,\dots,5} = -(\gamma + \rho), -\gamma + \rho(-0.31 \pm j0.95), -\gamma + \rho(0.81 \pm j0.59)$. The 5-gene regulatory network has a negative real eigenvalue. Along with it, one of the complex conjugate eigenvalues also has a negative real part. Guided by Theorem 2.1, we illustrated the existence of the limit cycle by considering the case where one pair of the complex conjugate eigenvalues has a positive real part.

Construction of an invariant box: We calculated the flow of the vector field through the different facets of the hypercube $0 \leq x_1, x_2, \dots, x_5 \leq \frac{\alpha}{\gamma}$. This is calculated along the unit normal vector of the various facets directed into the hypercube. Using the value of Ξ , we determined the flow of the vector field through the various facets and checked if the hypercube constituted a positively invariant set.

The vector field described by the system dynamics in (4) is: $\vec{G}(x_1, x_2, x_3, x_4, x_5) = [F(x_5) - \gamma x_1] \hat{x}_1 + [F(x_1) - \gamma x_2] \hat{x}_2 + [F(x_2) - \gamma x_3] \hat{x}_3 + [F(x_3) - \gamma x_4] \hat{x}_4 + [F(x_4) - \gamma x_5] \hat{x}_5$, where $\hat{x}_i$ (for $i = 1, 2, \dots, 5$) are the unit vectors in the direction of x_i axis. The value of the inner product between the vector field and the normal to the facets directed into the hypercube is obtained from Table I by substituting $m = 5$. The values of Ξ are positive for every outer facet of the hypercube. This indicated that the flow of the vector field described by the system dynamics is always directed inward and stays inside the hypercube, making it a positively invariant set.

Construction of a 5-dimensional Toroidal-like Manifold: The existence of a unique steady state in the given hypercube indicated that certain trajectories approach the steady state along the singular path defined by the eigenvector of the negative eigenvalue. So, we calculated the eigenvector for the real negative eigenvalue. Let the eigenvector be $[\hat{x}_1 \ \hat{x}_2 \ \hat{x}_3 \ \hat{x}_4 \ \hat{x}_5]^T$. So, using (2) the eigenvector is found to be $[1 \ 1 \ 1 \ 1 \ 1]^T$. In the direction of this eigenvector we cut out a hypervolume from the given hypercube, such that the steady state and the trajectories approaching it along this singular path are eliminated.

The regulatory network also has a pair of complex conjugate eigenvalues with negative real part. So, we calculated the eigenvectors associated with these complex eigenvalues and removed a hypervolume along it. Using the relation in (3) and the De Moivre's theorem, we get the eigenvector as: $[\tilde{x}_1 \ \tilde{x}_2 \ \tilde{x}_3 \ \tilde{x}_4 \ \tilde{x}_5]^T = [1 \ 0.31 + j0.95 \ -0.81 + j0.59 \ -0.81 - j0.59 \ 0.31 - j0.95]^T$. We divide the above eigenvector into its real and imaginary parts. These are the eigenvectors associated complex eigenvalue with negative real part: $v_{real} = [1 \ 0.31 \ -0.81 \ -0.81 \ 0.31]^T$ and $v_{im} = [0 \ 0.95 \ 0.59 \ -0.59 \ -0.95]^T$. The complex conjugate eigenvalues produce conjugate eigenvectors. So, the v_{real} remains same for the complex conjugate pair. v_{im} for the pair of eigenvalues is a conjugate pair. However, the conjugate gives the same eigenvector but in the opposite direction. In the sense of removing a hypervolume, the v_{im} can be considered to be same. So, we also removed the hypervolume along the eigenvectors v_{real} and v_{im} .

So, we obtained a toroidal-like manifold with no steady state. All the trajectories are inside the torus-like structure (because the hypercube is a positively invariant set).

TABLE III: Direction of flow of vector field within the smaller boxes.

Box	Coordinates of the hyperrectangle	Direction of the flow of vector field
12	$0 \leq x_1 \leq x_0, x_0 \leq x_2 \leq \frac{\alpha}{\gamma}, 0 \leq x_3 \leq x_0, x_0 \leq x_4 \leq \frac{\alpha}{\gamma}, x_0 \leq x_5 \leq \frac{\alpha}{\gamma}$	Box 12 $\rightarrow$ Box 24
13	$0 \leq x_1 \leq x_0, x_0 \leq x_2 \leq \frac{\alpha}{\gamma}, x_0 \leq x_3 \leq \frac{\alpha}{\gamma}, 0 \leq x_4 \leq x_0, x_0 \leq x_5 \leq \frac{\alpha}{\gamma}$	Box 13 $\rightarrow$ Box 23
16	$x_0 \leq x_1 \leq \frac{\alpha}{\gamma}, 0 \leq x_2 \leq x_0, x_0 \leq x_3 \leq \frac{\alpha}{\gamma}, 0 \leq x_4 \leq x_0, x_0 \leq x_5 \leq \frac{\alpha}{\gamma}$	Box 16 $\rightarrow$ Box 21
17	$x_0 \leq x_1 \leq \frac{\alpha}{\gamma}, 0 \leq x_2 \leq x_0, x_0 \leq x_3 \leq \frac{\alpha}{\gamma}, x_0 \leq x_4 \leq \frac{\alpha}{\gamma}, 0 \leq x_5 \leq x_0$	Box 17 $\rightarrow$ Box 28
19	$x_0 \leq x_1 \leq \frac{\alpha}{\gamma}, x_0 \leq x_2 \leq \frac{\alpha}{\gamma}, 0 \leq x_3 \leq x_0, x_0 \leq x_4 \leq \frac{\alpha}{\gamma}, 0 \leq x_5 \leq x_0$	Box 19 $\rightarrow$ Box 26
21	$0 \leq x_1 \leq x_0, 0 \leq x_2 \leq x_0, x_0 \leq x_3 \leq \frac{\alpha}{\gamma}, 0 \leq x_4 \leq x_0, x_0 \leq x_5 \leq \frac{\alpha}{\gamma}$	Box 21 $\rightarrow$ Box 13
23	$0 \leq x_1 \leq x_0, x_0 \leq x_2 \leq \frac{\alpha}{\gamma}, 0 \leq x_3 \leq x_0, 0 \leq x_4 \leq x_0, x_0 \leq x_5 \leq \frac{\alpha}{\gamma}$	Box 23 $\rightarrow$ Box 12
24	$0 \leq x_1 \leq x_0, x_0 \leq x_2 \leq \frac{\alpha}{\gamma}, 0 \leq x_3 \leq x_0, x_0 \leq x_4 \leq \frac{\alpha}{\gamma}, 0 \leq x_5 \leq x_0$	Box 24 $\rightarrow$ Box 19
26	$x_0 \leq x_1 \leq \frac{\alpha}{\gamma}, 0 \leq x_2 \leq x_0, 0 \leq x_3 \leq x_0, x_0 \leq x_4 \leq \frac{\alpha}{\gamma}, 0 \leq x_5 \leq x_0$	Box 26 $\rightarrow$ Box 17
28	$x_0 \leq x_1 \leq \frac{\alpha}{\gamma}, 0 \leq x_2 \leq x_0, x_0 \leq x_3 \leq \frac{\alpha}{\gamma}, 0 \leq x_4 \leq x_0, 0 \leq x_5 \leq x_0$	Box 28 $\rightarrow$ Box 16
Complete table in Supplementary (Part - I).		

Continuous Map of a Cross-Section Onto Itself: The hypercube is divided into 2^5 smaller hyperrectangles by using the hyperplanes $x_1 = x_0, x_2 = x_0, \dots$, and $x_5 = x_0$. We obtained the value of Ξ between the vector field describing the system dynamics and the unit normal vector through these hyperplanes from Table II by substituting $m = 5$.

Using the values of Ξ between (a) the vector field and the normal to the outer facets of the hypercube and (b) the vector field and the normal to the hyperplanes $x_i = x_0$ (for $i = 1, 2, \dots, 5$), we determined the direction of flow of the vector field within the smaller boxes. This is summarized in Table

III (For complete table refer to Supplementary (Part - I) available at: [GitHub](#)). Upon inspection of Table III, we find that the vector field formed a continuous map of a cross section onto itself. The continuous map is as follows: $12 \rightarrow 24 \rightarrow 19 \rightarrow 26 \rightarrow 17 \rightarrow 28 \rightarrow 16 \rightarrow 21 \rightarrow 13 \rightarrow 23 \rightarrow 12$. We obtained a toroidal-like manifold (which is a positively invariant set) and a cross section of this structure formed a continuous map onto itself. So, by Brouwer's Fixed Point theorem, there exists a limit cycle.

IV. LOCALIZATION OF LIMIT CYCLE

We extended the core idea of the Brouwer's Fixed Point theorem to an interval-based Reachability Analysis [22] to localize the limit cycle trajectory. This localization is the natural next step in rigorously refining the region of existence of the limit cycle. This method is based on rigorously solving the network dynamics using Interval Analysis. Starting from a set of initial conditions, a hypercube is obtained that rigorously contains all trajectories. The sets that mapped onto themselves under the vector field of the system dynamics, rigorously computed using Reachability Analysis, contain a limit cycle. This is a direct application of the Brouwer's Fixed Point theorem on refinement of the existence region.

The methodology involves discretizing an initial hypercube into smaller hyperrectangles, evolving the dynamics from these smaller hyperrectangles through Reachability Analysis (flowpipes), and examining the 'return map'. Each region is categorized as: (i) Does not contain the limit cycle, (ii) May contain the limit cycle, or (iii) Contains the limit cycle. This classification fundamentally depends on a geometric subset test to ascertain whether the rigorously computed return set X_t , is entirely encompassed within the original set X_0 .

This interval-based Reachability Analysis approach thus serves as an implementation of Brouwer's Fixed Point theorem's topological arguments using validated numerical techniques. While the theorem guarantees the presence of a fixed point and thus a periodic orbit, Reachability Analysis refines this conclusion by localizing the region where the limit cycle trajectory lies. In this approach, the geometric confinement test extends the theoretical proof by connecting the existence result to a rigorous characterization of the system's oscillatory dynamics.

The existence proof for the three-gene cyclic regulatory network within an invariant cube $0 \leq x_1, x_2, x_3 \leq \frac{\alpha}{\gamma}$, when divided into 2^3 smaller boxes by the surfaces $x_i = x_0$ (for $i = 1, 2, 3$), gives us six cross-sections of the smaller boxes that map onto themselves. We then illustrated the interval-based Reachability Analysis to localize periodic orbit through one of these cross-sections that map onto themselves in the three-gene cyclic regulatory network. The framework successfully divided the cross-section according to the underlying dynamical behaviour of the three-gene cyclic regulatory network. The methodology, based on partitioning the cross-section and analyzing the flowpipe of each partition, showed several findings, which are visually summarized in

Fig. 1 (For complete details about the methodology refer to Supplementary (Part - II) available at: [GitHub](#)).




Fig. 1: Projection of the cross-section, that maps onto itself, showing the geometric classification results. The black box contains the steady-state point and was excluded from the reachability analysis. The blue boxes is the region that does not contain the limit cycle. The cluster of yellow boxes signify the region that may contain the limit cycle. The green boxes are classified as the boxes that failed to return a solution for reachability analysis due to solver crash or blow up in the bounds due to overapproximation. The system parameters used in the simulation are: $\alpha = 26.26$, $\gamma = 0.92$, $k = 4$, $n = 5$.




Fig. 2: The numerical solution of the three-gene cyclic regulatory network with initial condition $(19, 3.5, 5.361)$ passes through the region classified as "May contain the limit cycle" in Fig. 1. The system parameters used in the simulation are: $\alpha = 26.26$, $\gamma = 0.92$, $k = 4$, $n = 5$. x, y, z axes represent the concentration of i -th (for $i = 1, 2, 3$) protein respectively.

The analysis classified the vast majority of the partitioned region of this cross-section as not containing a periodic orbit. The black box in Figure 1, contains the unstable steady state (x_0, x_0, x_0) (the value of x_0 for this simulation is 5.361) of the system and is excluded from the Reachability Anal-

ysis. A distinct cluster of initial sets, marked in yellow in Figure 1, was classified as “May contain the limit cycle”. A cluster of green boxes around the box containing the steady state is classified as “Failed boxes”. This is because of the failure of the reachability analysis to return a solution. This can be attributed to solver crash or blow up in the bounds due to overapproximation. Further in Fig. 2 the numerical solution of the three gene cyclic regulatory network model is plotted along with one of the cross-section that maps onto itself. Fig. 3 provides a magnified view of the trajectory passing through the region classified by the method as “May contain the limit cycle”. This finding validates the method’s efficacy in localizing the region in which the limit cycle exists.




Fig. 3: A magnified view of Fig. 2 showing the intersection of the numerical solution with the boxes classified as “May contain the limit cycle”.

V. CONCLUSION

We proved the existence of a limit cycle in a class of cyclic gene regulatory networks and provided a rigorous localization of this region of existence. The existence proof, based on Brouwer’s Fixed Point theorem, is an alternative and elementary proof for this class of systems, with the benefit of an enhanced geometric visualisation. This rigorous localization relies on an interval-based Reachability Analysis, which, in conjunction with Brouwer’s Fixed Point theorem, can provide a significantly tighter enclosure of the limit cycle trajectory.

For the existence proof, we constructed a positively invariant set for the network dynamics by exploiting the inherent boundedness and by creating a toroidal-like manifold region that eliminates hypervolume consisting of the unstable steady state and its associated stable manifolds. We found that a cross section of this manifold continuously maps onto itself under the system dynamics, showing via Brouwer’s Fixed Point theorem, the existence of a limit cycle. These results are illustrated using a 5-dimensional example. After establishing the existence of a limit cycle in cyclic gene regulatory networks, we employed interval-based Reachability Analysis to localize the region of its existence across the cross-sections that map onto themselves. The Reachability

Analysis was demonstrated using a 3-dimensional example, offering a visual representation of the categorized regions.

REFERENCES

- [1] H. K. Khalil, *Nonlinear control*, Vol. 406, New York: Pearson, 2015.
- [2] U. Alon, *An introduction to systems biology: design principles of biological circuits*, Chapman and Hall/CRC, 2006.
- [3] B. C. Goodwin, “Temporal organization in cells. A dynamic theory of cellular control processes”, *Temporal organization in cells. A dynamic theory of cellular control processes*, 1963.
- [4] B. C. Goodwin, “Oscillatory behavior in enzymatic control processes”, *Advances in enzyme regulation* 3, pp. 425-437, 1965.
- [5] J. S. Griffith, “Mathematics of cellular control processes I. Negative feedback to one gene”, *Journal of theoretical biology* 20.2, pp. 202-208, 1968.
- [6] R. Thomas and R. d’Ari, *Biological Feedback*, CRC Press, 1990.
- [7] M. Elowitz and S. Leibler, “A synthetic oscillatory network of transcriptional regulators”, *Nature* 403, pp. 335-338, 2000.
- [8] R. Thomas and M. Kaufman, “Multistationarity, the basis of cell differentiation and memory. ii. logical analysis of regulatory networks in terms of feedback circuits”, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 2001.
- [9] J. Tyson, “On the existence of oscillatory solutions in negative feedback cellular control processes”, *Journal of Mathematical Biology*, 1975.
- [10] S. Mohanty, A. Mohit, M. Aggarwal, and S. Sen, “An Elementary Proof for the Existence of Limit Cycle in Three Gene Cyclic Regulatory Network”, *2023 Ninth Indian Control Conference (ICC)*, Visakhapatnam, India, pp. 401-406, 2023.
- [11] S. Hastings, J. Tyson, and D. Webster, “Existence of periodic solutions for negative feedback cellular control systems”, *Journal of Differential Equations*, 1977.
- [12] H. El Samad, D. Del Vecchio, and M. Khammash, “Repressilators and promotilators: loop dynamics in synthetic gene networks”, *Proceedings of the 2005, American Control Conference*, vol. 6, pp. 4405-4410, 2005.
- [13] R. A. Smith, “Orbital stability for ordinary differential equations”, *Journal of differential equations*, Volume 69, Issue 2, pp. 265-287, 1987.
- [14] J. Mallet-Paret and H. Smith, “The Poincaré-Bendixson theorem for monotone cyclic feedback systems”, *Journal of Dynamics and Differential Equations* Volume 2, Issue 4, pp. 367-421, 1990.
- [15] D. Del Vecchio and R. M. Murray, *Biomolecular Feedback Systems*, Princeton University Press, Princeton, NJ, 2015.
- [16] Y. Hori, T. -H. Kim, and S. Hara, “Graphical and analytic criteria for the existence of protein level oscillations in cyclic gene regulatory networks”, *Proceedings of the 48th IEEE Conference on Decision and Control (CDC) held jointly with 2009 28th Chinese Control Conference*, pp. 3521-3526, 2009.
- [17] Y. Hori, S. Hara, and T. -H. Kim, “Periodic oscillations in cyclic repressor networks: Analytic existence criteria with biological insight”, *2010 IEEE International Conference on Control Applications*, pp. 824-829, 2010.
- [18] Y. Hori, T. -H. Kim and S. Hara, “Existence criteria of periodic oscillations in cyclic gene regulatory networks”, *Automatica*, Volume 47, Issue 6, pp. 1203-1209, 2011.
- [19] R. Katz, G. Giordano, and M. Margaliot, “Convergence to periodic orbits in 3-dimensional strongly 2-cooperative systems”, *arXiv preprint arXiv:2407.00461* (2024).
- [20] R. Katz, G. Giordano, and M. Margaliot, “Instability of equilibrium and convergence to periodic orbits in strongly 2-cooperative systems”, *Journal of Differential Equations*, Volume 444, Art. no. 133651, 2025.
- [21] L. A. Sanchez, “Global asymptotic stability of the Goodwin system with repression”, *Nonlinear Analysis: Real World Applications*, Volume 10, Issue 4, pp. 2151-2156, 2009b.
- [22] S. Bogomolov, M. Forets, G. Frehse, K. Potomkin, and C. Schilling, “JuliaReach: a toolbox for set-based reachability”, *Proceedings of the 22nd ACM International Conference on Hybrid Systems: Computation and Control*, pp. 39-44, 2019.