

Discriminating Seagrass From Green Macroalgae in European Intertidal areas using high resolution multispectral drone imagery

Simon Oiry^{a,*}, Bede Ffinian Rowe Davies^a, Pierre Gernez^a, Ana I. Sousa^b, Philippe Rosa^a, Maria Laura Zoffoli^c, Guillaume Brunier^d, Laurent Barillé^a

^a*Nantes University,*

^b*Aveiro University,*

^c*CNR ISMAR,*

^d*JeSaisPas,*

Abstract

In September 2021, a significant jump in seismic activity on the island of La Palma (Canary Islands, Spain) signaled the start of a volcanic crisis that still continues at the time of writing. Earthquake data is continually collected and published by the Instituto Geográfico Nacional (IGN). ...

Keywords: Drone, Remote Sensing

1. Introduction

Coastal areas are vital hotspots for marine biodiversity, with intertidal seagrass meadows playing a crucial role at the interface between land and ocean [1]. These meadows offer a myriad of ecosystem services to humanity, including limitation of coastline erosion, reducing the risk of eutrophication, carbon sequestration, and oxygen production. They serve as vital habitats for a diverse array of marine and terrestrial species, providing living, breeding, and feeding grounds [2, ; 3, ; 4]. Due to their proximity to human activities, seagrass meadows are directly exposed to and impacted by anthropogenic pressures. Global regression and fragmentation are currently observed due to diseases, disasters, coastal urbanization, sea reclamation, as well as fishing activities, dredging, sea level rise, coastal erosion, competition with alien species, and reduction in water quality [5, ; 6, ; 7, ; 8, ; 9]. While improvements in water quality have been recently reported in European sites, allowing an overall recovery of seagrass

*Corresponding author

Email address: oirysimon@gmail.com (Simon Oiry)

ecosystems at the local scale, many other coastal waters worldwide are still subjected to strong eutrophication processes [10, ; 11]. Coastal eutrophication has been associated to anomalous accumulation of green macroalgae, the so-called green tides. Green tides produce shade and suffocation over seagrass individuals, thus threatening the health of seagrass ecosystems [? , ; 12].

The importance of seagrass meadows and the variety of ecosystem services they provide have led to the enhancement of global and regional monitoring programs for systematically surveying different Essential Oceanic Variable [13] as seagrass coverage and composition; as well as Essential Biodiversity Variable [14] such as seagrass taxonomic diversity; species distribution, population abundance, and phenology. Monitoring programs also prioritize the identification of threats to these ecosystems, particularly during early stages, to facilitate effective mitigation actions. Traditionally, these ecological parameters have been quantified through in situ measurements, although this approach faces several constraints over intertidal zones. Intertidal meadows are only partially exposed during low tide and can be situated in difficult-to-reach mudflats, potentially leading to inaccurate and limited estimations with conventional sampling techniques [15]. However, satellite data have been proven effective in complementing in situ surveys, allowing for the rapid and consistent retrieval of EOVS over extensive seagrass meadows. [11, ; 16, ; 17, ; 18]

Satellite remote sensing offers the advantage of acquiring large-scale data in real-time but presents its inherent challenges. Free access satellite data (e.g., Sentinel-2 and Landsat8/9) provide relatively low spatial resolution data (10 - 30 m) across a limited number of spectral bands. These characteristics can be a limitation to accurately discriminating seagrass from others co-existing macrophytes over the meadow. Chlorophyceae (Green Algae) and marine Magnoliopsida (Seagrass) share the same pigment composition [19, ; 20]. As a result, their respective spectral signatures can be considered similar by a non-expert observer [21, ; 22]. Recently, using a hyperspectral library, [21] demonstrated that the spectral resolution of Sentinel-2, might be enough for the discrimination between Magnoliopsida and Chlorophyceae. However, green tide events occur at small spatial scales that are not observable using satellite imagery [23], especially during the initial stage of the event.

Remote sensing drone acquisitions are presented as a tool that can potentially fill gaps left by satellite and in situ data. Drone can cover large expanses while recording imagery at significantly higher spatial resolutions than satellite (pixel size from cm to mm) and still capturing data at multi-spectral resolution [24, ; 25]. The versatility of drones allows for their application across a diverse thematic range, from coastal zone management [26, ; 27, ; 28] to mapping the spatial distribution of species [29, ; 30, ; 31, ; 32, ; 33]. However, when applied to coastal habitat mapping, many studies showcase their findings with study case limited to a single flight, restricting the generalizability of their application to other sites [32, ; 34, ; 35, ; 33]. This study aimed to analyze the potential of a drone equipped with a multispectral sensor for mapping inter-

tidal macrophytes, with a particular focus on discriminating Magnoliopsida and Chlorophyceae. Ten drone flights were performed over soft-bottom intertidal areas along two European countries (France and Portugal), covering a wide range of habitats, from monospecific seagrass meadows to meadows mixed with green or red algae. A deep learning algorithm was trained and validated for macrophyte discrimination, emphasizing applicability across diverse sites without a loss of accuracy in predictions.

References

- [1] R. K. Unsworth, L. C. Cullen-Unsworth, B. L. Jones, R. J. Lilley, The planetary role of seagrass conservation, *Science* 377 (6606) (2022) 609–613.
- [2] R. C. Gardner, C. Finlayson, Global wetland outlook: state of the world’s wetlands and their services to people, 2018.
- [3] M. L. Zoffoli, P. Gernez, S. Oiry, L. Godet, S. Dalloyau, B. F. R. Davies, L. Barillé, Remote sensing in seagrass ecology: coupled dynamics between migratory herbivorous birds and intertidal meadows observed by satellite during four decades, *Remote Sensing in Ecology and Conservation* (12 2022). doi:10.1002/rse2.319.
- [4] E. Jankowska, L. N. Michel, G. Lepoint, M. Włodarska-Kowalczyk, Stabilizing effects of seagrass meadows on coastal water benthic food webs, *Journal of Experimental Marine Biology and Ecology* 510 (2019) 54–63.
- [5] H. M. Nguyen, P. J. Ralph, L. Marín-Guirao, M. Pernice, G. Procaccini, Seagrasses in an era of ocean warming: a review, *Biological Reviews* 96 (5) (2021) 2009–2030.
- [6] L. M. Soissons, E. P. Haanstra, M. M. Van Katwijk, R. Asmus, I. Auby, L. Barillé, F. G. Brun, P. G. Cardoso, N. Desroy, J. Fournier, et al., Latitudinal patterns in european seagrass carbon reserves: influence of seasonal fluctuations versus short-term stress and disturbance events, *Frontiers in Plant Science* 9 (2018) 88.
- [7] R. J. Orth, T. J. Carruthers, W. C. Dennison, C. M. Duarte, J. W. Fourqurean, K. L. Heck, A. R. Hughes, G. A. Kendrick, W. J. Kenworthy, S. Olyarnik, et al., A global crisis for seagrass ecosystems, *Bioscience* 56 (12) (2006) 987–996.
- [8] H. Lin, T. Sun, Y. Zhou, R. Gu, X. Zhang, W. Yang, Which genes in a typical intertidal seagrass (*zostera japonica*) indicate copper-, lead-, and cadmium pollution?, *Frontiers in Plant Science* 9 (2018) 1545.
- [9] J. E. Duffy, L. Benedetti-Cecchi, J. Trinanes, F. E. Muller-Karger, R. Ambo-Rappe, C. Boström, A. H. Buschmann, J. Byrnes, R. G. Coles, J. Creed, et al., Toward a coordinated global observing system for seagrasses and marine macroalgae, *Frontiers in Marine Science* 6 (2019) 317.

- [10] C. B. de Los Santos, D. Krause-Jensen, T. Alcoverro, N. Marbà, C. M. Duarte, M. M. Van Katwijk, M. Pérez, J. Romero, J. L. Sánchez-Lizaso, G. Roca, et al., Recent trend reversal for declining european seagrass meadows, *Nature communications* 10 (1) (2019) 3356.
- [11] M. L. Zoffoli, P. Gernez, L. Godet, S. Peters, S. Oiry, L. Barillé, [Decadal increase in the ecological status of a north-atlantic intertidal seagrass meadow observed with multi-mission satellite time-series](#), *Ecological Indicators* 130 (2021) 108033. doi:10.1016/j.ecolind.2021.108033.
URL <https://linkinghub.elsevier.com/retrieve/pii/S1470160X21006981>
- [12] Z. Wang, Z. Fang, J. Liang, X. Song, Assessment of global habitat suitability and risk of ocean green tides, *Harmful Algae* 119 (2022) 102324.
- [13] P. Miloslavich, N. J. Bax, S. E. Simmons, E. Klein, W. Appeltans, O. Aburto-Oropeza, M. A. Garcia, S. D. Batten, L. Benedetti-Cecchi, D. M. Checkley, S. Chiba, J. E. Duffy, D. C. Dunn, A. Fischer, J. Gunn, R. Kudela, F. Marsac, F. E. Muller-Karger, D. Obura, Y. J. Shin, [Essential ocean variables for global sustained observations of biodiversity and ecosystem changes](#), *Global Change Biology* 24 (2018) 2416–2433. doi:10.1111/GCB.14108.
URL <https://onlinelibrary.wiley.com/doi/full/10.1111/gcb.14108><https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.14108><https://onlinelibrary.wiley.com/doi/10.1111/gcb.14108>
- [14] H. M. Pereira, S. Ferrier, M. Walters, G. N. Geller, R. H. Jongman, R. J. Scholes, M. W. Bruford, N. Brummitt, S. H. Butchart, A. Cardoso, et al., Essential biodiversity variables, *Science* 339 (6117) (2013) 277–278.
- [15] W. Nijland, L. Reshitnyk, E. Rubidge, Satellite remote sensing of canopy-forming kelp on a complex coastline: a novel procedure using the landsat image archive, *Remote Sensing of Environment* 220 (2019) 41–50.
- [16] S. Xu, S. Xu, Y. Zhou, S. Yue, X. Zhang, R. Gu, Y. Zhang, Y. Qiao, M. Liu, Long-term changes in the unique and largest seagrass meadows in the bohai sea (china) using satellite (1974–2019) and sonar data: Implication for conservation and restoration, *Remote Sensing* 13 (5) (2021) 856.
- [17] D. Traganos, P. Reinartz, Mapping mediterranean seagrasses with sentinel-2 imagery, *Marine Pollution Bulletin* 134 (2018) 197–209. doi:10.1016/j.marpolbul.2017.06.075.
- [18] M. M. Coffey, D. D. Graybill, P. J. Whitman, B. A. Schaeffer, W. B. Salls, R. C. Zimmerman, V. Hill, M. C. Lebrasse, J. Li, D. J. Keith, et al., Providing a framework for seagrass mapping in united states coastal ecosystems using high spatial resolution satellite imagery, *Journal of Environmental Management* 337 (2023) 117669.

- [19] P. Ralph, S. Polk, K. Moore, R. Orth, W. Smith Jr, Operation of the xanthophyll cycle in the seagrass *zostera marina* in response to variable irradiance, *Journal of Experimental Marine Biology and Ecology* 271 (2) (2002) 189–207.
- [20] F. Douay, C. Verpoorter, G. Duong, N. Spilmont, F. Gevaert, [New hyperspectral procedure to discriminate intertidal macroalgae](#), *Remote Sensing* 14 (2) (2022). doi:10.3390/rs14020346.
URL <https://www.mdpi.com/2072-4292/14/2/346>
- [21] B. F. R. Davies, P. Gernez, A. Geraud, S. Oiry, P. Rosa, M. L. Zofoli, L. Barillé, [Multi- and hyperspectral classification of soft-bottom intertidal vegetation using a spectral library for coastal biodiversity remote sensing](#), *Remote Sensing of Environment* 290 (2023) 113554. doi:10.1016/j.rse.2023.113554.
URL <https://www.sciencedirect.com/science/article/pii/S0034425723001050>
- [22] A. Bannari, T. S. Ali, A. Abahussain, The capabilities of sentinel-msi (2a/2b) and landsat-oli (8/9) in seagrass and algae species differentiation using spectral reflectance, *Ocean Science* 18 (2) (2022) 361–388.
- [23] F. Tuya, H. Hernandez-Zerpa, F. Espino, R. Haroun, Drastic decadal decline of the seagrass *cymodocea nodosa* at gran canaria (eastern atlantic): interactions with the green algae *caulerpa prolifera*, *Aquatic Botany* 105 (2013) 1–6.
- [24] I. Fairley, B. J. Williamson, J. McIlvenny, N. King, I. Masters, M. Lewis, S. Neill, D. Glasby, D. Coles, B. Powell, et al., Drone-based large-scale particle image velocimetry applied to tidal stream energy resource assessment, *Renewable Energy* 196 (2022) 839–855.
- [25] J. Oh, D.-j. Kim, H. Lee, Use of a drone for mapping and time series image acquisition of tidal zones, *Journal of the Korean Institute of Intelligent Systems* 27 (2) (2017) 119–125.
- [26] R. Adade, A. M. Aibinu, B. Ekumah, J. Asaana, Unmanned aerial vehicle (uav) applications in coastal zone management—a review, *Environmental Monitoring and Assessment* 193 (2021) 1–12.
- [27] E. Casella, J. Drechsel, C. Winter, M. Benninghoff, A. Rovere, Accuracy of sand beach topography surveying by drones and photogrammetry, *Geo-Marine Letters* 40 (2020) 255–268.
- [28] D. B. Angnuureng, K. Brempong, P. Jayson-Quashigah, O. Dada, S. Akuoko, J. Frimpomaa, P. Mattah, R. Almar, Satellite, drone and video camera multi-platform monitoring of coastal erosion at an engineered pocket beach: A showcase for coastal management at elmina bay, ghana (west africa), *Regional Studies in Marine Science* 53 (2022) 102437.

- [29] K. E. Joyce, K. C. Fickas, M. Kalamandeen, The unique value proposition for using drones to map coastal ecosystems, *Cambridge Prisms: Coastal Futures* 1 (2023) e6.
- [30] K. Tallam, N. Nguyen, J. Ventura, A. Fricker, S. Calhoun, J. O’Leary, M. Fitzgibbons, I. Robbins, R. K. Walter, Application of deep learning for classification of intertidal eelgrass from drone-acquired imagery, *Remote Sensing* 15 (9) (2023) 2321.
- [31] M. Roca, M. B. Dunbar, A. Román, I. Caballero, M. L. Zoffoli, P. Gernez, G. Navarro, Monitoring the marine invasive alien species *rugulopteryx okamurae* using unmanned aerial vehicles and satellites, *Frontiers in Marine Science* 9 (10 2022). doi:[10.3389/fmars.2022.1004012](https://doi.org/10.3389/fmars.2022.1004012).
- [32] A. Román, A. Tovar-Sánchez, I. Olivé, G. Navarro, Using a uav-mounted multispectral camera for the monitoring of marine macrophytes, *Frontiers in Marine Science* (2021) 1225.
- [33] G. Brunier, S. Oiry, Y. Gruet, S. F. Dubois, L. Barillé, [Topographic analysis of intertidal polychaete reefs \(sabellaria alveolata\) at a very high spatial resolution](#), *Remote Sensing* 2022, Vol. 14, Page 307 14 (2022) 307. doi:[10.3390/RS14020307](https://doi.org/10.3390/RS14020307).
URL <https://www.mdpi.com/2072-4292/14/2/307>
- [34] A. Collin, S. Dubois, D. James, T. Houet, Improving intertidal reef mapping using uav surface, red edge, and near-infrared data, *Drones* 3 (3) (2019) 67.
- [35] T. Rossiter, T. Furey, T. McCarthy, D. B. Stengel, Uav-mounted hyperspectral mapping of intertidal macroalgae, *Estuarine, Coastal and Shelf Science* 242 (2020) 106789.