

DISCOV-Gracilaria Paper

Simon Oiry¹ Bede Ffinian Rowe Davies¹ Pierre Gernez¹
Laurent Barillé¹

2024-11-17

To be Written

1

1 Introduction

Coastal ecosystems are among the most dynamic and productive environments on Earth, providing ecosystem services and supporting biodiversity (Barbier et al., 2011; Unsworth et al., 2022; Watanabe et al., 2018). These ecosystems, spanning mangroves, salt marshes, seagrass meadows, and rocky intertidal zones, play a pivotal role in carbon sequestration, nutrient cycling, and shoreline stabilization (Liquete et al., 2013; Mehvar et al., 2018). They also serve as habitats for numerous species, many of which are commercially or ecologically significant (De Valck et al., 2023; Seitz et al., 2014). Coastal areas are densely populated, with billions of people globally depending on their resources for livelihoods, fisheries, and tourism (Mukherjee et al., 2023; Small and Nicholls, 2003). However, coastal ecosystems face mounting pressures from human activities such as land reclamation, pollution, and overfishing, compounded by the impacts of climate change (Hall-Spencer and Harvey, 2019; Lu et al., 2018). Sea level rise, ocean acidification, and increasing storm intensity further exacerbate the vulnerability of these systems, threatening their resilience and the services they provide (He and Silliman, 2019).

One of the significant threats to coastal ecosystems is biological invasions by non-native species, which can disrupt native biodiversity and alter ecosystem functions (Capdevila et al., 2019; Krueger-Hadfield, 2018; Liu et al., 2020). *Gracilaria vermiculophylla*, an invasive red macroalga native to the northwest Pacific, exemplifies this issue. Over the last century, this species has spread extensively across temperate estuaries in North America, Europe, and other regions, facilitated by aquaculture and maritime activities (Krueger-Hadfield et al., 2017;

¹Institut des Substances et Organismes de la Mer, ISOMer, Nantes Université, UR 2160, F-44000 Nantes, France

Rueness, 2005; Weinberger et al., 2008). Its success as an invader stems from its tolerance to a wide range of environmental stressors, including temperature (Sotka et al., 2018), salinity (Weinberger et al., 2008), and nutrient variability (Abreu et al., 2011), as well as its ability to establish in soft sediment habitats traditionally devoid of macroalgae (Ramus et al., 2017). While *G. vermiculophylla* can provide some ecosystem services, such as habitat for invertebrates and juvenile fish, it often outcompetes native vegetation, alters sediment composition (Nyberg et al., 2009), and disrupts trophic interactions (Ginneken et al., 2018). In regions like the Baltic Sea and the eastern United States, it has been documented to negatively affect native fucoids and seagrasses (Thomsen et al., 2013; Van Katwijk, 2003). These impacts underscore the importance of monitoring and managing the spread of *G. vermiculophylla*, particularly as climate change and anthropogenic pressures continue to facilitate biological invasions.

Remote sensing has revolutionized our ability to monitor and manage ecosystems, offering efficient and scalable methods for detecting environmental changes over large areas (Davies et al., 2024a, 2024b; Zoffoli et al., 2021). Among these technologies, drone-based remote sensing has emerged as a particularly promising tool for studying coastal environments (Román et al., 2024, 2023). Equipped with high-resolution cameras and multispectral or hyperspectral sensors, drones can capture fine-scale spatial and spectral data, enabling researchers to identify and map vegetation, detect stress in plants, and monitor changes over time [Román et al. (2021) ; Oiry et al. 2024]. Unlike traditional satellite imagery, drones provide the flexibility to operate in overcast conditions, achieve higher spatial resolution, and target specific areas of interest. For invasive species like *G. vermiculophylla*, drones equipped with multispectral sensors can differentiate it from native vegetation based on its unique spectral reflectance characteristics (Davies et al. 2025). This capability not only enhances detection accuracy but also reduces the time and labor associated with traditional field surveys. As the cost of drone technology continues to decrease and advancements in machine learning facilitate data analysis, drone-based remote sensing is becoming increasingly accessible and impactful for ecological research and management.

In this study, we aim to harness the potential of drone-based multispectral remote sensing to map *Gracilaria vermiculophylla* in intertidal zones. Bla bla what are we going to do ? bla bla .

- Abreu, M.H., Pereira, R., Buschmann, A., Sousa-Pinto, I., Yarish, C., 2011. Nitrogen uptake responses of *Gracilaria vermiculophylla* (ohmi) papenfuss under combined and single addition of nitrate and ammonium. *Journal of Experimental Marine Biology and Ecology* 407, 190–199.
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C., Silliman, B.R., 2011. The value of estuarine and coastal ecosystem services. *Ecological monographs* 81, 169–193.
- Capdevila, P., Hereu, B., Salguero-Gómez, R., Rovira, G. la, Medrano, A., Cebrian, E., Garrabou, J., Kersting, D.K., Linares, C., 2019. Warming impacts on early life stages increase the vulnerability and delay the population recovery of a long-lived habitat-forming macroalga. *Journal of Ecology* 107, 1129–1140.

- Davies, B.F.R., Oiry, S., Rosa, P., Zoffoli, M.L., Sousa, A.I., Thomas, O.R., Smale, D.A., Austen, M.C., Biermann, L., Attrill, M.J., others, 2024b. A sentinel watching over intertidal seagrass phenology across western europe and north africa. *Communications Earth & Environment* 5, 382.
- Davies, B.F.R., Oiry, S., Rosa, P., Zoffoli, M.L., Sousa, A.I., Thomas, O.R., Smale, D.A., Austen, M.C., Biermann, L., Attrill, M.J., others, 2024a. Intertidal seagrass extent from sentinel-2 time-series show distinct trajectories in western europe. *Remote Sensing of Environment* 312, 114340.
- De Valck, J., Jarvis, D., Coggan, A., Schirru, E., Pert, P., Graham, V., Newlands, M., 2023. Valuing ecosystem services in complex coastal settings: An extended ecosystem accounting framework for improved decision-making. *Marine Policy* 155, 105761.
- Ginneken, V. van, Vries, E. de, others, 2018. The global dispersal of the non-endemic invasive red alga *gracilaria vermiculophylla* in the ecosystems of the euro-asia coastal waters including the wadden sea unesco world heritage coastal area: Awful or awesome? *Oceanography & Fisheries Open Access Journal* 8, 4–26.
- Hall-Spencer, J.M., Harvey, B.P., 2019. Ocean acidification impacts on coastal ecosystem services due to habitat degradation. *Emerging Topics in Life Sciences* 3, 197–206.
- He, Q., Silliman, B.R., 2019. Climate change, human impacts, and coastal ecosystems in the anthropocene. *Current Biology* 29, R1021–R1035.
- Krueger-Hadfield, S., 2018. Everywhere you look, everywhere you go, there’s an estuary invaded by the red seaweed *gracilaria vermiculophylla* (ohmi) papenfuss, 1967. *BioInvasions Records* 7.
- Krueger-Hadfield, S.A., Kollars, N.M., Strand, A.E., Byers, J.E., Shainker, S.J., Terada, R., Greig, T.W., Hammann, M., Murray, D.C., Weinberger, F., others, 2017. Genetic identification of source and likely vector of a widespread marine invader. *Ecology and evolution* 7, 4432–4447.
- Liquete, C., Piroddi, C., Drakou, E.G., Gurney, L., Katsanevakis, S., Charef, A., Egoh, B., 2013. Current status and future prospects for the assessment of marine and coastal ecosystem services: A systematic review. *PloS one* 8, e67737.
- Liu, C., Zou, D., Liu, Z., Ye, C., 2020. Ocean warming alters the responses to eutrophication in a commercially farmed seaweed, *gracilariopsis lemaneiformis*. *Hydrobiologia* 847, 879–893.
- Lu, Y., Yuan, J., Lu, X., Su, C., Zhang, Y., Wang, C., Cao, X., Li, Q., Su, J., Ittekkot, V., others, 2018. Major threats of pollution and climate change to global coastal ecosystems and enhanced management for sustainability. *Environmental Pollution* 239, 670–680.
- Mehvar, S., Filatova, T., Dastgheib, A., De Ruyter van Steveninck, E., Ranasinghe, R., 2018. Quantifying economic value of coastal ecosystem services: A review. *Journal of marine science and engineering* 6, 5.
- Mukherjee, S., Ghosh, K.K., Chanda, A., 2023. Coastal pollution—an overview. *Environmental Oceanography and Coastal Dynamics: Current Scenario and Future Trends* 99–107.
- Nyberg, C.D., Thomsen, M.S., Wallentinus, I., 2009. Flora and fauna associated with the introduced red alga *gracilaria vermiculophylla*. *European Journal of Phycology* 44, 395–403.
- Ramus, A.P., Silliman, B.R., Thomsen, M.S., Long, Z.T., 2017. An invasive foundation species

- enhances multifunctionality in a coastal ecosystem. *Proceedings of the national academy of sciences* 114, 8580–8585.
- Román, A., Oiry, S., Davies, B.F., Rosa, P., Gernez, P., Tovar-Sánchez, A., Navarro, G., Méléder, V., Barillé, L., 2024. Mapping intertidal microphytobenthic biomass with very high-resolution remote sensing imagery in an estuarine system. *Science of The Total Environment* 177025.
- Román, A., Prasyad, H., Oiry, S., Davies, B.F., Brunier, G., Barillé, L., 2023. Mapping intertidal oyster farms using unmanned aerial vehicles (UAV) high-resolution multispectral data. *Estuarine, Coastal and Shelf Science* 291, 108432.
- Román, A., Tovar-Sánchez, A., Olivé, I., Navarro, G., 2021. Using a UAV-mounted multi-spectral camera for the monitoring of marine macrophytes. *Frontiers in Marine Science* 8, 722698.
- Rueness, J., 2005. Life history and molecular sequences of *gracilaria vermiculophylla* (gracilariales, rhodophyta), a new introduction to european waters. *Phycologia* 44, 120–128.
- Seitz, R.D., Wennhage, H., Bergström, U., Lipcius, R.N., Ysebaert, T., 2014. Ecological value of coastal habitats for commercially and ecologically important species. *ICES Journal of Marine Science* 71, 648–665.
- Small, C., Nicholls, R.J., 2003. A global analysis of human settlement in coastal zones. *Journal of coastal research* 584–599.
- Sotka, E.E., Baumgardner, A.W., Bippus, P.M., Destombe, C., Duermit, E.A., Endo, H., Flanagan, B.A., Kamiya, M., Lees, L.E., Murren, C.J., others, 2018. Combining niche shift and population genetic analyses predicts rapid phenotypic evolution during invasion. *Evolutionary Applications* 11, 781–793.
- Thomsen, M.S., Stæhr, P.A., Nejrup, L., Schiel, D.R., 2013. Effects of the invasive macroalgae *gracilaria vermiculophylla* on two co-occurring foundation species and associated invertebrates. *Aquatic Invasions* 8, 133–145.
- Unsworth, R.K., Cullen-Unsworth, L.C., Jones, B.L., Lilley, R.J., 2022. The planetary role of seagrass conservation. *Science* 377, 609–613.
- Van Katwijk, M., 2003. Reintroduction of eelgrass (*zostera marina* l.) in the dutch wadden sea: A research overview and management vision, in: *Challenges to the Wadden Sea Area*. In: *Proceedings of the 10th International Scientific Wadden Sea Symposium*, Groningen, the Netherlands. pp. 173–195.
- Watanabe, Y., Kawamura, T., Yamashita, Y., 2018. Introduction: The coastal ecosystem complex as a unit of structure and function of biological productivity in coastal areas. *Fisheries science* 84, 149–152.
- Weinberger, F., Buchholz, B., Karez, R., Wahl, M., 2008. The invasive red alga *gracilaria vermiculophylla* in the baltic sea: Adaptation to brackish water may compensate for light limitation. *Aquatic Biology* 3, 251–264.
- Zoffoli, M.L., Gernez, P., Godet, L., Peters, S., Oiry, S., Barillé, L., 2021. Decadal increase in the ecological status of a north-atlantic intertidal seagrass meadow observed with multi-mission satellite time-series. *Ecological Indicators* 130, 108033.