Spatial distribution of *Gracillaria* vermiculophylla in French and Spanish estuaries using Unmaned Aerial Vehicule.

Simon Oiry¹

Bede Ffinian Rowe Davies¹ Laurent Barillé¹ Pierre Gernez¹

2024-12-03

To be Written

1

1 Introduction

Coastal ecosystems are among the most dynamic and productive environments on Earth, providing ecosystem services and supporting biodiversity (Barbier et al., 2011; Unsworth et al., 2022; Watanabe et al., 2018). These ecosystems, spanning mangroves, salt marshes, seagrass meadows, and rocky intertidal zones, play a pivotal role in carbon sequestration, nutrient cycling, and shoreline stabilization (Liquete et al., 2013; Mehvar et al., 2018). They also serve as habitats for numerous species, many of which are commercially or ecologically significant (De Valck et al., 2023; Seitz et al., 2014). Coastal areas are densely populated, with billions of people globally depending on their resources for livelihoods, fisheries, and tourism (Mukherjee et al., 2023; Small and Nicholls, 2003). However, coastal ecosystems face mounting pressures from human activities such as land reclamation, pollution, and overfishing, compounded by the impacts of climate change (Hall-Spencer and Harvey, 2019; Lu et al., 2018). Sea level rise, ocean acidification, and increasing storm intensity further exacerbate the vulnerability of these systems, threatening their resilience and the services they provide (He and Silliman, 2019).

One of the significant threats to coastal ecosystems is biological invasions by non-native species, which can disrupt native biodiversity and alter ecosystem functions (Capdevila et

¹Institut des Substances et Organismes de la Mer, ISOMer, Nantes Université, UR 2160, F-44000 Nantes, France

al., 2019; Krueger-Hadfield, 2018; Liu et al., 2020). Gracilaria vermiculophylla, an invasive red macroalga native to the northwest Pacific, exemplifies this issue. Over the last century, this species has spread extensively across temperate estuaries in North America, Europe, and other regions, facilitated by aquaculture and maritime activities (Krueger-Hadfield et al., 2017; Rueness, 2005; Weinberger et al., 2008). Its success as an invader stems from its tolerance to a wide range of environmental stressors, including temperature (Sotka et al., 2018), salinity (Weinberger et al., 2008), and nutrient variability (Abreu et al., 2011), as well as its ability to establish in soft sediment habitats traditionally devoid of macroalgae (Ramus et al., 2017). While G. vermiculophylla can provide some ecosystem services, such as habitat for invertebrates and juvenile fish, it often outcompetes native vegetation, alters sediment composition (Nyberg et al., 2009), and disrupts trophic interactions (Ginneken et al., 2018). In regions like the Baltic Sea and the eastern United States, it has been documented to negatively affect native fucoids and seagrasses (Thomsen et al., 2013; Van Katwijk, 2003). These impacts underscore the importance of monitoring and managing the spread of G. vermiculophylla, particularly as climate change and anthropogenic pressures continue to facilitate biological invasions.

Remote sensing has revolutionized our ability to monitor and manage ecosystems, offering efficient and scalable methods for detecting environmental changes over large areas (Davies et al., 2024a, 2024b; Zoffoli et al., 2021). Among these technologies, drone-based remote sensing has emerged as a particularly promising tool for studying coastal environments (Román et al., 2024, 2023). Equipped with high-resolution cameras and multispectral or hyperspectral sensors, drones can capture fine-scale spatial and spectral data, enabling researchers to identify and map vegetation, detect stress in plants, and monitor changes over time [Román et al. (2021)] ; Oiry et al. 2024. Unlike traditional satellite imagery, drones provide the flexibility to operate in overcast conditions, achieve higher spatial resolution, and target specific areas of interest. For invasive species like G. vermiculophylla, drones equipped with multispectral sensors can differentiate it from native vegetation based on its unique spectral reflectance characteristics (Davies et al. 2025). This capability not only enhances detection accuracy but also reduces the time and labor associated with traditional field surveys. As the cost of drone technology continues to decrease and advancements in machine learning facilitate data analysis, dronebased remote sensing is becoming increasingly accessible and impactful for ecological research and management.

In this study, we aim to harness the potential of drone-based multispectral remote sensing to map *Gracilaria vermiculophylla* in intertidal zones. A Neural Network classification model (DISCOV;Oiry et al. (2024a)), developed by Oiry et al. (2024b) to map intertidal vegetation in Europe has been update to improve red algae detection capability of the DISCOV model. Bla bla what are we going to do? bla bla .

2 Materiel & Methods

2.1 Study sites

Field campaigns were conducted at three study sites in France and Spain. At each site, two locations were investigated Figure 1. The Aven & Belon Estuary in South Brittany, France (Figure 1 A & C), is a dynamic ria-type system hosting diverse habitats, including sandy tidal flats and subtidal zones with coarse, marine-origin sediments (Castaing and Guilcher, 1995; Michel et al., 2021). These habitats support key benthic species such as *Scrobicularia plana*, *Cerastoderma edule*, and *Tellina tenuis*, which play essential roles in sediment bioturbation and nutrient cycling (Blanchet et al., 2014; Tankoua et al., 2011). The estuary serves as a nursery for juvenile fish and a feeding ground for migratory birds, with its ecological productivity driven by a mix of euryhaline and marine species adapted to salinity gradients (Blanchet et al., 2014). Oyster farming, particularly *Crassostrea gigas*, is a dominant activity, altering sediment dynamics and local biodiversity (Michel et al., 2021). Despite its ecological richness, the estuary faces pressures from nutrient loading and physical alterations, with bioindicators like *S. plana* used to monitor the impacts of salinity, sediment quality, and pollution (Tankoua et al., 2011).

The Saja-Besaya Estuary, situated along the Cantabrian Sea in northern Spain, is characterized by the confluence of the Saja and Besaya rivers near Torrelavega (Figure 1 C). The estuary, also known as San Martín de la Arena or Suances Estuary, has been subject to significant anthropogenic pressures, including industrial developments throughout the 20th century. These activities have led to contamination from mining, paper manufacturing, and carbonate discharges, classifying the estuary as highly polluted near its upper reaches (Ortega et al., 2005). This contamination impacts the estuarine ecosystem, including water quality and biodiversity, with minimal aquatic life and sparse riverbank vegetation in its lower sections (Romero et al., 2008).




Figure 1: Location of the drone flights. A: Flights made in Aven Estuary, France; B: Flights made in Bélon Estuary, France; C: Flights made in Saja Estuaries, Spain. Golden polygons represent intertidal areas.

2.2 Remote sensing data acquisition and pre-processing

A total of 6 drone flights were done spread in the 3 study sites. Each time, flights were done at an altitude of 120 m and at a speed of 10 m.s⁻¹ (Table 1).

Table 1: List of drone flights, summarising the location, the date, and the total extent of each flight (in hectars).

Country	Site	Flight	Date	
France	Aven & Belon	Notre-Dame De Tremor	2024-04-11	
		Pont du Guilly	2024-04-11	
Spain	Saja Estuary	Marisma de Cortiguera	2024-06-25	
		Marisma de Cudón	2024-06-25	

2.2.1 Multispectral data

At each location, reflectance images with a resolution of 1.2 million pixels were captured using a DJI Matrice 300 quadcopter drone equipped with a Micasense RedEdge Dual MX multispectral camera. The camera recorded data across ten spectral bands, spanning from blue to near-infrared (NIR) wavelengths (444, 475, 531, 560, 660, 668, 705, 717, 740, and 840 nm) (). To ensure consistent lighting conditions, the drone's flight trajectory was aligned to maintain a solar azimuth angle of 90 degrees. Image acquisition was carried out with an overlap of 70% between side-by-side images and 80% between successive images along the flight path. A downwelling light sensor (DLS2) was used to measure real-time irradiance, enabling the correction of reflectance values for variations in light intensity caused by cloud cover during the flight. The raw image data were subsequently calibrated to reflectance using a calibration panel with ~50% reflectivity, provided by the camera's manufacturer. Images were processed using structure-from-motion photogrammetry software (Agisoft, 2019) to generate multispectral ortho-mosaics for each flight. The ortho-mosaicking workflow was consistent across all flights. Initially, key tie points were identified within each image and across overlapping images to create a sparse point cloud. This point cloud was refined by removing noisy points using a reprojection accuracy metric. Subsequently, a dense point cloud was generated using a structure-from-motion algorithm. A digital surface model (DSM) was then created through surface interpolation of the dense point cloud, which served as the basis for reconstructing the multispectral ortho-image (Nebel et al., 2020). The resolution of the multispectral orthomosaic obtained were 8 cm per pixel.

2.2.2 LiDAR data

LiDAR standing for Light Detection and Ranging uses lasers to measure distances by timing reflected pulses, creating detailed 3D maps of surfaces.

Using the Matrice 300 Series Dual Gimbal Connector, a DJI Zenmuse L1 LiDAR and RGB sensor was mounted on the drone alongside a multispectral camera. This setup enabled the simultaneous capture of LiDAR point clouds, high-resolution RGB images, and multispectral images collected by the MicaSense RedEdge Dual MX during the same flight. The same processing workflow as Section 2.2.1 was applied to process LiDAR RGB images, resulting in ortho-mosaic with a resolution of 2.5 cm per pixel. Since the mapping focused solely on flat surfaces without dense vegetation, the LiDAR measured only a single return. Operating in repetitive scanning mode with a sampling rate of 240 kHz, the system achieved a point density of 350 points per square meter. The LiDAR point cloud was extracted and converted into LAS format using DJI Terra software. The LAS point cloud was then imported into Agisoft Metashape (Agisoft, 2019) to generate a DEM with a resolution of 2.5 cm.

2.3 Scene classification

A neural network classification model (DISCOV; Oiry et al. (2024b); Oiry et al. (2024a)), previously applied with success to Micasense reflectance data for mapping intertidal vegetation along the Portuguese and French Atlantic coasts, has been used in this study. The training dataset of DISCOV v1.0 has been updated. As shown by Oiry et al. (2024b) the DISCOV v1.0 model (Oiry et al., 2024a) was trained using only 5771 Rhodophyceae pixel (3% of the training dataset). To fill this gap the original training dataset of DISCOV v1.0 was updated using new training pixel coming from the 5 drone flights (Section 2.2).

3 Results




Figure 2: DISCOV Prediction (A), RGB composition (B) and Bathymetry (C) of the Bélon estuary site in Brttany, France. The total extent of this flight was 21 hectars with a resolution of 8 mm per pixel. Bathymetry is represented as the height above mean sea level.

4 Discussion

5 Conclusion

References

- Abreu, M.H., Pereira, R., Buschmann, A., Sousa-Pinto, I., Yarish, C., 2011. Nitrogen uptake responses of gracilaria vermiculophylla (ohmi) papenfuss under combined and single addition of nitrate and ammonium. Journal of Experimental Marine Biology and Ecology 407, 190–199.
- Agisoft, 2019. Agisoft metashape.
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C., Silliman, B.R., 2011. The value of estuarine and coastal ecosystem services. Ecological monographs 81, 169–193.
- Blanchet, H., Gouillieux, B., Alizier, S., others, 2014. Multiscale patterns in the diversity and organization of benthic intertidal fauna among french atlantic estuaries. Journal of Sea Research 90, 95–110. https://doi.org/10.1016/j.seares.2014.02.014
- Capdevila, P., Hereu, B., Salguero-Gómez, R., Rovira, G. la, Medrano, A., Cebrian, E., Garrabou, J., Kersting, D.K., Linares, C., 2019. Warming impacts on early life stages increase the vulnerability and delay the population recovery of a long-lived habitat-forming macroalga. Journal of Ecology 107, 1129–1140.
- Castaing, P., Guilcher, A., 1995. Morphosedimentary evolution of ria-type estuaries. Earth Surface Processes and Landforms 20, 361–376. https://doi.org/10.1002/esp.3290200408
- Davies, B.F.R., Oiry, S., Rosa, P., Zoffoli, M.L., Sousa, A.I., Thomas, O.R., Smale, D.A., Austen, M.C., Biermann, L., Attrill, M.J., others, 2024b. A sentinel watching over intertidal seagrass phenology across western europe and north africa. Communications Earth & Environment 5, 382.
- Davies, B.F.R., Oiry, S., Rosa, P., Zoffoli, M.L., Sousa, A.I., Thomas, O.R., Smale, D.A., Austen, M.C., Biermann, L., Attrill, M.J., others, 2024a. Intertidal seagrass extent from sentinel-2 time-series show distinct trajectories in western europe. Remote Sensing of Environment 312, 114340.
- De Valck, J., Jarvis, D., Coggan, A., Schirru, E., Pert, P., Graham, V., Newlands, M., 2023. Valuing ecosystem services in complex coastal settings: An extended ecosystem accounting framework for improved decision-making. Marine Policy 155, 105761.
- Ginneken, V. van, Vries, E. de, others, 2018. The global dispersal of the non-endemic invasive red alga gracilaria vermiculophylla in the ecosystems of the euro-asia coastal waters including the wadden sea unesco world heritage coastal area: Awful or awesome? Oceanography & Fisheries Open Access Journal 8, 4–26.
- Hall-Spencer, J.M., Harvey, B.P., 2019. Ocean acidification impacts on coastal ecosystem services due to habitat degradation. Emerging Topics in Life Sciences 3, 197–206.
- He, Q., Silliman, B.R., 2019. Climate change, human impacts, and coastal ecosystems in the anthropocene. Current Biology 29, R1021–R1035.

- Krueger-Hadfield, S., 2018. Everywhere you look, everywhere you go, there's an estuary invaded by the red seaweed gracilaria vermiculophylla (ohmi) papenfuss, 1967. BioInvasions Records 7.
- Krueger-Hadfield, S.A., Kollars, N.M., Strand, A.E., Byers, J.E., Shainker, S.J., Terada, R., Greig, T.W., Hammann, M., Murray, D.C., Weinberger, F., others, 2017. Genetic identification of source and likely vector of a widespread marine invader. Ecology and evolution 7, 4432–4447.
- Liquete, C., Piroddi, C., Drakou, E.G., Gurney, L., Katsanevakis, S., Charef, A., Egoh, B., 2013. Current status and future prospects for the assessment of marine and coastal ecosystem services: A systematic review. PloS one 8, e67737.
- Liu, C., Zou, D., Liu, Z., Ye, C., 2020. Ocean warming alters the responses to eutrophication in a commercially farmed seaweed, gracilariopsis lemaneiformis. Hydrobiologia 847, 879–893.
- Lu, Y., Yuan, J., Lu, X., Su, C., Zhang, Y., Wang, C., Cao, X., Li, Q., Su, J., Ittekkot, V., others, 2018. Major threats of pollution and climate change to global coastal ecosystems and enhanced management for sustainability. Environmental Pollution 239, 670–680.
- Mehvar, S., Filatova, T., Dastgheib, A., De Ruyter van Steveninck, E., Ranasinghe, R., 2018. Quantifying economic value of coastal ecosystem services: A review. Journal of marine science and engineering 6, 5.
- Michel, G., Le Bot, S., Lesourd, S., Lafite, R., 2021. Morpho-sedimentological and dynamic patterns in a ria type estuary: The belon estuary (south brittany, france). Journal of Maps 17, 389–400. https://doi.org/10.1080/17445647.2021.1925170
- Mukherjee, S., Ghosh, K.K., Chanda, A., 2023. Coastal pollution—an overview. Environmental Oceanography and Coastal Dynamics: Current Scenario and Future Trends 99–107.
- Nebel, S., Beege, M., Schneider, S., Rey, G.D., 2020. A review of photogrammetry and photorealistic 3D models in education from a psychological perspective, in: Frontiers in Education. Frontiers Media SA, p. 144.
- Nyberg, C.D., Thomsen, M.S., Wallentinus, I., 2009. Flora and fauna associated with the introduced red alga gracilaria vermiculophylla. European Journal of Phycology 44, 395–403.
- Oiry, S., Davies, B.F.R., Pierre, G., Laurent, B., 2024a. DISCOV: Drone Intertidal Sediment Classification Of Vegetation. https://doi.org/10.5281/zenodo.14218984
- Oiry, S., Davies, B.F.R., Sousa, A.I., Rosa, P., Zoffoli, M.L., Brunier, G., Gernez, P., Barillé, L., 2024b. Discriminating seagrasses from green macroalgae in european intertidal areas using high-resolution multispectral drone imagery. Remote Sensing 16. https://doi.org/10.3390/rs16234383
- Ortega, T., Ponce, R., Forja, J., Gómez-Parra, A., 2005. Fluxes of dissolved inorganic carbon in three estuarine systems of the cantabrian sea (north of spain). Journal of Marine Systems 53, 125–142.
- Ramus, A.P., Silliman, B.R., Thomsen, M.S., Long, Z.T., 2017. An invasive foundation species enhances multifunctionality in a coastal ecosystem. Proceedings of the national academy of sciences 114, 8580–8585.
- Román, A., Oiry, S., Davies, B.F., Rosa, P., Gernez, P., Tovar-Sánchez, A., Navarro, G., Méléder, V., Barillé, L., 2024. Mapping intertidal microphytobenthic biomass with very

- high-resolution remote sensing imagery in an estuarine system. Science of The Total Environment 177025.
- Román, A., Prasyad, H., Oiry, S., Davies, B.F., Brunier, G., Barillé, L., 2023. Mapping intertidal oyster farms using unmanned aerial vehicles (UAV) high-resolution multispectral data. Estuarine, Coastal and Shelf Science 291, 108432.
- Román, A., Tovar-Sánchez, A., Olivé, I., Navarro, G., 2021. Using a UAV-mounted multispectral camera for the monitoring of marine macrophytes. Frontiers in Marine Science 8, 722698.
- Romero, M., Andrés, A., Alonso, R., Viguri, J., Rincón, J.M., 2008. Sintering behaviour of ceramic bodies from contaminated marine sediments. Ceramics International 34, 1917–1924.
- Rueness, J., 2005. Life history and molecular sequences of gracilaria vermiculophylla (gracilariales, rhodophyta), a new introduction to european waters. Phycologia 44, 120–128.
- Seitz, R.D., Wennhage, H., Bergström, U., Lipcius, R.N., Ysebaert, T., 2014. Ecological value of coastal habitats for commercially and ecologically important species. ICES Journal of Marine Science 71, 648–665.
- Small, C., Nicholls, R.J., 2003. A global analysis of human settlement in coastal zones. Journal of coastal research 584–599.
- Sotka, E.E., Baumgardner, A.W., Bippus, P.M., Destombe, C., Duermit, E.A., Endo, H., Flanagan, B.A., Kamiya, M., Lees, L.E., Murren, C.J., others, 2018. Combining niche shift and population genetic analyses predicts rapid phenotypic evolution during invasion. Evolutionary Applications 11, 781–793.
- Tankoua, O.F., Buffet, P.-E., Amiard, J.-C., Amiard-Triquet, C., Mouneyrac, C., Berthet, B., 2011. Potential influence of confounding factors (size, salinity) on biomarkers in the sentinel species scrobicularia plana used in programmes monitoring estuarine quality. Environmental Science and Pollution Research 18, 1253–1263. https://doi.org/10.1007/s11356-011-0479-3
- Thomsen, M.S., Stæhr, P.A., Nejrup, L., Schiel, D.R., 2013. Effects of the invasive macroalgae gracilaria vermiculophylla on two co-occurring foundation species and associated invertebrates. Aquatic Invasions 8, 133–145.
- Unsworth, R.K., Cullen-Unsworth, L.C., Jones, B.L., Lilley, R.J., 2022. The planetary role of seagrass conservation. Science 377, 609–613.
- Van Katwijk, M., 2003. Reintroduction of eelgrass (zostera marina l.) in the dutch wadden sea: A research overview and management vision, in: Challenges to the Wadden Sea Area. In: Proceedings of the 10th International Scientific Wadden Sea Symposium, Groningen, the Netherlands. pp. 173–195.
- Watanabe, Y., Kawamura, T., Yamashita, Y., 2018. Introduction: The coastal ecosystem complex as a unit of structure and function of biological productivity in coastal areas. Fisheries science 84, 149–152.
- Weinberger, F., Buchholz, B., Karez, R., Wahl, M., 2008. The invasive red alga gracilaria vermiculophylla in the baltic sea: Adaptation to brackish water may compensate for light limitation. Aquatic Biology 3, 251–264.
- Zoffoli, M.L., Gernez, P., Godet, L., Peters, S., Oiry, S., Barillé, L., 2021. Decadal increase

in the ecological status of a north-atlantic intertidal seagrass meadow observed with multimission satellite time-series. Ecological Indicators 130, 108033.