

Remote Sensing of Environment

Heatwave impacts on intertidal seagrass reflectance: from laboratory experiment to satellite mapping of seagrass heat shock index --Manuscript Draft--

Manuscript Number:	RSE-D-25-00697
Article Type:	Research Paper
Section/Category:	Ecological applications (wetland, habitat, animal population, etc.)
Keywords:	Spectral change; Air exposition; Coastal Ecosystems; Extreme events; Zostera noltei; Leaf browning
Corresponding Author:	Simon Oiry Nantes Université FRANCE
First Author:	Simon Oiry
Order of Authors:	Simon Oiry Bede Ffinian Rowe Davies Philippe Rosa Augustin Debly Maria Laura Zoffoli Anne-Laure Barillé Nicolas Harin Marta Román Pierre Gernez Laurent Barillé
Abstract:	Seagrasses play a vital role in coastal ecosystems, providing habitat, stabilizing sediments, and contributing to carbon sequestration. However, climate change has increased the frequency and intensity of heatwaves, posing a significant threat to seagrass health. This study investigates the effects of marine and atmospheric heatwaves on the spectral reflectance of the intertidal seagrass <i>Zostera noltei</i>. Laboratory experiments were conducted under controlled heatwave conditions, where hyperspectral reflectance measurements were taken to assess the impacts over time. Heatwaves caused a substantial decline in seagrass reflectance, particularly in the green and near-infrared regions, corresponding to the browning of green leaves. Key vegetation indices, including the Normalized Difference Vegetation Index (NDVI) and Green Leaf Index (GLI), showed pronounced reductions under heatwave stress, with NDVI values decreasing by up to 34% and GLI by 57%. A novel metric, the Seagrass Heat Shock Index (SHSI), was developed to quantify the transition of seagrass leaves from green to brown, demonstrating a strong ability to capture the effects of heatwave exposure on seagrass coloration. Multispectral satellite observations corroborated the laboratory results, revealing widespread browning of seagrass leaves during marine and atmospheric heatwave events in South Brittany, France. Notably, darkened seagrass patches were observed in intertidal areas exposed to temperatures exceeding 32°C for over 13.5 hours per day. These findings highlight the potential of spectral reflectance as a tool for detecting early signs of heatwave-induced stress in seagrasses, offering a valuable method for remote sensing-based habitat assessment. The present study underscores the potential of remote sensing to capture rapid environmental changes in intertidal zones, enabling for continuous monitoring of seagrass meadows under the current and future climate regimes.
Opposed Reviewers:	



Simon Oiry
PhD student
Nantes University
France
Email: oirysimon@gmail.com

Remote Sensing of Environment
Editor-in-Chief

Dear Editor-in-Chief,

26th February 2025

Please find attached a manuscript entitled "**Heatwave impacts on intertidal seagrass reflectance: from laboratory experiment to satellite mapping of seagrass heat shock index**", for consideration for publication in *Remote Sensing of Environment*.

Seagrasses play a crucial role in coastal ecosystems, providing essential services such as habitat provision, sediment stabilization, and carbon sequestration. They contribute significantly to maintaining water quality, supporting fisheries, and mitigating coastal erosion. However, the increasing frequency and intensity of marine and atmospheric heatwaves, driven by climate change, present a major threat to the health of the habitats they form. These extreme climatic events induce physiological stress, disrupt ecosystem functions, and accelerate habitat degradation, ultimately threatening the biodiversity and ecological resilience of intertidal seagrass meadows. Remote sensing is a vital tool for seagrass monitoring, offering a non-invasive and large-scale approach to detecting changes in seagrass extent and density. This capability is crucial for understanding the rapid impacts of climate change on intertidal ecosystems and developing targeted conservation strategies.

The impact of heatwaves on subtidal seagrasses has been extensively studied; however, research on intertidal seagrasses, which experience both marine and terrestrial stressors, remains limited. Furthermore, while previous studies have predominantly explored the relationship between environmental pressures and physiological responses, none have specifically investigated the effects of extreme climatic events on the spectral reflectance of seagrasses. This study addresses this research gap by analyzing how heatwave-induced stress alters seagrass reflectance using laboratory experiment. From our experimental results, we developed a new seagrass heat shock index (SHSI) specifically designed for remote sensing application. The SHSI is a novel spectral index able to quantify the severity of heatwave impact by measuring and quantifying the transition from green to brown leaves observed in the lab and in the field. We applied and validated the SHSI index to Sentinel-2 images in order to monitor a heatwave event and documented the heatwave impact on seagrass spatial distribution and temporal variations. Altogether, our results demonstrate the potential of remote sensing to detect early signs of heatwave-induced stress in seagrasses and enable large-scale monitoring of vulnerable intertidal habitats.

This new capacity to detect heatwave-impacted seagrass at large scale using remote sensing will be particularly significant for conservation and rewilding initiatives. Seagrass restoration is a nature-based solution for greenhouse gas mitigation that could be negatively affected by heatwaves. Restoration is currently increasing worldwide supporting the Sustainable Development Goals (SDGs) of the United Nations.

For the consideration of this manuscript, we suggest the following reviewers:

- **Dimitris Poursanidis**, Foundation for Research and Technology—Hellas, Greece
(dpoursanidis@terrasolutions.eu)
- **Bárbara Ondiviela**, University of Cantabria, Spain (barbara.ondiviela@unican.es)
- **Francesco Boscutti**, University of Udine, Italy (francesco.boscutti@uniud.it)
- **Laura Govers**, Royal Netherlands Institute for Sea Research, Netherlands
(lauralgovers@gmail.com)
- **Isabel Caballero de Frutos**, Spanish National Research Council, Spain
(isabel.caballero@icman.csic.es)

The first author, Simon Oiry, is also the corresponding author. The work described, its production, and authorship conform to Nantes University's policies on ethical and responsible research conduct. We confirm that this manuscript has not been published elsewhere and is not under consideration by another journal. All authors have approved the manuscript and agree with its submission to *Remote Sensing of Environment*.

We look forward to hearing from you in due course.

Yours sincerely,

Simon Oiry
On behalf of the authors

- First study investigating heatwave impacts on intertidal seagrass reflectance.
- Heatwaves reduce seagrass reflectance, especially in green and NIR regions.
- A novel Seagrass Heat Shock Index (SHSI) detects heatwave-induced leaf browning.
- Sentinel-2 satellite data confirm large-scale seagrass darkening during heatwaves.
- Seagrass exposed to >13h of 32°C air per day showed the most severe browning.

[Click here to view linked References](#)

Heatwave impacts on intertidal seagrass reflectance: from laboratory experiment to satellite mapping of seagrass heat shock index

Simon Oiry¹, Bede Ffinian Rowe Davies¹, Philippe Rosa¹, Augustin Debly¹, Maria Laura Zoffoli², Anne-Laure Barillé³, Nicolas Harin³, Marta Román^{1, 4}, Pierre Gernez¹, Laurent Barillé¹

Keywords:

Spectral change, Air exposition, Coastal Ecosystems, Extreme events, *Zostera noltei*, Leaf browning

¹ Institut des Substances et Organismes de la Mer, ISOMer, Nantes Université, UR 2160, F-44000 Nantes, France

² Consiglio Nazionale delle Ricerche, Istituto di Scienze Marine (CNR-ISMAR), 00133 Rome, Italy

³ Bio-littoral, Immeuble Le Nevada, 2 Rue du Château de l'Eraudière, 44300 Nantes, France

⁴ Centro de Investigación Mariña, Departamento de Ecoloxía e Bioloxía Animal, Facultade de Ciencias do Mar, Universidade de Vigo, Campus Universitario As Lagoas-Marcosende, 36310 Vigo, Spain

Abstract

Seagrasses play a vital role in coastal ecosystems, providing habitat, stabilizing sediments, and contributing to carbon sequestration. However, climate change has increased the frequency and intensity of heatwaves, posing a significant threat to seagrass health. This study investigates the effects of marine and atmospheric heatwaves on the spectral reflectance of the intertidal seagrass *Zostera noltei*. Laboratory experiments were conducted under controlled heatwave conditions, where hyperspectral reflectance measurements were taken to assess the impacts over time. Heatwaves caused a substantial decline in seagrass reflectance, particularly in the green and near-infrared regions, corresponding to the browning of green leaves. Key vegetation indices, including the Normalized Difference Vegetation Index (NDVI) and Green Leaf Index (GLI), showed pronounced reductions under heatwave stress, with NDVI values decreasing by up to 34% and GLI by 57%. A novel metric, the Seagrass Heat Shock Index (SHSI), was developed to quantify the transition of seagrass leaves from green to brown, demonstrating a strong ability to capture the effects of heatwave exposure on seagrass coloration. Multispectral satellite observations corroborated the laboratory results, revealing widespread browning of seagrass leaves during marine and atmospheric heatwave events in South Brittany, France. Notably, darkened seagrass patches were observed in intertidal areas exposed to temperatures exceeding

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

31 32°C for over 13.5 hours per day. These findings highlight the potential of spectral
32 reflectance as a tool for detecting early signs of heatwave-induced stress in seagrasses,
33 offering a valuable method for remote sensing-based habitat assessment. The present
34 study underscores the potential of remote sensing to capture rapid environmental
35 changes in intertidal zones, enabling for continuous monitoring of seagrass meadows
36 under the current and future climate regimes.

37

1. Introduction

Seagrasses play a crucial role in coastal ecosystems by providing habitats and feeding grounds for various marine species, supporting marine biodiversity, and contributing to primary production and carbon sequestration (Sousa et al., 2019; Unsworth et al., 2022). Seagrasses are essential for several ecological functions, such as sediment stabilization (Infantes et al., 2022) or eutrophication mitigation by consuming nutrients (Gladstone-Gallagher et al., 2018). This justifies their use as indicators of environmental changes due to their sensitivity to water quality variations (Zoffoli et al., 2021). The interactions between seagrass meadows and their associated herbivores further enhance the delivery of ecosystem services, including coastal protection, fisheries support and provision of habitat and resources for birds (Gardner and Finlayson, 2018; Jankowska et al., 2019; Unsworth and Butterworth, 2021; Zoffoli et al., 2023). Understanding and preserving seagrass is vital for maintaining the biodiversity and productivity of coastal regions (Ramesh and Mohanraju, 2020; Scott et al., 2018).

Despite their crucial role in marine ecosystems, seagrasses face numerous threats that compromise their health and functionality. Intertidal seagrasses are subjected to a combination of aquatic and aerial conditions linked with tidal cycles, and face disturbance from terrestrial and aquatic stressors. Coastal development and human activities are primary threats, reducing the available habitat for seagrasses and

1
2
3
4 57 increasing water turbidity, limiting light penetration, and photosynthesis (Waycott et
5
6
7 58 al., 2009). Seagrasses are also threatened by runoff from agricultural fields and urban
8
9
10 59 areas leading to nutrient enrichment. Eutrophication promotes the growth of seaweed
11
12
13 60 in coastal waters, causing macroalgal blooms that compete with seagrasses for light and
14
15
16 61 nutrients (Brun et al., 2003; Oiry et al., 2024; Thomsen et al., 2023). Pollution from
17
18
19 62 industrial and agricultural sources introduces harmful chemicals and heavy metals into
20
21
22 63 coastal waters, posing toxic risks to seagrass health (Bastos et al., 2023; Green et al.,
23
24
25 64 2021; Zahoor and Mushtaq, 2023). Among manifold anthropogenic stressors, heatwaves
26
27
28 65 (HWs), exacerbated by climate change, pose a severe threat to seagrasses, with
29
30
31 66 catastrophic dieback events observed worldwide (Carlson et al., 2018; Marbà and
32
33
34 67 Duarte, 2010; Moore and Jarvis, 2008; Strydom et al., 2020; Thomson et al., 2015).

35
36 68 Marine Heatwaves (MHWs) are defined by Hobday et al. (2016) as prolonged
37
38
39 69 discrete anomalously warm water events, while Atmospheric Heatwaves (AHW) are
40
41
42 70 defined by Perkins and Alexander (2013) as periods of at least three consecutive days
43
44
45 71 with temperatures exceeding the 90th percentile of a time series covering at least 30
46
47
48 72 years. Subtidal seagrass meadows are exposed to MHWs, whereas at the interface
49
50
51 73 between land and ocean, intertidal seagrasses are exposed to both MHWs and AHWs.
52
53
54 74 HWs profoundly impact seagrass physiology, with effects varying between species and
55
56
57 75 geographic location. Widespread seagrass species such as *Zostera marina* exhibits high
58
59
60 76 susceptibility to elevated sea surface temperatures during winter and spring, leading to
61
62
63
64
65

advanced flowering, high mortality rates, and reduced biomass (Sawall et al., 2021). Similarly, *Cymodocea nodosa* shows increased photosynthetic activity during HWs but suffers negative effects on photosynthetic performance and leaf biomass during recovery (Deguette et al., 2022). Additionally, different populations of *Zostera marina* along the European thermal gradient exhibit varied photophysiological responses during the recovery phase of HWs, indicating differential adaptation capabilities among populations (Winters et al., 2011). High-latitude populations exhibited prolonged declines in photophysiological performance even after temperatures returned to control levels, whereas the low-latitude Adriatic population showed full recovery (Winters et al., 2011). These events intensify other stressors, such as overgrazing and seed burial, compromising recruitment (Guerrero-Meseguer et al., 2020). Although extensive research exists on marine heatwaves' effects on subtidal seagrasses (Arias-Ortiz et al., 2018; Deguette et al., 2022; Strydom et al., 2020), less attention has been given to intertidal habitats and even less to the effect of atmospheric extreme events on intertidal seagrass. Nonetheless, recent research showed that the low tide exposure of *Zostera noltei* to a simulated four-day atmospheric HW caused significant decreases in its photosynthetic efficiency, resulting in leaf necrosis and decay (Román et al., 2023).

The increased occurrence of extreme climate events calls for the implementation of monitoring strategies able to provide detailed and spatially explicit assessments of HWs effects on seagrass meadows. In such context, remote sensing, whose ability to

map seagrass distribution over a variety of spatio-temporal scales has been demonstrated (Davies et al., 2024a, 2024b; Oiry et al., 2024; Román et al., 2021), proved useful to study the changes in seagrass coverage caused by extreme HW event (Strydom et al., 2020). The pigment composition of plants, such as chlorophylls, carotenoids, and anthocyanins, significantly influences their spectral signature in the visible range due to their specific light absorption properties (Davies et al., 2023; Douay et al., 2022; Olmedo-Masat et al., 2020; Ustin and Jacquemoud, 2020). During the senescence phase of seagrass' life-cycle, the degradation of chlorophyll and the unmasking of accessory pigments result in noticeable changes in leaf coloration and reflectance, including increased reflectance in the red and green wavelengths and shifts in the red-edge position (Boyer et al., 1988; Mariën et al., 2019; Peñuelas et al., 2004). Leaf browning, often observed after stress events, produces reflectance changes similar to those caused by senescence, enabling the detection of vegetation stress through remote sensing (Boyer et al., 1988; Peñuelas et al., 2004). Spectral indices such as the Brown Pigment Index (BPI) and the Photochemical Reflectance Index (PRI) have been developed to assess changes in terrestrial plant physiological status, including oxidative and drought stress (Garbulsky et al., 2011; Skendzic, 2023). While these effects are well-documented in terrestrial plants, the spectral reflectance changes associated with senescence and stress events such as MHWs or AHWs remain poorly studied on intertidal seagrasses.

This study aims to experimentally test the hypothesis that HWs alter the reflectance of the intertidal seagrass *Zostera noltei*. Controlled experiments in intertidal chambers were conducted to evaluate the direct impact of heat stress on seagrass reflectance. The findings will then be applied to satellite remote sensing images, providing critical insights into the spatial extent and temporal dynamics of HW effects on seagrass meadows. By linking experimental results with large-scale observations of seagrass leaves' browning, the study aims to underscore the potential of remote sensing to enhance our understanding of seagrass responses to extreme thermal events across diverse settings and timescales.

2. Material & Methods

2.1 Laboratory Experiment

2.1.1 Sampling and acclimation of seagrasses

Seagrass samples were taken in summer 2024, at low tide, from a *Zostera noltei* (dwarf eelgrass) meadow located in Bourgneuf Bay, France (46°57'32.0"N, 2°10'37.0"W). A metal coring device was used to sample seagrass from an area of 30x15x5 cm (length x width x depth, respectively), maintaining the sediment structure and avoiding damage to seagrass rhizomes and leaves (Figure 1 A). These coring device enabled the collection of sediment samples at a consistent depth, minimizing variability between samples. A total of six samples were collected. Samples including seagrass, sediment, meiofauna, and macrofauna, were placed in plastic trays. Keeping the entire biota allowed for natural interactions between components and reduced stress on the seagrass. Seawater was added to each tray to avoid hydric stress caused by insufficient moisture during transportation (1h drive from the laboratory). Simultaneously, seawater was sampled 4km away from the seagrass sampling site and transported to the lab, where it was filtered using a 0.22 µm nitrocellulose filter to remove suspended particulate matter. The filtered seawater was used in the acclimation tank and the intertidal chambers. The seagrasses were acclimated for one week with a water temperature of 17°C, matching

the *in situ* temperature during sampling, and a photosynthetically active radiation (PAR) of 150 $\mu\text{mol}\cdot\text{s}^{-1}\cdot\text{m}^{-2}$.

2.1.2 Experimental design

A tidal cycle (i.e. regularly alternating 6h of low-tide and 6h of high-tide) was simulated in the laboratory using an intertidal chamber system from ElectricBlue® (Figure 1B ; Electric Blue (2023)). The experimental setup allowed only two tidal states: high tide or low tide, with no intermediate stages. The transition between these states took about 15 minutes to complete after initiation. During the phase of high tide, a volume of 30 L of filtered seawater was pumped and circulated through the chamber (Figure 1 B, C). During low tide, the seagrass sample was emerged. The acclimated seagrasses were split into two subsets and placed in two independent chambers used in parallel, with one chamber used for control and the other for the experimental treatment. The intertidal chambers were equipped with LED lights that emitted a low amount of red and infrared radiation. To achieve a Photosynthetically Active Radiation (PAR) intensity of up to 400 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$, a filament bulb was added inside the chambers. During the diurnal phase of the experiment, the PAR was kept constant in both intertidal chambers. To follow the circadian cycle, these lights (both LED and filament bulb) were turned on and off each day, at the time of sunrise and sunset, respectively.

Air temperature and water temperature were controlled inside the experiment chambers in order to reproduce the range of variability observed in the field (Figure 2).

Field temperature was measured using *in situ* sensors (T7.3 EnvLoggers from ElectricBlue®) deployed at the sampling site in August 2024. The loggers were positioned along a transect from the upper to the lower intertidal zone, attached to pre-existing wooden poles at the sediment surface. In complement, the daily temperature maxima recorded *in situ* were compared with measurements from the nearest Météo France weather station (Annex A). The control chamber was kept at temperatures representing typical summer conditions, with water temperatures at 18°C and air temperatures from 19°C to 23°C, following natural daily temperature fluctuations (Figure 2). In the treatment chamber, the air temperature was adjusted to mimic an AHW that affected the seagrass meadow in Quiberon, South Brittany, France (47°35'40.0"N, 3°07'30.0"W), from September 2 to September 6, 2021. Air temperature in the experimental chamber was set to vary from 23°C (at night) to 35°C (daytime) during the first day of the experiment, and increase by 1°C daily during three consecutive days. Water temperature in the experimental chamber was also adjusted to mimic MHW conditions, starting at the seasonal baseline (18°C) and rising incrementally by 0.5°C daily to simulate the increasing temperatures during the event. This aimed to reproduce the thermal stress experienced by the seagrass meadow during a MHW (Figure 2). The experiment, with both treatment and control chambers, was repeated three times to obtain replicates (hereafter referred to as "Run").

2.1.3 Optical measurements

3.1.3.1 Hyperspectral reflectance measurements

Throughout the experiment, the hyperspectral reflectance, $R(\lambda)$, of both the control and treatment seagrasses was measured using an ASD HandHeld 2 equipped with a fiber optic extension placed inside the chamber. The measurement set up made it possible to automatically acquire $R(\lambda)$ without opening the chamber. An average of five $R(\lambda)$ spectra, each with an integration time of 544 ms, was taken every minute during daytime (Malvern Panalytical, 2023). Every 10 minutes, the fiber optic was switched from one intertidal chamber to the other, in order to measure $R(\lambda)$ in both the treatment and control. Light conditions were controlled inside of the chambers and the reflectance calibration was performed each morning at the very first moment of low tide using a Spectralon white reference with 99% Lambertian reflectivity.

3.1.3.2 Spectrum post-processing

A Savitzky-Golay smoothing function with a 5 nm moving window was applied to each spectrum using the “hsdar” package in R (Lehnert et al., 2017). The second derivative at 665 nm ($\delta\delta 665$), showing the highest variability between the control and the treatment, was tested as an indicator of the spectral changes following HWs.

The effect the HW on $R(\lambda)$ was also quantified using two radiometric indices:

- The Normalized Difference Vegetation Index (NDVI, Rouse et al. (1974)), a proxy of chlorophyll-a concentration (Equation 1)

$$NDVI = \frac{R(840) - R(668)}{R(840) + R(668)} \quad (1)$$

where $R(840)$ and $R(668)$ are the reflectance at 840 and 668 nm respectively.

- The Green Leaf Index (GLI, Louhaichi et al. (2001)), a quantification of the seagrass leaves greenness (Equation 2)

$$GLI = \frac{[R(550) - R(668)] + [R(550) - R(450)]}{(2 \times R(550)) + R(668) + R(450)} \quad (2)$$

where $R(550)$ and $R(450)$ are the reflectance in the green (at 550 nm) and in the blue (at 450 nm) spectral bands, respectively.

Based on the observed spectral changes in seagrasses exposed to HWs, we developed a new radiometric index to better detect the radiometric change caused by the HW. The browning of the leaves was characterized by substantial radiometric changes in both the green and red-edge spectral regions. The Seagrass Heat Shock Index (SHSI) was introduced as the reflectance line height at 740 nm, compared to the 560 - 842 nm baseline (Figure 3). Namely the SHSI subtracts the reflectance observed at 740 nm to the linearly interpolated reflectance between 560 and 842 nm, so that the index is positive in the case of brown, HW-impacted seagrass leaves, and negative in the case of green, non-impacted leaves:

$$SHSI = I_{SHSI} - R(740) \quad (3)$$

where :

$$I_{SHSI} = R(560) + \tau[R(842) - R(560)]$$

and :

$$\tau = \frac{740 - 560}{842 - 560}$$

where $R(560)$, $R(740)$, and $R(842)$ represent the reflectance at 560, 740, and 842 nm, respectively, and τ is equal to a constant of 0.64. These wavelengths were selected to align with the spectral resolution of satellites missions such as Sentinel-2, for broader remote sensing application.

2.2 Observation of a seagrass bed impacted by a HWs

Field measurements were taken the 10th of September 2021 after an atmospheric and marine HW in order to assess the impact of heat stress on seagrass. The study site was a seagrass meadow near Quiberon (France : 46°57'32.0"N, 2°10'37.0"W, Figure 4). Brown seagrass leaves were observed over large patches of the meadow alongside areas covered by green seagrass (Figure 5). A total of 96 Quadrat Points (QPs) were collected as georeferenced quadrat images across the meadow. These images allowed for visual assessment of vegetation type, density, and coloration. The quadrats were then divided into two categories: green seagrasses (henceforth: unimpacted QPs) and brown seagrasses (henceforth: impacted QPs), based on a visual estimation of the leaf coloration (Figure 4).

2.2.1 Temperature data and HW detection

3.2.1.1 Air temperature

Hourly air temperature data from 1952 to 2024 (more than 395,000 observation) from a nearby weather station (Lorient-Lann Bihoue, 47°45'46"N 3°26'11"W) was retrieved from Meteo France (<https://portail-api.meteofrance.fr>).

3.2.1.2 Water temperature

Sea Surface Temperature (SST) data from 1982 -2022 over the Quiberon coastal area was downloaded from the Copernicus Marine Data Store (Copernicus Marine Environment Monitoring Service, CMEMS (2024)). An area of 2700 km² was extracted and analyzed. This area was large enough to minimize missing values caused by cloud cover and small enough to limit the influence of offshore SST stability.

3.2.1.3 Heatwave detection and characterization

MHW and AHW detection was performed using the HeatwaveR package in R (Schlegel and Smit, 2018). This package utilizes the methodology proposed by Hobday et al. (2016) to detect HW events. The annual climatology (i.e. the average temperature of each day of the year since the start of the time serie) of both air and water temperature was computed. HWs were defined as events when the temperature exceeded the 90th percentile of the climatology during three consecutive days. Furthermore, the severity of each event was assessed using the methodology proposed by Hobday et al. (2018).

2.2.2 Satellite observations

Three 2021 Sentinel-2 images of the study site were selected to assess the effect of the combined AHW and MHW (“HW event”, henceforth) on the seagrass meadow: the first image was taken 5 days before the HW (1st of September 2021), the second image during the HW (6th of September 2021) and the third image one month later (8th of October 2021). Level-2 data were downloaded from the Copernicus open access hub (European Space Agency, 2024a) provided by the European Space Agency (ESA). Level-2 images consist of orthorectified surface reflectance corrected from the effect of the atmosphere using ESA’s standard correction (i.e., Sen2cor ; European Space Agency, 2024b).

The SHSI (Equation 3) was computed and mapped for each image. For the pixel containing a field QP (Figure 4), the satellite-derived reflectance was extracted, and compared before and after the HW event.

2.2.3 Emersion time of the seagrass meadow

The spatial distribution of seagrass emersion time during low tide was estimated using bathymetric and water level data. High resolution bathymetry data (Litto3D® product) for the Quiberon intertidal meadow were sourced from the “Service Hydrographique et Océanographique de la Marine” (SHOM, 2021), while one-minute interval water level data were downloaded from Intergovernmental Oceanographic Commission data

portal (IOC, 2024), using measurements from the nearest tide gauge at Le Crouesty. A 2.85 m vertical correction was applied to the Litto3D data to align its zero reference with that of the water level data (RAM, SHOM (2022))

Once aligned, the corrected elevation was compared to water height for each pixel and each time step during the HW event. The emersion time was then calculated as the daily total time each pixel remained exposed along the duration of the AHW.

2.3 Statistics

General Linear Mixed effects Models (GLMMs) were used to assess relative differences over time in response variables (Spectral Indices) with different treatments (Impacted vs Unimpacted). To analyze the effect of HW on the reflectance indices observed during the lab experiment, the relative change was modeled as a function of Days (1-3: Discrete) with Runs (1-3: Factor) and Timestep within Run (1-6: Factor) as cross random factors. Satellite-derived SHSI were modeled as a function of Date (1-3: Discrete) and Treatment (Impacted vs Unimpacted: Categorical). A General Additive Model (GAM) was used to assess the relationship between relative SHSI change with emersion time. SHSI was modeled as a function of emersion time with a basis spline. All model parameters were estimated within a Bayesian framework using the “brms” and “RStan” packages in R to leverage the stan language (Bürkner, 2021; Carpenter et al., 2017; R Core Team, 2023; Stan Development Team et al., 2020). The response variables were

modeled assuming a Gaussian distribution, with weakly informative priors (Student-
T(3,0,2.5)). Model parameters were estimated using Markov Chain Monte Carlo
(MCMC) sampling, with 4 chains of 5000 iterations and a warm-up of 500.

3. Results

3.1 Laboratory Experiment

3.1.1 Heatwave effect on seagrass reflectance

During the laboratory HW experiment, the seagrass reflectance was drastically
impacted by the increase in both air and water temperature (Figure 6). The Control
 $R(\lambda)$, displayed a spectral shape typical of seagrass, with a green peak around 560 nm, a
valley associated with chlorophyll-a absorption around 665 nm, and a high near-
infrared (NIR) plateau beyond 705 nm. This remained stable over time, with only
minimal changes in magnitude and spectral features. In contrast, the Treatment $R(\lambda)$
showed severe changes throughout the experiment. During day 1, the Treatment $R(\lambda)$
was generally similar to the Control $R(\lambda)$, despite a slightly less marked peak around
560 nm and slightly lower NIR values from 750 – 900 nm. A drastic decrease was then
observed during days 2 and 3 across all wavelengths, particularly in the green – yellow
spectral region (from 500 – 650 nm) and in the NIR (from 750 - 900 nm). During day 3,
the collapse in $R(\lambda)$ appeared to stabilize in the NIR, whereas it slightly continued in

the green spectral region. At the end of the HW experiment, the $R(\lambda)$ valley around 665 nm was also less pronounced, suggesting a decrease of chlorophyll-a concentration.

3.1.2 Heatwave effect on radiometric indices

All radiometric indices, $R''_{665\text{ nm}}$, NDVI, GLI and SHSI changed after the experimental heatwave (Figure 7 ; Annex B).

At the start of the experiment (day 1), there was no notable difference in $R''_{665\text{ nm}}$, NDVI and GLI between the Treatment and Control groups. (Figure 7 A, B & C). During days 2 and 3 the radiometric indices all decreased strongly, with an overall decline of 68%, 31% and 54 for $R''_{665\text{ nm}}$, NDVI and GLI.

Unlike the other metrics, the SHSI of the Treatment was on average 55 % higher than that of the Control in day 1 (Figure 7 D). By day 2, the SHSI exhibited a rapid increase of approximately 241 %, eventually reaching an overall rise of 420 % by day 3.

With a maximum deviation of 420 %, SHSI emerges as the most sensitive index for detecting seagrass browning. Consequently, only this index was considered for the next steps of the study.

Looking at raw SHSI values revealed clear distinctions between the Control and Treatment groups (Figure 8). On day 1, the SHSI of the Control and Treatment groups were comparable, with median values of -0.11 and -0.08, respectively. By the end of the experiment, seagrasses in the Treatment group exhibited a median SHSI of 0.15, consistent with their visibly darkened appearance. In contrast, the Control group

retained a green appearance throughout the experiment, with a median SHSI of -0.07. A negative SHSI was considered indicative of non-impacted seagrasses, while a positive SHSI was used as a marker for impacted seagrasses.

3.2 HW of September 2021 in Quiberon, South Brittany

3.2.1 Spectral changes

Sentinel-2 images acquired the 1st and 6th of September 2021 were analysed to assess the short-term impact of a HW on seagrass leaves in South Brittany, France (Figure 9 A and C). Both an AHW and a MHW occurred during the summer of 2021, from the 4th to the 7th and from the 3rd to the 8th of September, respectively (Figure 9 B). Within just a few days, the temperature experienced a sharp increase, from 22.2 to 30.8°C in air, and from 17.7 to 19.3 °C in water temperature. During this period, the 90th percentile of the air temperatures was 25.3 °C and 18.8 °C for the water temperature. The air temperature anomaly of 9.9 °C classified the AHW as a strong event, whereas the 1.7 °C anomaly in water temperature classified the MHW as a moderate event.

Two days after the start of the AHW, the Sentinel-2 image from the 6th of September revealed patches of brown seagrass in the true-color composition (Figure 9 C). Such brown patches were absent from the image from the 1st of September, taken before the HW began (Figure 9 A). Before the HW, all QPs appeared green on the Sentinel-2 image, with similar reflectance spectra, typical of green seagrass leaves

(Figure 9 A and D). Their reflectance spectra showed a peak at 560 nm (in the green part of the spectra), low values at 665 nm and a high infrared plateau (> 705 nm). However, on the 6th of September, QPs classified as impacted during the field campaign, showed significant differences in their reflectance spectral shape compared to unimpacted QPs (Figure 9 C and E). The reflectance spectra of brown seagrass were characterized by the loss of the reflectance peak at 560 nm and a decrease in the infrared plateau, which was replaced by a steadily increasing slope up to 940 nm. The darkening of large seagrass patches could also be observed in the true color composition (Figure 9 C)

3.2.2 SHSI metric applied to Sentinel-2

Using Sentinel-2 data and the QPs, SHSI of green seagrass areas that appeared unimpacted by the HW (Unimpacted QPs ; Figure 9 C), showed minimal change between 1st and 6th of September (3 % ; Figure 10). In contrast, seagrass impacted by the HW that turned brown (Impacted QPs ; Figure 9 C) exhibited significant SHSI changes, showing an increase of 97 % during the HW exposure (Figure 10). One month after the event, on the 8th of October 2021, the SHSI of unimpacted seagrass had increased by 14 % compared to the 1st of September. Regarding impacted seagrass, one month after the event, the SHSI decreased to values comparable to those of unimpacted seagrass. This change reflects an increase of 15 % compared to values recorded on the 1st of September.

Using the SHSI (Equation 3), we detected large darkened seagrass patches in the meadow on 6th of September (Figure 11). A total of 26.9 hectares of seagrass turned brown between 1st and 6th of September. The largest brown patch covered nearly 8 hectares. Overall, 24 % of the total seagrass meadow area showed signs of darkening between 1st and 6th of September. Comparing the spatial distribution of darkened patches with the site's topography revealed that 94.6 % of darkened areas were located above a bathymetric level of 3.9 meters (Figure 11, A and B). One month later, on 8th of October, the previously darkened areas appeared to have regained their green coloration (Figure 11 C).

Additionally, there was a clear relationship between seagrass emersion time and seagrass darkening (Figure 12 ; Annex C). Seagrass emerged during less than 13 hours a day were not impacted, whereas seagrass emerged during more than 13 hours turned brown with the maximum darkening occurring on seagrass emerged during more than 14.5 hours daily.

4. Discussion

4.1 Effect of heatwaves on *Zostera noltei* reflectance

This study explored the dual effect of MHWs and AHWs on spectral reflectance of intertidal seagrasses. Significant changes in the reflectance of seagrasses exposed to HWs were observed both in field measurements and the laboratory experiment. Effects were drops of reflectance around 560 nm and around 740 nm (Figure 7) and of the second derivative at 665 nm ($R''665$), NDVI, and GLI (Figure 7 A, B & C) indexes. By contrast, SHSI showed a marked increase, reaching up to 600% for some samples by day 3 (Figure 7 D), which demonstrated its sensitivity at quantifying seagrass browning. These changes suggest a progressive reduction in photosynthetic activity as well as potential structural or physiological changes to the leaves, such as degraded pigmentation or altered light absorption. Moreover, we observed a considerable decrease in the reflectance in the NIR region (700 - 1300 nm) in stressed plants, which is mainly related with the destruction of the membranes (chloroplasts, thylakoids, cell walls) due to thermal oxidative stress, and the consequent decrease of light scattering through the leaf surface (Knipling, 1970). The leaf darkening observed in plants subjected to thermal stress was mainly explained by the oxidation of phenolic compounds by the enzyme polyphenol oxidase and with damage of the photosynthetic apparatus (Allakhverdiev et al., 2008). High temperatures destabilize chloroplasts

membranes, disrupting photosystem II and impairing recovery of photosynthetic function. As chlorophyll-a degrades, the ratio of pigments shifts, with pigments like carotenoids becoming more prominent (Dascaliuc et al., 2007; Jones and Clayton-Greene, 1992). These processes can eventually result in growth impairment and plant death.

The Seagrass Heat Shock Index (Equation 3) developed in this study makes it possible to detect reductions in the green and the Red-Edge regions caused by thermal stress using only three reflectance bands (560 740 and 840 nm). These spectral bands are available in current (i.e. Sentinel-2, Pleiades-Neo, WorldView-3, SkySat, and GeoSat-2) and future multi-spectral missions (i.e. Sentinel-2 Next Generation and Landsat Next), opening new perspectives for the detection and monitoring of heatwave effects on intertidal seagrass. As shown in the next section, SHSI can be used to assess the extent and severity of darkening events across intertidal seagrass meadows from high resolution satellite images.

4.2 Satellite observations of HW effects on intertidal seagrass

Using our new Seagrass Heat Shock Index, Sentinel-2 images acquired before, during, and after a HW event made it possible to detect seagrass leaf darkening at high spatial resolution, as well as to describe the spatial distribution of the heatwave impact over the whole meadow. The analyse of satellite images in Quiberon Bay during the 2021

1
2
3
4 427 HW revealed that most of the meadow darkening took place in the upper intertidal
5
6
7 428 area, where plants were exposed to air for more than 13 hours per day, while no
8
9
10 429 significant browning was observed in the lower intertidal zone. This contrasted with the
11
12
13 430 experiment, in which seagrasses were exposed to a symmetric tidal cycle, spending
14
15
16 431 equal amounts of time submerged and exposed to air. However, in the field, this
17
18
19 432 configuration occurs only in meadows located at hydrographic zero height (mean sea
20
21
22 433 level). Thus, under the same atmospheric temperatures, bathymetry, and therefore,
23
24
25 434 exposure times will modulate the stress effects observed, as seagrasses located lower in
26
27
28 435 the intertidal may be more resistant to HWs than those in upper fringes. Field
29
30 436 observations further revealed that leaf detachment began after the heatwave, causing an
31
32
33 437 apparent decline in seagrass density at the upper intertidal. *Zostera noltei*, as a species
34
35
36 438 inhabiting the intertidal zone and regularly exposed to air, has developed adaptations
37
38
39 439 to minimize hydric stress, such as its narrow leaves, which help reduce water loss
40
41
42 440 during air exposure periods (Cabaço et al., 2009). Nonetheless, if plants at the upper
43
44
45 441 subtidal are exposed to intense heating and irradiance for prolonged periods, these
46
47
48 442 mechanisms may not be able to prevent degradation of photosynthetic pigments,
49
50 443 decreases in photosynthetic capacity and tissue degradation.

51
52
53 444 During the HW, Sentinel-2 imagery indicated that impacted seagrass patches
54
55
56 445 (i.e., areas exhibiting seagrass darkening) experienced a sharp decline in NDVI.
57
58
59 446 Moreover, the Sentinel-2 image acquired one month after the HW showed that the

NDVI of impacted seagrass patches had not recovered to its initial level (Figure 10). NDVI is commonly used as a proxy for seagrass cover (Davies et al., 2024a; Zoffoli et al., 2020), with high NDVI associated with greater seagrass coverage and low NDVI indicating reduced cover. Our findings demonstrate that heatwaves altered the spectral signature of seagrasses, causing a decrease in NDVI—and consequently, in the satellite-derived estimate of seagrass cover—while the actual coverage observed in the field remained stable (see Figure 5). This discrepancy underscores a potential limitation of remote sensing in accurately capturing true seagrass cover during stress events. We demonstrate that such bias could be corrected by the combined analysis of both NDVI and SHSI indices, as proxies of seagrass cover and heatwave impact.

One month after the heatwave, the SHSI in impacted quadrats recovered to similar values to those of the unimpacted ones. Moreover, Sentinel-2 images revealed a partial recovery in areas of the mid to low intertidal, as shown by their return to green color. These results suggested a certain the resilience capacity of *Z. noltei* patches after thermal disturbance. *Zostera noltei* is a dynamic, fast-growing colonizer species (Roca et al., 2016), with fast leaf turnover times and growth rates (Borum et al., 2004) and fast rhizome elongation rates (Duarte, 1991). The return to green color in some parts of the meadow after recovery might be due to new leaf growth, which was likely sustained by the energy reserves of the surviving below ground biomass (Hemminga, 1998).

4.3 Ecological implications of heatwaves impact on *Zostera noltei*

In the present study we observed that heatwave-driven thermal stress changed the spectral reflectance of *Z. noltei* leaves, indicating photosynthetic pigments degradation, with potential negative consequences on health, growth and survival of the above ground biomass. In this sense, previous experiments on the effects of warming on *Zostera noltei* have shown contrasting results, depending on the duration and intensity of heating. In some cases, there were no significant changes in photosynthetic performance or survival under short-term moderate stress (Franssen et al., 2014), or under gradual temperature increases (Román et al., 2022). By contrast, studies focused on extreme temperature events have evidenced decreases of photosynthetic performance and leaf tissue integrity (Massa et al., 2009; Román et al., 2023). Therefore, our findings, jointly with previous research, suggest that *Z. noltei* meadows at the upper intertidal could be a vulnerable species under the future global warming scenarios, in which heatwaves will be more frequent and intense (Climate Change (IPCC), 2023; Stillman, 2019).

The observed tissue degradation, both in the laboratory experiment and the field, can eventually result in decreases of meadows density and cover under recurrent heatwaves. This decline could have cascading effect on species that rely on the provision of refuge and/or food by the seagrass canopy (Zoffoli et al., 2023).

Commercially important fish and shellfish populations that rely on seagrass for

1
2
3
4 486 sustenance and refuge could also be impacted, with negative consequences on local
5
6
7 487 fisheries' productivity and the livelihoods of coastal communities (Unsworth and
8
9
10 488 Cullen-Unsworth, 2014). Furthermore, as seagrass meadows decline, their capacity to
11
12
13 489 act as a blue carbon sink—critical for climate mitigation—could diminish, contributing
14
15
16 490 to increased atmospheric carbon levels (Armitage and Fourqurean, 2016; Samper-
17
18
19 491 Villarreal et al., 2020). Moreover, the sediment stabilization and wave attenuation could
20
21
22 492 be hampered, ultimately increasing the risk of coastal erosion (Calleja et al., 2007;
23
24 493 Folmer et al., 2012; Gacia et al., 1999).

25
26
27 494 Given the combined effects of temperature extremes, eutrophication, and other
28
29
30 495 anthropogenic pressures, targeted management strategies are essential for enhancing
31
32
33 496 seagrass resilience (Loarie et al., 2009). Approaches such as reducing local stressors,
34
35
36 497 cultivating heat-tolerant genotypes, and investing in restoration initiatives are vital to
37
38
39 498 supporting these ecosystems in a warming climate. Although challenges remain, the
40
41
42 499 adaptability and potential resilience of certain seagrass species offer hope for their
43
44 500 persistence amid accelerating ecological shifts.
45
46
47 501

5. Conclusion

This research investigated the effects of both marine and atmospheric heatwaves on the intertidal seagrass *Zostera noltei*, a critical component of coastal ecosystems facing increased thermal stress due to climate change. Heat stress effects on seagrass reflectance were documented in the laboratory using a controlled experiment in order to understand how extreme heat events affect seagrass health and to assess the potential of remote sensing for HW impacts monitoring. Our findings revealed that heatwaves lead to substantial declines in seagrass reflectance, particularly in the green and near-infrared regions, likely driven by pigment degradation and structural damage. This change was reflected in significant reductions in key vegetation indices such as NDVI and GLI. The Seagrass Heat Shock Index (SHSI), developed in this study, successfully detected seagrass darkening, a visible symptom of heatwave stress, demonstrating the viability of spectral monitoring to capture early-stage impacts of heat events on intertidal ecosystems. We applied and validated the newly developed Seagrass Heat Shock Index to Sentinel-2 satellite images acquired during an heatwave event, and documented the broad spatial impact of heatwaves on seagrass meadows in Quiberon Bay, France. The correlation between heatwave exposure and darkening of seagrass suggests that remote sensing, combined with targeted field observations, can enhance our understanding of ecosystem responses to climate-driven thermal events. These results advocate for integrating regular spectral monitoring into conservation strategies,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

522 as it can help predict seagrass resilience and guide adaptive management practices.

523 Future work should focus on refining remote sensing tools and examining the

524 cumulative effects of repeated heatwave events to support the conservation of intertidal

525 seagrass meadows in a warming world.

526

6. Annexes

6.1 Annex A - Temperatures of the experiment

On average, *in situ* temperatures were $3 \pm 3.2^{\circ}\text{C}$ higher than those recorded by Meteo France (Figure A). Additionally, temperatures recorded by Meteo France were more stable than those from the *in situ* sensors, likely due to the sheltered and shaded location of the Meteo France equipment. This difference was used to adjust heatwave temperatures measured by Meteo France to better reflect the conditions experienced by the seagrasses.

1
2
3
4
5 536 **6.2 Annex B - Outputs of GLMM**
6
7

8 Table B
9

10 537
11
12
13

14 538
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

539 **6.3 Annex C- Outputs of GAM**

540 Table C

Figure Captions

Figure 1: Illustration of the experiment. A: Seagrass field sampling using a coring device; B: Intertidal chamber used during the experiment; C: Seagrass sample inside a chamber during the experiment at high tide; D: Treatment sample at the start of the experiment; E: Treatment sample at the end of the experiment, 3 days after the start of the HW event.

Figure 2: Temperature variation in the control (left) and treatment (right) intertidal chambers, during the HW experiment. The red line indicates air temperature, and the blue line water temperature. Due to the tidal cycle of immersion / emersion, the seagrasses experienced the temperatures represented by solid lines.

Figure 3: Computation of the reflectance Seagrass Heat Shock Index (SHSI) for Impacted (A) and Unimpacted (B) seagrass leaves. The dashed line represents the reflectance interpolation between 560 and 842 nm. The red vertical at 740 nm represents the SHSI line height.

Figure 4: Location of field observations in a seagrass meadow impacted by a HW that occurred on the 10th of September 2021 in Quiberon, South Brittany, France. The red line indicates the intertidal zone (Zone between high tide and low tide, exposed during low tide), the dark green area indicates the extent of the seagrass meadow and the olive area indicate saltmarshes. Green points indicate the location of quadrat pictures over

unimpacted seagrasses (i.e. showing a green colour on the field), and orange points indicate the location of quadrats taken over impacted seagrasses (i.e. showing a brown color on the field).

Figure 5: Illustration of the two colorations of seagrass leaves observed in situ on the 10th of September 2021 after a heatwave in Quiberon, South Brittany (France). A: Picture of a zone with both green and brown seagrass; B: Seagrass quadrat with green leaves; C: Seagrass quadrat with brown leaves; D: Picture of a zone where all leaves turned brown.

Figure 6: Standardized hyperspectral reflectance of *Z. noltei* leaves during the HW experiment, showing the Control (Left) and Treatment (Right) measurements. The color indicates the progression along the experiment from the beginning (Day 1: Green), middle (Day 2: Yellow) and end (Day 3: Brown). A min-max standardization was applied to each individual spectrum.

Figure 7: Comparison of spectral metrics for detecting reflectance changes of seagrass leaves after a HW. A: Relative difference between the Treatment and the Control over time for A) the second derivative at 665 nm B) the NDVI C) the GLI and D) the SHSI. Points indicate raw data, the line represents a GLM estimates, while the shaded area is the model's 89 % confidence interval. The dashed lines represent no difference between the Control and the Treatment.

Figure 8: Median of the SHSI across experimental runs, on each day of the experiment.

Error structure represent the 89 % confidence interval (Quantiles 0.055 and 0.945 for lower and upper ribbon, respectively). The green line shows values of the Control group while the orange line indicates values of the Treatment group.

Figure 9: Intertidal seagrass meadow in South Brittany (France) observed before and

during a heatwave (HW). A: RGB color composition of the Sentinel-2 image of the 1st of

September 2021 before the HW; C: RGB color composition of the Sentinel-2 image of the

6th of September 2021 on the second day of a strong AHW. The circles correspond to

QPs collected on the 10th of September 2021, with unimpacted seagrass in green and

impacted seagrass in orange; B: Detection of HW events based on both Air Temperature

and Sea Surface Temperature (SST). The solid line represents the daily average

temperature, while the dashed line indicates the 90th percentile of the climatology.

Coloured areas identify HWs (marine in blue and atmospherical in red). The two

vertical dashed lines represent the acquisition dates of the two Sentinel-2 images (01-09-

2021 and 06-09-2021); D: Sentinel-2 reflectance of seagrass leaves before the HW for both

categories of QPs; E: Sentinel-2 reflectance of seagrass leaves during the HW for both

categories of QPs. Average spectral signatures were obtained in areas where QPs

corresponded to green and brown seagrasses leaves (green and orange circles,

respectively) as identified during the field survey. The shaded areas around the

reflectance spectra represent the standard deviation.

Figure 10: Changes in the relative SHSI estimated from Sentinel-2, before (1st of September 2021), during (6th of September 2021) and after (8th of October 2021) a HW in the seagrass meadow of Quiberon (South Brittany, France). The relative SHSI difference was calculated using the 1st of September as a reference. SHSI was calculated for two categories of Quadrat Points (QPs; Figure 9): unimpacted seagrass (green) and impacted seagrass (orange). Points represent the estimated value of the SHSI using a GLM, while the error bar represents the 89% confidence interval.

Figure 11: Sentinel-2 color composition of the seagrass meadow of Quiberon, South Brittany, France, Before (A ; 1st of September 2021), During (B ; 6th of September 2021) and After (C ; 8th of October 2021) the HW and SHSI applied to the same Sentinel-2 images Before (D), During (E) and After (F) the HW.

Figure 12: Relative change of the SHSI before and during the HW events as a function of the daily emersion time of seagrass. The line represents a GAM estimate, and the shaded area indicates the standard error. Shaded points represent raw data, each corresponding to a single pixel of the meadow.

Figure A: Comparison of daily maximum temperatures in August measured using an in-situ sensor (blue) and retrieved from Meteo France (orange). The solid line in the middle of the boxplot represents the median, the two ends of the box represent the 25th

and 75th percentiles, and the whiskers represent values that are no more than 1.5 times the interquartile range.

Table Captions

Table B: Outputs of the Generalised Linear mixed effects model assessing the relationship between time of heatwave (days) and the relative change to the control of different vegetation indices (Figure 7, Second Derivative, NDVI, GLI, and SHSI). Each index has two terms: Intercept and Slope. The Intercept represents the expected value of the index at the reference time point (e.g., day 0), while the Slope quantifies the effect of time of heatwave, indicating how the index changes per unit increase in days. The Estimate represents the posterior mean of the regression coefficient, while Std. Error indicates its standard deviation. The Lower 95% CI and Upper 95% CI define the 95% credible interval, showing the range within which the true parameter is expected to fall with 95% probability. The Rhat statistic assesses model convergence (values close to 1 indicate proper convergence). Bulk_ESS and Tail_ESS denote the effective sample sizes for bulk estimation and tail uncertainty, respectively, reflecting the reliability of the posterior estimates.

Table C: Outputs of the Generalized Additive Model (GAM) assessing the relationship between emersion time per day (hours) and the relative change in SHSI (Figure 12). The

Intercept represents the expected value of the relative change of SHSI at the reference time point, while the $s(emersion_per_day)$ is a function of the effect of emersion time per day on the relative change of SHSI. The Estimate represents the posterior mean of the regression coefficient, while Std. Error indicates its standard deviation. The Lower 95% CI and Upper 95% CI define the 95% credible interval, showing the range within which the true parameter is expected to fall with 95% probability. The Rhat statistic assesses model convergence (values close to 1 indicate proper convergence). Bulk_ESS and Tail_ESS denote the effective sample sizes for bulk estimation and tail uncertainty, respectively, reflecting the reliability of the posterior estimates.

8. References

- Allakhverdiev, S.I., Kreslavski, V.D., Klimov, V.V., Los, D.A., Carpentier, R., Mohanty, P., 2008. Heat stress: An overview of molecular responses in photosynthesis. *Photosynthesis research* 98, 541–550.
- Arias-Ortiz, A., Serrano, O., Masqué, P., Lavery, P.S., Mueller, U., Kendrick, G.A., Rozaimi, M., Esteban, A., Fourqurean, J.W., Marbà, N., others, 2018. A marine heatwave drives massive losses from the world's largest seagrass carbon stocks. *Nature Climate Change* 8, 338–344.
- Armitage, A., Fourqurean, J.W., 2016. Carbon storage in seagrass soils: Long-term nutrient history exceeds the effects of near-term nutrient enrichment. *Biogeosciences* 13, 313–321.
- Bastos, M., Roebeling, P., Alves, F.L., Villasante, S., Magalhães Filho, L., 2023. High risk water pollution hazards affecting aveiro coastal lagoon (portugal)—a habitat risk assessment using InVEST. *Ecological Informatics* 76, 102144.
- Borum, J., Duarte, C.M., Krause-Jensen, D., Greve, T.M., 2004. European seagrasses: An introduction to monitoring and management. *Monitoring; Managing of European Seagrasses Project*.

- Boyer, M., Miller, J., Belanger, M., Hare, E., Wu, J., 1988. Senescence and spectral reflectance in leaves of northern pin oak (*quercus palustris muenchh.*). *Remote Sensing of Environment* 25, 71–87.
- Brun, F.G., Vergara, J.J., Navarro, G., Hernández, I., Pérez-Lloréns, J.L., 2003. Effect of shading by *ulva rigida* canopies on growth and carbon balance of the seagrass *zostera noltii*. *Marine Ecology Progress Series* 265, 85–96.
- Bürkner, P.-C., 2021. Bayesian item response modeling in r with brms and stan. *Journal of Statistical Software* 100, 1–54. <https://doi.org/10.18637/jss.v100.i05>
- Cabaço, S., Machás, R., Santos, R., 2009. Individual and population plasticity of the seagrass *zostera noltii* along a vertical intertidal gradient. *Estuarine, Coastal and Shelf Science* 82, 301–308.
- Calleja, M.L., Marbà, N., Duarte, C.M., 2007. The relationship between seagrass (*posidonia oceanica*) decline and sulfide porewater concentration in carbonate sediments. *Estuarine, Coastal and Shelf Science* 73, 583–588.
- Carlson, D.F., Yarbrow, L.A., Sclaro, S., Poniatowski, M., McGee-Absten, V., Carlson Jr, P.R., 2018. Sea surface temperatures and seagrass mortality in florida bay: Spatial and temporal patterns discerned from MODIS and AVHRR data. *Remote Sensing of Environment* 208, 171–188.

- Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M.,
Brubaker, M., Guo, J., Li, P., Riddell, A., 2017. Stan: A probabilistic programming
language. *Journal of Statistical Software* 76, 1–32. <https://doi.org/10.18637/jss.v076.i01>
- Climate Change (IPCC), I.P. on, 2023. Summary for policymakers. IPCC, Geneva,
Switzerland. <https://doi.org/10.59327/IPCC/AR6-9789291691647.001>
- CMEMS, 2024. European north west shelf/iberia biscay irish seas – high resolution
ODYSSEA sea surface temperature multi-sensor L3 observations reprocessed, e.u.
Copernicus marine service information (CMEMS). Marine data store (MDS). (Accessed
on 17-10-2024). <https://doi.org/10.48670/moi-00311>
- Dascaliuc, A., Ralea, T., Cuza, P., 2007. Influence of heat shock on chlorophyll
fluorescence of white oak (*quercus pubescens* willd.) leaves. *Photosynthetica* 45, 469–
471. <https://doi.org/10.1007/s11099-007-0084-3>
- Davies, B.F.R., Gernez, P., Geraud, A., Oiry, S., Rosa, P., Zoffoli, M.L., Barillé, L.,
2023. Multi-and hyperspectral classification of soft-bottom intertidal vegetation using a
spectral library for coastal biodiversity remote sensing. *Remote Sensing of Environment*
290, 113554.
- Davies, B.F.R., Oiry, S., Rosa, P., Zoffoli, M.L., Sousa, A.I., Thomas, O.R., Smale,
D.A., Austen, M.C., Biermann, L., Attrill, M.J., others, 2024a. A sentinel watching over

inter-tidal seagrass phenology across western europe and north africa. Communications
Earth & Environment 5, 382. <https://doi.org/10.1038/s43247-024-01543-z>

Davies, B.F.R., Oiry, S., Rosa, P., Zoffoli, M.L., Sousa, A.I., Thomas, O.R., Smale,
D.A., Austen, M.C., Biermann, L., Attrill, M.J., others, 2024b. Intertidal seagrass extent
from sentinel-2 time-series show distinct trajectories in western europe. Remote Sensing
of Environment 312, 114340. <https://doi.org/10.1016/j.rse.2024.114340>

Deguet, A., Barrote, I., Silva, J., 2022. Physiological and morphological effects of a
marine heatwave on the seagrass *Cymodocea nodosa*. Scientific Reports 12, 7950.

Douay, F., Verpoorter, C., Duong, G., Spilmont, N., Gevaert, F., 2022. New
hyperspectral procedure to discriminate intertidal macroalgae. Remote Sensing 14, 346.

Duarte, C.M., 1991. Allometric scaling of seagrass form and productivity. Marine
ecology progress series. Oldendorf 77, 289–300.

Electric Blue, 2023. Intertidal chamber.

European Space Agency, 2024b. Sen2Cor: Sentinel-2 atmospheric correction
processor.

European Space Agency, 2024a. Copernicus open access hub.

Folmer, E.O., Geest, M. van der, Jansen, E., Olff, H., Michael Anderson, T., Piersma, T., Gils, J.A. van, 2012. Seagrass–sediment feedback: An exploration using a non-recursive structural equation model. *Ecosystems* 15, 1380–1393.

Franssen, S.U., Gu, J., Winters, G., Huylmans, A.-K., Wienpahl, I., Sparwel, M., Coyer, J.A., Olsen, J.L., Reusch, T.B., Bornberg-Bauer, E., 2014. Genome-wide transcriptomic responses of the seagrasses *zostera marina* and *nanozostera noltii* under a simulated heatwave confirm functional types. *Marine Genomics* 15, 65–73.

Gacia, E., Granata, T., Duarte, C., 1999. An approach to measurement of particle flux and sediment retention within seagrass (*posidonia oceanica*) meadows. *Aquatic Botany* 65, 255–268.

Garbulsky, M.F., Peñuelas, J., Gamon, J., Inoue, Y., Filella, I., 2011. The photochemical reflectance index (PRI) and the remote sensing of leaf, canopy and ecosystem radiation use efficiencies: A review and meta-analysis. *Remote sensing of environment* 115, 281–297.

Gardner, R.C., Finlayson, C., 2018. Global wetland outlook: State of the world's wetlands and their services to people, in: Ramsar Convention Secretariat. pp. 2020–5.

- Gladstone-Gallagher, R.V., Hughes, R.W., Douglas, E.J., Pilditch, C.A., 2018. Biomass-dependent seagrass resilience to sediment eutrophication. *Journal of Experimental Marine Biology and Ecology* 501, 54–64.
- Green, A.E., Unsworth, R.K., Chadwick, M.A., Jones, P.J., 2021. Historical analysis exposes catastrophic seagrass loss for the united kingdom. *Frontiers in plant science* 12, 629962.
- Guerrero-Meseguer, L., Marín, A., Sanz-Lázaro, C., 2020. Heat wave intensity can vary the cumulative effects of multiple environmental stressors on *Posidonia oceanica* seedlings. *Marine Environmental Research* 159, 105001.
- Hemminga, M., 1998. The root/rhizome system of seagrasses: An asset and a burden. *Journal of Sea Research* 39, 183–196.
- Hobday, A.J., Alexander, L.V., Perkins, S.E., Smale, D.A., Straub, S.C., Oliver, E.C., Benthuyssen, J.A., Burrows, M.T., Donat, M.G., Feng, M., others, 2016. A hierarchical approach to defining marine heatwaves. *Progress in oceanography* 141, 227–238.
- Hobday, A.J., Oliver, E.C., Gupta, A.S., Benthuyssen, J.A., Burrows, M.T., Donat, M.G., Holbrook, N.J., Moore, P.J., Thomsen, M.S., Wernberg, T., others, 2018. Categorizing and naming marine heatwaves. *Oceanography* 31, 162–173.

- Infantes, E., Hoeks, S., Adams, M.P., Heide, T. van der, Katwijk, M.M. van, Bouma, T.J., 2022. Seagrass roots strongly reduce cliff erosion rates in sandy sediments. *Marine Ecology Progress Series* 700, 1–12.
- IOC, 2024. Intergovernmental oceanographic commission ; sea level monitoring station - le conquet, france (LECY).
- Jankowska, E., Michel, L.N., Lepoint, G., Włodarska-Kowalczyk, M., 2019. Stabilizing effects of seagrass meadows on coastal water benthic food webs. *Journal of Experimental Marine Biology and Ecology* 510, 54–63.
- Jones, R.B., Clayton-Greene, K.A., 1992. The role of photosynthesis and oxidative reactions in leaf blackening of protea neriifolia r. Br. leaves. *Scientia Horticulturae* 50, 137–145. [https://doi.org/10.1016/S0304-4238\(05\)80016-0](https://doi.org/10.1016/S0304-4238(05)80016-0)
- Knipling, E.B., 1970. Physical and physiological basis for the reflectance of visible and near-infrared radiation from vegetation. *Remote Sensing of Environment* 1, 155–159. [https://doi.org/10.1016/S0034-4257\(70\)80021-9](https://doi.org/10.1016/S0034-4257(70)80021-9)
- Lehnert, L.W., Meyer, H., Bendix, J., 2017. Hsdar: Manage, analyse and simulate hyperspectral data in r.
- Loarie, S.R., Duffy, P.B., Hamilton, H., Asner, G.P., Field, C.B., Ackerly, D.D., 2009. The velocity of climate change. *Nature* 462, 1052–1055.

- Louhaichi, M., Borman, M.M., Johnson, D.E., 2001. Spatially located platform and aerial photography for documentation of grazing impacts on wheat. *Geocarto International* 16, 65–70.
- Malvern Panalytical, 2023. RS3 software.
- Marbà, N., Duarte, C.M., 2010. Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality. *Global change biology* 16, 2366–2375.
- Mariën, B., Balzarolo, M., Dox, I., Leys, S., Lorène, M.J., Geron, C., Portillo-Estrada, M., AbdElgawad, H., Asard, H., Campioli, M., 2019. Detecting the onset of autumn leaf senescence in deciduous forest trees of the temperate zone. *New Phytologist* 224, 166–176.
- Massa, S., Arnaud-Haond, S., Pearson, G., Serrão, E., 2009. Temperature tolerance and survival of intertidal populations of the seagrass *Zostera noltii* (Hornemann) in southern Europe (Ria Formosa, Portugal). *Hydrobiologia* 619, 195–201.
- Moore, K.A., Jarvis, J.C., 2008. Environmental factors affecting recent summertime eelgrass diebacks in the lower Chesapeake Bay: Implications for long-term persistence. *Journal of Coastal Research* 135–147.
- Oiry, S., Davies, B.F.R., Sousa, A.I., Rosa, P., Zoffoli, M.L., Brunier, G., Gernez, P., Barillé, L., 2024. Discriminating seagrasses from green macroalgae in European

intertidal areas using high-resolution multispectral drone imagery. Remote Sensing 16.
<https://doi.org/10.3390/rs16234383>

Olmedo-Masat, O.M., Raffo, M.P., Rodríguez-Pérez, D., Arijón, M., Sánchez-Carnero, N., 2020. How far can we classify macroalgae remotely? An example using a new spectral library of species from the south west atlantic (argentine patagonia). Remote Sensing 12, 3870.

Peñuelas, J., Munné-Bosch, S., Llusià, J., Filella, I., 2004. Leaf reflectance and photo- and antioxidant protection in field-grown summer-stressed *phillyrea angustifolia*. Optical signals of oxidative stress? New Phytologist 162, 115–124.

Perkins, S.E., Alexander, L.V., 2013. On the measurement of heat waves. Journal of climate 26, 4500–4517.

R Core Team, 2023. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Ramesh, C., Mohanraju, R., 2020. Seagrass ecosystems of andaman and nicobar islands: Status and future perspective. Environmental & Earth Sciences Research Journal 7.

Roca, G., Alcoverro, T., Krause-Jensen, D., Balsby, T.J.S., Katwijk, M.M. van, Marbà, N., Santos, R., Arthur, R., Mascaró, O., Fernández-Torquemada, Y., others, 2016.

Response of seagrass indicators to shifts in environmental stressors: A global review
and management synthesis. *Ecological Indicators* 63, 310–323.

Román, A., Tovar-Sánchez, A., Olivé, I., Navarro, G., 2021. Using a UAV-mounted
multispectral camera for the monitoring of marine macrophytes. *Frontiers in Marine
Science* 8, 722698.

Román, M., Gilbert, F., Viejo, R.M., Román, S., Troncoso, J.S., Vázquez, E.,
Olabarria, C., 2023. Are clam-seagrass interactions affected by heatwaves during
emersion? *Marine Environmental Research* 186, 105906.

Román, S., Vázquez, E., Román, M., Viejo, R.M., Woodin, S.A., Wetthey, D.S.,
Troncoso, J.S., Olabarria, C., 2022. Effects of warming on biological interactions between
clams and the seagrass *zostera noltei*: A case study using open top chambers. *Estuarine,
Coastal and Shelf Science* 276, 108027.

Rouse, J.W., Haas, R.H., Schell, J.A., Deering, D.W., others, 1974. Monitoring
vegetation systems in the great plains with ERTS. *NASA Spec. Publ* 351, 309.

Samper-Villarreal, J., Bolaños, R.C., Heidemeyer, M., Vargas, M.M., Vargas, R.M.,
2020. Characterization of seagrasses at two new locations in the eastern tropical pacific
(el jobo and matapalito, costa rica). *Aquatic botany* 165, 103237.

Sawall, Y., Ito, M., Pansch, C., 2021. Chronically elevated sea surface temperatures revealed high susceptibility of the eelgrass *zostera marina* to winter and spring warming. *Limnology and Oceanography* 66, 4112–4124.

Schlegel, R.W., Smit, A.J., 2018. heatwaveR: A central algorithm for the detection of heatwaves and cold-spells. *Journal of Open Source Software* 3, 821. <https://doi.org/10.21105/joss.00821>

Scott, A.L., York, P.H., Duncan, C., Macreadie, P.I., Connolly, R.M., Ellis, M.T., Jarvis, J.C., Jinks, K.I., Marsh, H., Rasheed, M.A., 2018. The role of herbivory in structuring tropical seagrass ecosystem service delivery. *Frontiers in Plant Science* 9, 127.

SHOM, 2022. Service hydrographique et océanographique de la marine ; références altimétriques maritimes: Ports de france métropolitaine et d'outre-mer, cotes du zéro hydrographique et niveaux caractéristiques de la marée. Shom, Brest, France.

SHOM, 2021. Service hydrographique et océanographique de la marine ; bathymétrie Litto3D® bretagne 2018-2021.

Skendzic, S., 2023. Drought stress in winter wheat-physiological responses and detection using remote and proximal sensing techniques, in: *Smart Life Sciences and Technology for Sustainable Development*. pp. 39–39.

Sousa, A.I., Silva, J.F. da, Azevedo, A., Lillebø, A.I., 2019. Blue carbon stock in
zostera noltei meadows at ria de aveiro coastal lagoon (portugal) over a decade.
Scientific reports 9, 14387.

Stan Development Team, C., others, 2020. RStan: The r interface to stan. R package
version 2.21. 2.

Stillman, J.H., 2019. Heat waves, the new normal: Summertime temperature
extremes will impact animals, ecosystems, and human communities. Physiology 34, 86–
100.

Strydom, S., Murray, K., Wilson, S., Huntley, B., Rule, M., Heithaus, M., Bessey, C.,
Kendrick, G.A., Burkholder, D., Fraser, M.W., others, 2020. Too hot to handle:
Unprecedented seagrass death driven by marine heatwave in a world heritage area.
Global change biology 26, 3525–3538.

Thomsen, E., Herbeck, L.S., Viana, I.G., Jennerjahn, T.C., 2023. Meadow trophic
status regulates the nitrogen filter function of tropical seagrasses in seasonally eutrophic
coastal waters. Limnology and Oceanography 68, 1906–1919.

Thomson, J.A., Burkholder, D.A., Heithaus, M.R., Fourqurean, J.W., Fraser, M.W.,
Statton, J., Kendrick, G.A., 2015. Extreme temperatures, foundation species, and abrupt

ecosystem change: An example from an iconic seagrass ecosystem. *Global change biology* 21, 1463–1474.

Unsworth, R., Cullen-Unsworth, L.C., 2014. Biodiversity, ecosystem services, and the conservation of seagrass meadows. *Coast. Conserv* 19, 95.

Unsworth, R.K., Butterworth, E.G., 2021. Seagrass meadows provide a significant resource in support of avifauna. *Diversity* 13, 363.

Unsworth, R.K., Cullen-Unsworth, L.C., Jones, B.L., Lilley, R.J., 2022. The planetary role of seagrass conservation. *Science* 377, 609–613.

Ustin, S.L., Jacquemoud, S., 2020. How the optical properties of leaves modify the absorption and scattering of energy and enhance leaf functionality. *Remote sensing of plant biodiversity* 349–384.

Waycott, M., Duarte, C.M., Carruthers, T.J., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A., Fourqurean, J.W., Heck Jr, K.L., Hughes, A.R., others, 2009. Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceedings of the national academy of sciences* 106, 12377–12381.


Winters, G., Nelle, P., Fricke, B., Rauch, G., Reusch, T.B., 2011. Effects of a simulated heat wave on photophysiology and gene expression of high-and low-latitude populations of *zostera marina*. *Marine Ecology Progress Series* 435, 83–95.


- Zahoor, I., Mushtaq, A., 2023. Water pollution from agricultural activities: A critical global review. *Int. J. Chem. Biochem. Sci* 23, 164–176.
- Zoffoli, M.L., Gernez, P., Godet, L., Peters, S., Oiry, S., Barillé, L., 2021. Decadal increase in the ecological status of a north-atlantic intertidal seagrass meadow observed with multi-mission satellite time-series. *Ecological Indicators* 130, 108033.
- Zoffoli, M.L., Gernez, P., Oiry, S., Godet, L., Dalloyau, S., Davies, B.F.R., Barillé, L., 2023. Remote sensing in seagrass ecology: Coupled dynamics between migratory herbivorous birds and intertidal meadows observed by satellite during four decades. *Remote Sensing in Ecology and Conservation* 9, 420–433.
- <https://doi.org/10.1002/rse2.319>
- Zoffoli, M.L., Gernez, P., Rosa, P., Le Bris, A., Brando, V.E., Barillé, A.-L., Harin, N., Peters, S., Poser, K., Spaias, L., others, 2020. Sentinel-2 remote sensing of zostera noltei-dominated intertidal seagrass meadows. *Remote Sensing of Environment* 251, 112020.




Figure2


[Click here to access/download;Figure;Figure2.png](#)














Figure8

[Click here to access/download;Figure;Figure8.png](#)




Figure9

[Click here to access/download;Figure;Figure9.png](#)




Figure10






Figure11


[Click here to access/download;Figure;Figure11.png](#)







Click here to access/download
Supplementary Material
FigureA.png






Click here to access/download
Supplementary Material
TableB.docx



Click here to access/download
Supplementary Material
TableC.docx



Declaration of interests

☒The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: