

# Effect of Marine and Atmospheric Heatwaves on Reflectance and Pigment Composition of Intertidal *Nanozostera noltei*

Simon Oiry<sup>1</sup>, Bede Ffinian Rowe Davies<sup>1</sup>, Philippe Rosa<sup>1</sup>, Augustin Debly<sup>1</sup>, Maria Laura Zoffoli<sup>2</sup>, Anne-Laure Barillé<sup>3</sup>, Nicolas Harin<sup>3</sup>, Pierre Gernez<sup>1</sup>, and Laurent Barillé<sup>1</sup>

<sup>1</sup>Institut des Substances et Organismes de la Mer, ISOMer, Nantes Université, UR 2160, F-44000 Nantes, France

<sup>2</sup>Consiglio Nazionale delle Ricerche, Istituto di Scienze Marine (CNR-ISMAR), 00133 Rome, Italy

<sup>3</sup>Bio-littoral, Immeuble Le Nevada, 2 Rue du Château de l'Eraudière, 44300 Nantes, France

## Abstract

Seagrasses are critical to coastal ecosystems, providing habitat, stabilizing sediments, and aiding carbon sequestration. Climate change has increased the frequency and intensity of heatwaves, potentially threatening seagrass health. This study investigates the impact of marine and atmospheric heatwaves on the pigment composition and reflectance of the intertidal seagrass *Nanozostera noltei*. We performed laboratory experiments, exposing *N. noltei* samples to controlled heatwave conditions and measured hyperspectral reflectance and pigment concentration to assess its impact over time. Results revealed that heatwaves induce significant declines in seagrass reflectance, particularly in the green and near-infrared regions, linked to (likely due to) pigment degradation. Key vegetation indices, such as the Normalized Difference Vegetation Index (NDVI) and Green Leaf Index (GLI), also displayed marked reductions under heatwave stress, with NDVI values decreasing by up to 34% and GLI by 57%. A novel Seagrass Darkening Index (SDI) was developed to identify seagrass darkening, showing a strong correlation with heatwave exposure. This research suggests that spectral monitoring can effectively track the early impacts of heatwaves on seagrasses, providing a valuable tool for remote sensing-based habitat assessment. Satellite observations confirmed these findings, showing widespread seagrass darkening during atmospheric and marine heatwave events in Quiberon, France. Darkened seagrasses observed after heatwaves were exposed more than 13.5 hours daily. This work highlights the need for continuous monitoring of seagrass meadows under the current climate regime, underscoring the potential of remote sensing in capturing rapid environmental changes in intertidal zones.

**Keywords:** Remote Sensing, Pigment Composition, Seagrass, Coastal Ecosystems, Heatwaves

## 1 Introduction

Intertidal seagrasses play a crucial role in the ecosystem by providing habitats and feeding grounds for various marine species, supporting rich marine biodiversity, and contributing significantly to primary production and carbon sequestration (Sousa et al., 2019; Unsworth et al., 2022). These seagrasses are es-

5 sential in maintaining the health of coastal ecosystems by stabilizing sediments, filtering water, and serv-  
 6 ing as indicators of environmental changes due to their sensitivity to water quality variations (Zoffoli et al.,  
 7 2021). The interactions between seagrass meadows and their associated herbivores further enhance the deliv-  
 8 ery of ecosystem services, including coastal protection and fisheries support (Gardner and Finlayson, 2018;  
 9 Jankowska et al., 2019; Zoffoli et al., 2023). Understanding and preserving these ecosystems are vital for  
 10 maintaining the biodiversity and productivity of coastal regions (Ramesh and Mohanraju, 2020; Scott et al.,  
 11 2018).

12 Despite their crucial role in marine ecosystems, intertidal seagrasses face numerous threats that com-  
 13 promise their health and functionality. Coastal development and human activities are primary threats. These  
 14 activities not only reduce the available habitat for seagrasses but also increase water turbidity, which lim-  
 15 its light penetration and hampers photosynthesis (Waycott et al., 2009). Seagrasses are also threatened by  
 16 nutrient enrichment from agricultural and urban runoff, which can lead to eutrophication. This condition pro-  
 17 motes the overgrowth of algal blooms that compete with seagrasses for light and nutrients, further stressing  
 18 these important plants (Thomsen et al., 2023) (Oiry et al. 2024). Pollution from industrial and agricultural  
 19 fields sources introduces harmful chemicals and heavy metals into coastal waters, posing toxic risks to sea-  
 20 grass health. These pollutants can affect the physiological processes of seagrasses, reducing their growth and  
 21 survival rates (Sevgi and Leblebici, 2022). Additionally, invasive species can out compete native seagrasses  
 22 for resources, altering community structure and function (Simpson et al., 2016).

23 Heatwaves, exacerbated by climate change, pose a growing threat to seagrasses. Marine Heatwaves  
 24 (MHW), defined by Hobday et al. (2016) as prolonged discrete anomalously warm water events, and At-  
 25 mospheric Heatwaves (AHW), defined by Perkins and Alexander (2013) as periods of at least three consec-  
 26 utive days with temperatures exceeding the 90th percentile, cause severe physiological stress on seagrasses  
 27 (Deguette et al., 2022; Sawall et al., 2021). At the interface between the land and oceans, intertidal sea-  
 28 grasses are exposed to both MHW and AHW. Heatwaves have profound impacts on seagrasses, with their  
 29 effects varying based on species and geographic location. For instance, the seagrass *Zostera marina* ex-  
 30 hibits high susceptibility to elevated sea surface temperatures during winter and spring, leading to advanced  
 31 flowering, high mortality rates, and reduced biomass (Sawall et al., 2021). Similarly, *Cymodocea nodosa*  
 32 shows increased photosynthetic activity during heatwaves but suffers negative effects on photosynthetic per-  
 33 formance and leaf biomass during recovery (Deguette et al., 2022). Additionally, different populations of  
 34 *Zostera marina* along the European thermal gradient exhibit varied photophysiological responses during the  
 35 recovery phase of heatwaves, indicating differential adaptation capabilities among populations (Winters et  
 36 al., 2011). These events intensify other stressors, such as overgrazing and seed burial, compromising sexual  
 37 recruitment (Guerrero-Meseguer et al., 2020).

38 Remote sensing is increasingly being used to monitor marine ecosystems, including seagrass mead-  
 39 ows. Spectral indices such as the Normalized Difference Vegetation Index (NDVI) and the Soil-Adjusted

---

Simon Oiry  <https://orcid.org/0000-0001-7161-5246>

Bede Ffinian Rowe Davies  <https://orcid.org/0000-0001-6462-4347>

Philippe Rosa  <https://orcid.org/0000-0001-6621-3445>

Augustin Debly  <https://orcid.org/0009-0000-7194-7099>

Maria Laura Zoffoli  <https://orcid.org/0000-0003-1892-0051>

Pierre Gernez  <https://orcid.org/0000-0003-2055-410X>

Laurent Barillé  <https://orcid.org/0000-0001-5138-2684>

Author roles were classified using the Contributor Role Taxonomy (CRediT; <https://credit.niso.org/>) as follows: Simon Oiry: Investigation, Writing, Data acquisition; Bede Ffinian Rowe Davies: Investigation, Data acquisition, Revision; Philippe Rosa: Data acquisition, Revision; Augustin Debly: Investigation, Data acquisition, Revision; Maria Laura Zoffoli: Data acquisition, Revision; Anne-Laure Barillé: Investigation, Data acquisition, Revision; Nicolas Harin: Investigation, Data acquisition, Revision; Pierre Gernez: Investigation, Data acquisition, Revision; Laurent Barillé: Investigation, Data acquisition, Revision

Vegetation Index (SAVI) are effective for quantifying vegetation health over time (Akbar et al., 2020; Câr-lan et al., 2020; Huete, 2012; Kloos et al., 2021). By analyzing specific spectral patterns, remote sensing can track changes in seagrass meadows with high temporal and spatial resolution. Europe promotes these techniques through frameworks like the Water Framework Directive and the Marine Strategy Framework Directive, which advocate for habitat mapping using remote sensing technologies to cover large areas and detect long-term trends (Papathanasopoulou et al., 2019).

This study aims to experimentally test the hypothesis that warm events, such as atmospheric heat-waves, alter the pigment composition and reflectance of the intertidal seagrass *Nanozostera noltei*. By linking these changes with satellite remote sensing data, the study seeks to provide insights into how heatwaves affect seagrass meadows and to assess the potential of remote sensing in monitoring these impacts

## 2 Material & Methods

### 2.1 Laboratory Experiment


#### 2.1.1 Sampling and acclimation of seagrasses

Seagrass was sampled from a *Nanozostera noltei* (dwarf eelgrass, syn. *Zostera noltei*) meadow on Noirmoutier Island, France (46°57'32.0"N, 2°10'37.0"W) at low tide in June 2024. A home-made inox sampling box was used to sample seagrass from an area of 30 cm by 15 cm and 5 cm deep, maintaining the sediment structure and avoiding damage to the rhizomes and the leafs of the seagrass (Figure 1 A). This sampling box allowed to limitate sampling variability between replicates. The entire system, including seagrass, sediment, meiofauna, and macrofauna, was placed in plastic trays together. Keeping the entire mesocosm allowed for the natural interactions between components to be maintained and reduced stress on the seagrass. To avoid hydric stress during transportation, seawater was added to each tray. Simultaneously, seawater was sampled from a nearby site and transported to the lab, where it was filtered using a 0.22 µm nitrocellulose filter to remove all suspended particulate matter. This seawater was used in the acclimation tank and the intertidal chambers. The seagrasses were acclimated at high tide for one weeks with a water temperature of 17°C, matching the temperature at the time of sampling, and with light of 150 µmol.s-1.m-2 of PAR photons (Akbar et al., 2020). A wave generator was used in the tank to circulate and reoxygenate the water.



**Figure 1**

*Illustrations of the various steps of the experiment. A: Field sampling of seagrass using a homemade sampling box; B: Intertidal chambers used during the experiment; C: Seagrass sample inside a chamber during the experiment at high tide; D: Photo of the treatment sample at the start of the experiment; E: Photo of the treatment sample at the end of the experiment.*




### 67 2.1.2 Experimental design

68 Two intertidal chambers from [ElectricBlue](#) were used to simulate tidal cycles and control water  
 69 temperature during high tide and air temperature during low tide (Figure 1 B,C). One chamber served as the  
 70 control, while the other was used for the experimental treatment.



**Figure 2**

Comparison of daily maximum temperatures in August measured using an *in-situ* sensor (blue) and retrieved from Meteo France (orange). The solid line in the middle of the boxplot represents the median, the two ends of the box represent the 25th and 75th percentiles, and the whiskers represent values that are no more than 1.5 times the interquartile range.




To closely replicate field temperatures experienced by seagrasses during the experiment, *in situ* temperature sensors were installed at the sampling site in Bourgneuf Bay in August 2024. Daily maximum temperatures recorded by these *in situ* sensors were compared to measurements from the nearest Meteo France weather station (Figure 2). On average, *in situ* temperatures were  $3 \pm 3.2^{\circ}\text{C}$  higher than those recorded by Meteo France. Additionally, temperatures recorded by Meteo France were more stable than those from the *in situ* sensors, likely due to the sheltered and shaded location of the Meteo France equipment. This difference was used to adjust heatwave temperatures measured by Meteo France to better reflect the conditions experienced by the seagrasses.

The control chamber was kept at temperatures representing typical seasonal conditions, with water temperatures ranging from  $18^{\circ}\text{C}$  to  $19^{\circ}\text{C}$  and air temperatures from  $18^{\circ}\text{C}$  to  $23^{\circ}\text{C}$ , following natural circadian temperature fluctuations (Figure 3 left). For the experimental treatment, air temperature was adjusted to replicate an atmospheric heatwave that affected the seagrass meadow in Quiberon, France ( $47^{\circ}35'40.0''\text{N}$ ,  $3^{\circ}07'30.0''\text{W}$ ), from September 2 to September 6, 2021. On the first day of the experiment, air temperatures

in the experimental chamber were set to range from 23°C at night to 35°C during the day, increasing by 1°C daily. Water temperature in the experimental chamber was also adjusted to mimic heatwave conditions, starting at the seasonal baseline (18°C) and rising incrementally by 0.5°C daily to simulate the increasing temperatures during the heatwave. This setup aimed to replicate the thermal stress experienced by the seagrass meadow during the actual heatwave event (Figure 3 right). The experiment concluded only when no changes in the reflectance of the treatment were observed for two consecutive days.

**Figure 3**

Temperature profiles of both the control (left) and the treatment (right) followed during the heatwave experiment. The red line indicates air temperature, whereas the blue line indicates water temperature. Due to the tidal cycle followed during the experiment, the seagrasses only experience temperatures represented by solid lines.



### 2.1.3 Bio-optical measurements over seagrass leaves

**2.1.3.1 Hyperspectral measurements.** Throughout the experiment, hyperspectral signatures of both the control and treatment seagrasses were taken using an ASD HandHeld 2 equipped with a fiber optic, allowing measurements to be taken directly inside the chamber without opening it. Automatic spectra acquisition has been done using the [RS3 software](#) developed by the instrument manufacturer. An average of five reflectance spectrum ( $R(\lambda)$ ), each with an integration time of 544 ms, was taken every minute. Every 10 minutes, the fiber optic was switched from one benthic chamber to the other, in order to measure reflectance in both treatment and control. Because light conditions were controlled inside of the chambers, reflectance calibration of the instrument was performed only each morning at the very first moment of low tide using a Spectralon white reference with 99% Lambertian reflectivity.

The second derivative of  $R$  was calculated to retrieve absorption features and compare their variability over time. Two radiometric indices were also monitored throughout the experiment :

- The Normalized Difference Vegetation Index (NDVI, Rouse et al. (1974)), as a proxy of the concentration of chlorophyll-a (Equation 1)

$$NDVI = \frac{R(840) - R(668)}{R(840) + R(668)} \quad (1)$$

where  $R(840)$  and  $R(668)$  are the reflectance at 840 nm and 668 nm respectively.

- The Seagrass Percent Covers, developed by Zoffoli et al. (2020), used to assess proportion of seagrass inside of a pixel (Equation 2):

$$SPC = 172.06 \times NDVI - 22.18 \quad (2)$$

- The Green Leaf Index (GLI, Louhaichi et al. (2001)), as a measurement of the greenness of seagrass leafs (Equation 3)

$$GLI = \frac{[R(550) - R(668)] + [R(550) - R(450)]}{(2 \times R(550)) + R(668) + R(450)} \quad (3)$$

where  $R(550)$  and  $R(450)$  are the reflectance in green at 550 nm and in the blue at 450 nm, respectively.

**2.1.3.2 WILL BE REMOVED - Multispectral imagery measurement.** Parallel to hyperspectral measurements, multispectral images were taken at the beginning and end of each diurnal low tide (09:00 am and 03:00 pm). A Micasense RedEdge-MX Dual multispectral camera, originally designed to be mounted on a drone, was modified for use without a drone. A 3D-printed mount was designed to attach the camera to the intertidal chamber and ensure that each picture was captured under the same conditions. At each time step (09:00 am and 03:00 pm), a first picture of the Spectralon was taken to allow for image correction in reflectance, followed by a second picture of the target. DISCOV, a Neural Network classification model previously developed to map intertidal vegetation using drone imagery, has been applied to each image taken inside the intertidal chambers. To understand the behavior of the model on seagrasses affected by heatwaves, classification images from before and after the heatwave have been compared.

**2.1.3.3 Pigment concentration measurements.** At the beginning and the end of each diurnal low tide (09:00 am and 03:00 pm) leaves samples have been took in both the test and the control. leaves sampled have been stored under -80°C waiting for analysis. Before pigment extraction, each sample was lyophilized and its dry weight was measured. Pigment composition and biomass were analyzed using high-performance liquid chromatography (HPLC). The HPLC system (Alliance HPLC 248 System, Waters) was equipped with a reverse-phase C-18 separating column (SunFire C-18 Column, 100Å, 3.5 µm, 2.1 mm x 50 mm, Waters), preceded by a precolumn (VanGuard 3.9 mm x 5 mm, Waters). The system also included a photodiode array detector (2998 PDA) and a fluorimeter (Ex: 425 nm, Em: 655 nm; RF-20A, SH

**Au secours Philippe !!!**




## 2.2 Observation of seagrass leaves darkening.

### 2.2.1 Field Observation

**Figure 4**

*Location of the fieldtrip campaign that occurred in the 10th of September 2021. The light grey polygons indicates the intertidal zone (Zone between High tide and low tide, that is totally emerged at low tide) and the dark grey polygon indicate the extent of the drone flight. Green dots indicate location of quadrat picture over green seagrasses while red dots indicate location of quadrat took over darkened seagrasses.*




A fieldtrip, aiming to map a seagrass meadow near Quiberon (France : 46°57'32.0"N, 2°10'37.0"W), occurred in the 10th of September 2021 (Figure 4). During this fieldtrip, significant darkening of seagrass



leaves was observed over large areas of the meadow (Figure 5 C & D). A total of 122 Ground Control Points (GCPs) were collected in the form of georeferenced quadrat images across the meadow. These images allow for visual assessment of vegetation type, density, and health status. The images were then divided into two categories: green seagrasses and darkened seagrasses, based on a visual estimation of the leaf condition (Figure 4).

**Figure 5**

*Illustrations of the two health conditions of seagrasses observed in the field. A: Global view of a green green meadow; B: Quadrat images of green seagrasses; C: Quadrat images of darkened seagrasses; D: Global view of an darkened meadow. All images were taken on September 10th, 2021, in Quiberon.*



## 2.2.2 Temperature Data and identification of heatwaves

**2.2.2.1 Air temperature.** Since January 1, 2024, Météo France weather data has been freely and openly accessible. Hourly air temperature data (°C) for the French Atlantic and Channel coasts was retrieved using a [custom script](#) as no API was available at the time of this study. Weather stations located within 10 kilometers of the coastline were considered, but only those with at least 30 years of data were included to ensure reliable climatological reconstruction. Of the 156 weather stations within 10 kilometers of the coast, only 36 had sufficient data for climatology reconstruction. The hourly data was then aggregated into daily mean temperatures for each station.

Heatwave detection was performed using the HeatwaveR package in R ([Schlegel and Smit, 2018](#)). This package utilizes the methodology proposed by Hobday et al. (2016) to detect heatwave events. The climatology for the year was computed using the temperature time series. An event was considered a heatwave each time the temperature exceeded the 90th percentile of the climatology for three consecutive days. The severity of each event has been assessed using the methodology proposed by Hobday et al. (2018). Fol-

lowing the methodology of Schlegel et al. (2017), coldspell events corresponding to temperature bellow the percentile 10th of the climatology during 3 consecutives days have also been measured.

Concerning heatwaves event near Quiberon, this methodology has been apply solely on the nearest weather station of our study site (Lorient-Lann Bihoue, 47°45'46"N 3°26'11"W , more than 395000 observation since the 1st of February 1952).

**2.2.2.2 Water temperature.** SST data from the Copernicus CMEMS platform were downloaded for the French coast near Quiberon (47°29 03 N, 3°07 09 W), covering 1982-2022 (CMEMS, 2024). Only pixels within an area of 2700 km<sup>2</sup> around Quiberon, Brittany, France (47°29 03 N, 3°07 09 W) were extracted and analyzed. This area was large enough to minimize missing values caused by cloud cover, yet small enough to avoid being influenced by the stability of offshore SST. The multi-sensor Level 3S Sea Surface Temperature (SST) product that we used is built from nighttime infrared satellite data, including sources like ESA SST CCI, C3S, and EUMETSAT. Data from AVHRR, ATSR, and SLSTR radiometers are used, but only those with a quality level above 4. An intercalibration method ensures consistency, and a daily reference SST field is constructed by combining the best data with median values from other sources. A large-scale bias correction is applied, and daily single-sensor composites are merged into a multi-sensor file (L3S), selecting the best sensor for each grid cell. A daily average was computed for the time series, and the HeatwaveR package (Schlegel and Smit, 2018) was used to compute SST climatology and detect SST events using the same method as air temperature event detection.

### 2.2.3 Satellite Observation

Two Sentinel-2 images covering the field trip site (Figure 4) were downloaded from the Copernicus platform (European Space Agency, 2024a). The first image was taken before the heatwave event on September 1, 2021, and the second one after the event on September 6, 2021. Both images were acquired at Level-2 processing, in surface reflectance, and were orthorectified and atmospherically corrected to account for the effects of the atmosphere on reflectance values (European Space Agency, 2024b).

Reflectance value of each GCPs (Figure 4) have been extracted and compared before and after the heatwave event.

### 2.2.4 Emersion Time of Seagrasses

To estimate the emersion time of seagrasses during low tide, bathymetric and tidal data were used. Bathymetry data for the Quiberon intertidal meadow were sourced from the “Service Hydrographique et Océanographique de la Marine” (SHOM, 2021). It corresponds to the Litto3D® product, which is a three-dimensional land-sea elevation database, accurately depicting the coastal terrain along certain regions of the French shores. Litto3D data were collected through airborne bathymetric lidar over the sea and airborne topographic lidar extending up to 2 km inland and is available as a gridded digital terrain model. it uses the NGF/IGN69 reference “zero”, which corresponds to the mean sea level recorded at the Marseille tide gauge between 1885 and 1897, commonly known as the “Terrestrial Altimetric Zero”.

Tidal data during the heatwave event were downloaded from the Intergovernmental Oceanographic Commission data portal (IOC, 2024), using measurements from the nearest tide gauge at Le Conquet. In this dataset, the reference “zero” corresponds to the lowest astronomical tide, also known as the Hydrographic Zero.

Before calculating the emersion time, both datasets were intercalibrated to a common altitude reference. This involved applying a correction factor to the Litto3D data to align it with the Hydrographic zero. SHOM annually publishes a document called “Références Altimétriques Marines” (RAM, Shom, 2022), which provides these correction factors for each seaport along the French coastline. For Le Conquet port data from 2022, the correction factor was 2.85 m.

Once aligned, the corrected elevation was compared to water height for each pixel and each time step during the heatwave event. The emersion time was then calculated as the daily total time each pixel remained exposed.

## 3 Results

### 3.1 Laboratory Experiment

#### 3.1.1 Spectral inspection


Hyperspectral signatures were acquired throughout the experiment for both the Control and Treatment groups. In the Control group, reflectance remained relatively stable over time, with only minimal changes in magnitude and spectral features. Slight variations in spectral shape observed in this group were attributed to differing levels of water evaporation during low tides. Immediately after the water level receded from high to low tide, samples were very wet but dried progressively, reaching maximum dryness just before the tide rose again. This fluctuation in water content induced variations in the reflectance spectra. Overall, the Control group's reflectance spectra displayed a peak at 560 nm (green region), low values at 665 nm (indicative of strong chlorophyll-a absorption), and a high plateau in the near-infrared (NIR), beyond 705 nm (Figure 6, left).

In contrast, the Treatment group showed more pronounced changes in reflectance throughout the experiment. On the first day (T0), reflectance values were comparable to those in the Control group, especially in the visible region, with a notable peak around 560 nm and a pronounced trough at 665 nm. However, starting from T24 (the second day), reflectance began to decrease across all wavelengths, particularly around 560 nm and in the NIR. Around T48, the NIR reflectance appeared to stabilize at values like those observed at T24. In the green region, however, reflectance continued to decline slightly until the experiment's conclusion. Additionally, a slight increase in reflectance at 665 nm between T24 and T48 was observed, possibly associated with a reduction in chlorophyll-a concentration (Figure 6, right).



**Figure 6**

*Hyperspectral Reflectance signature collected during the three days of the experiment for both the Control (Left) and the Treatment (Right). A min-max standardization has been applied to each individual spectrum. Temperature profiles applied to both scenarios are shown in Figure 3.*



### 3.1.2 Radiometric indices comparison

The second derivative of the spectral signature for both Control and Treatment groups was computed, highlighting key absorption features. Figure 7 A shows the average second derivative for the Control group (top) and the Treatment group (bottom) over the experiment's duration. Clear differences emerge between the two groups: the Control group exhibits more variability, reflecting more pronounced spectral features, whereas the Treatment group shows less variability, indicating fewer absorption features in its spectral signature. The maximum absorption for both groups occurs at 665 nm, with values of  $1.63 \times 10^{-4}$  for the Control group and  $8.08 \times 10^{-5}$  for the Treatment group—a reduction of 50.31% by the experiment's end.

Monitoring the second derivative at 665 nm ( $R''_{665 \text{ nm}}$ ) over time reveals a rapid drop in replicates 1 and 3, decreasing by 26 and 54 % respectively after 24 hours (T24) of treatment (Figure 7, B). After T24  $R''_{665 \text{ nm}}$  in these replicates stabilizes for the remainder of the experiment. In contrast, replicate 2 shows a slower but more substantial decrease of 110 % by the experiment's end. Overall, between the start and the end of the experiment, the  $R''_{665 \text{ nm}}$  have decreased by 72 %.


NDVI shows the same trends, exhibit an initial rapid decline of approximately 25% between T24 and T30, after which their values stabilize, with minimal further changes observed through the end of the experiment. In contrast, Replicate 2 remains relatively stable between 0 and 48 hours, showing little change during this period (drop of 9% at T24). However, after T48, Replicate 2 undergoes a substantial drop, ultimately reaching a reduction of approximately 48% (Figure 7, C).

The relative GLI shows similar trends across the replicates. Replicates 1 and 3 exhibit an initial rapid decline of approximately 40 and 57 %, respectively between T24 and T30, after which their values stabilize with minimal further changes observed until the end of the experiment. In contrast, Replicate 2 remains relatively constant between 0 and 48 hours, showing only a minor decline of around 5% by T24. However, after T48 mark, Replicate 2 experiences a substantial decrease, ultimately reaching a reduction of

245 73% (Figure 7, D).

**Figure 7**

*Second Derivative of the Control and the Treatment (A) and evolution of the second derivative (B), the NDVI (C) and the GLI (D) over Time. A: Mean and standard deviation of the second derivative for both Control and Treatment groups on the final day of the experiment; B: Relative evolution of the second derivative at 665 nm between Treatment and Control groups over time across three replicates. Points indicate raw data, the line represents a generalized linear model, and the shaded area the model's standard error; C: Relative evolution of NDVI for the Treatment group compared to the Control group over time across three replicates. Points indicate raw data, the line represents a generalized linear model, and the shaded area the model's standard error; D: Relative evolution of GLI for the Treatment group relative to the Control group over time across three replicates. Points indicate raw data, the line represents a generalized linear model, and the shaded area the model's standard error.*





### 3.1.3 Buidling the Seagrass Darkening radiometric index

Based on the observed spectral changes in seagrasses exposed to extreme events, a radiometric index can be developed to detect seagrass darkening events through remote sensing. This darkening is characterized by a reduction in reflectance within the green region, along with a substantial decrease in the red-edge portion of the electromagnetic spectrum. (Figure 7).


The Seagrass Darkening Index (SDI) is introduced here as a measure of the difference between the interpolated value between 560 and 842 nm and the observed value at 740 nm. This approach calculates the line height using a triplet of bands, where 560 nm and 842 nm serve as the ‘baseline’ and 740 nm represents the central peak. The SDI can be mathematically expressed as:

$$SDI = [R(560) + 0.64 \cdot (R(842) - R(560))] - R(740) \quad (4)$$

Where  $R(560)$ ,  $R(740)$ , and  $R(842)$  represent reflectance values at 560, 740, and 842 nm, respectively. These wavelengths were selected to align with the spectral resolution of satellites like Sentinel-2, enabling the index to be used for mapping darkening events from satellite data. From SDI, negative values indicate green, healthy seagrasses, while positive values correspond to darkened seagrasses.

**Figure 8**

*Development and Application of the Seagrass Darkening Index (SDI). The left panel displays the average reflectance values for both the Control and Treatment groups on the final day of the experiment. Shaded areas represent the standard deviation of reflectance, while markers indicate the spectral bands used to construct the SDI. The right panel shows the application of the SDI for both groups on the last day of the experiment.*



When applied to reflectance data collected during the experiment, the SDI revealed clear distinctions between the Control and Treatment groups. By the end of the experiment, seagrasses in the Treatment group exhibited a median SDI of 0.01, with no samples showing a negative SDI value, consistent with their visibly darkened appearance. In contrast, the Control group retained a green appearance throughout the experiment, with a median SDI of -0.003, and only 3% of samples displaying a positive SDI value (Figure 8, right).

### 3.1.4 *Pigment composition evolution along the experiment*

bla bla bla bla


## 3.2 Heatwaves of Quiberon

### 3.2.1 *Spectral inspection*

The Sentinel-2 images used in this study are dated from the 1st of September 2021 and the 6th of September 2021 (Figure 9 A and C, respectively). The Atmospheric Heat Wave (AHW) started on 4 September and lasted until 7 September, while the Marine Heat Wave (MHW) started on 3 September and ended on 8 September 2021 (Figure 9 B). The AHW has been classified as a strong event, and the MHW has been classified as moderate. Even though the Sentinel-2 of the 6th of September was captured only a few days after the start of the events, darkening of seagrass can already be observed in the true-color composition (Figure 9 C). Before the event, all GCPs appeared green on the images, with similar reflectance spectra, regardless of the class attributed to the GCPs on the field (Figure 9 A and D left). Their reflectance spectra showed a peak at 560 nm (in the green part of the spectra), low values at 665 nm, corresponding to the strong absorption by chlorophyll-a and a high infrared plateau (> 705 nm). However, GCPs classified as dark seagrass during the fieldtrip, showed significant differences in reflectance spectral shape compared to GCPs classified as green seagrass on the 10th of September, with corresponding differences in the true color composition (Figure 9 C and D right). In fact these dark seagrass GCPs were affected by extreme events, which altered their color and impacted the satellite reflectance (Figure 9 C). The reflectance spectra over dark seagrass were characterized by the loss of the reflectance peak at 560 nm and a decrease in the infrared plateau, which was observed as a steady increasing slope up to 940 nm.

**Figure 9**

*Spectral Comparison of Sentinel-2 Reflectance (D) over Ground Control Points (GCPs) Before (A) and During (B) Heatwave Events (C); A: RGB color composition of the Sentinel-2 image on the 1st of September 2021. The points correspond to GCPs collected on September 10, 2021 Figure 4; B: Detection of heatwave events based on both Air Temperature and Sea Surface Temperature (SST). The solid line represents the daily average temperature, while the dashed line indicates the 90th percentile of the climatology. Colored polygons identify heatwaves (marine in blue and atmospheric in red). The two vertical dashed lines represent the acquisition dates of the two Sentinel-2 images used in this analysis ; C : RGB color composition of the Sentinel-2 image on the 6th of September 2021. The points correspond to GCPs collected on September 10, 2021 Figure 4 ; D : Average spectral signatures of GCPs where dark and green seagrasses (blue and gold lines, respectively) were identified during the field survey. The left plot shows the reflectance of these GCPs before the heatwave impact, while the right plot shows the spectral signature during the heatwaves. The ribbons around the lines represent the standard deviation.*



### 3.2.2 Radiometric indices comparison


Using Sentinel-2 data, the spectral indices (NDVI and GLI) and SPC estimated for pixels unaffected by the heatwave showed no significant differences before or during the event (Figure 10, left column) In

contrast, seagrass darkened by the heatwave exhibited a significant decrease in both index values and SPC (Figure 10, right column). In this case, the NDVI dropped by approximately 34%, from a median of 0.61 to 0.40, while also becoming more homogeneous (the standard deviation decreased from 0.09 to 0.06) (Figure 10 B). Although the heatwave did not affect the observed seagrass density in the field, the drop in NDVI led to an underestimation of satellite-based SPC, from 83 to 48% (Figure 10 D). The GLI, the index that reflects the greenness of leaves, was the most affected by the event, with an important reduction from a median of 0.23 to 0.03 during the event (Figure 10 F). Interestingly, when comparing the initial seagrass density of both targets (i.e., seagrass density before the event) pixels affected by the heatwave (median SPC = 85%; [Figure 10 D green) were denser than the those remained unchanged (median SPC = 65%; [Figure 10 C green).



**Figure 10**

Comparison of Sentinel-2 based radiometric indices at two different dates, before (2021-09-01, green) and after (2021-09-06, red) the heatwaves events of Quiberons, for the two category of GCPs seen in the field : Green seagrasses (i.e seagrass not affected by the event, Left Column) and Darkened Seagrasses (i.e seagrass affected by the event, Right column). The Normalised Difference Vegetation Index (NDVI) has been computed using the Equation 1, the Seagrass Percent Cover (SPC) with the Equation 2 and the Green Leaf Index (GLI) using Equation 3,




### 3.2.3 Mapping of the seagrass darkening

Using the Equation 4, we can detect seagrass patches that have darkened. Following the heatwave events, a total of 26.9 hectares of seagrass darkened, with the largest affected patch covering nearly 8 hectares. Overall, 25 % of the total seagrass meadow area showed signs of darkening between the 1st and 6th of September 2021. Comparing the spatial distribution of darkened patches with the site's bathymetry reveals that 94.6 % of darkened areas are located at above 3.8 meters of bathymetry (Figure 11, A and B).

**Figure 11**

*Map of the Darkening of seagrasses during the heatwave events of September 2021 in Quiberon. A: 3.9 m bathymetry isobar (golden line); B: Result of the detection of darkened seagrass patches (Blue polygons)*




Additionally, analysis of bathymetry data to determine seagrass emersion time shows a clear relationship between the time seagrass is exposed to air and its darkening. During the heatwave, no significant darkening occurred with daily exposure below 13.5 hours. However, when exposure reached 13.5 hours or

306 more per day, seagrasses began to darken, peaking at around 14.5 hours of daily exposure (Figure 12).

**Figure 12**

*Seagrass Darkening Index as a function of the daily emergence time of seagrasses. The line represents a Generalized Additive Model (GAM) prediction, and the shaded area indicates the standard error. Shaded points represent raw data, with each point corresponding to a single pixel of the meadow.*




4 Discussion

4.1 Frenquency Intensity of Heatwaves and implication for seagrasses

Figure 13

Frequency and intensity of atmospheric heatwaves (red) and cold spells (blue) over time at the Lorient-Lann Bihoué weather station. The left column displays the annual frequency of heatwaves and cold spells as the total number of events per year. The right column illustrates event intensity, represented by the cumulative temperature anomaly throughout the year, measured in degrees Celsius.






## 4.2 Resilience of seagrasses to extreme events

**Figure 14**

Comparison of the seagrass percent cover before (left), during (center) and after (right) the heatwaves for seagrasses that have not been impacted by heatwaves (i.e. remains green) and seagrasses that have been impacted by the events (i.e. seagrasses that darkened).



## 5 Bibliography

- Akbar, M., Arisanto, P., Sukirno, B., Merdeka, P., Priadhi, M., Zallesa, S., 2020. Mangrove vegetation health index analysis by implementing NDVI (normalized difference vegetation index) classification method on sentinel-2 image data case study: Segara anakan, kabupaten cilacap, in: IOP Conference Series: Earth and Environmental Science. IOP Publishing, p. 012069.
- Cârlan, I., Mihai, B.-A., Nistor, C., Große-Stoltenberg, A., 2020. Identifying urban vegetation stress factors based on open access remote sensing imagery and field observations. *Ecological Informatics* 55, 101032.
- CMEMS, 2024. European north west shelf/iberia biscay irish seas – high resolution ODYSSEA sea surface temperature multi-sensor L3 observations reprocessed, e.u. Copernicus marine service information (CMEMS). Marine data store (MDS). (Accessed on 17-10-2024). <https://doi.org/10.48670/moi-00311>
- Deguet, A., Barrote, I., Silva, J., 2022. Physiological and morphological effects of a marine heatwave on the seagrass *Cymodocea nodosa*. *Scientific Reports* 12, 7950.
- European Space Agency, 2024b. Sen2Cor: Sentinel-2 atmospheric correction processor.
- European Space Agency, 2024a. Copernicus open access hub.
- Gardner, R.C., Finlayson, C., 2018. Global wetland outlook: State of the world's wetlands and their services to people, in: Ramsar Convention Secretariat. pp. 2020–5.
- Guerrero-Meseguer, L., Marín, A., Sanz-Lázaro, C., 2020. Heat wave intensity can vary the cumulative effects of multiple environmental stressors on *Posidonia oceanica* seedlings. *Marine Environmental Research* 159, 105001.
- Hobday, A.J., Alexander, L.V., Perkins, S.E., Smale, D.A., Straub, S.C., Oliver, E.C., Benthuyssen, J.A., Burrows, M.T., Donat, M.G., Feng, M., others, 2016. A hierarchical approach to defining marine heatwaves. *Progress in oceanography* 141, 227–238.

- Hobday, A.J., Oliver, E.C., Gupta, A.S., Benthuisen, J.A., Burrows, M.T., Donat, M.G., Holbrook, N.J., Moore, P.J., Thomsen, M.S., Wernberg, T., others, 2018. Categorizing and naming marine heatwaves. *Oceanography* 31, 162–173.
- Huete, A.R., 2012. Vegetation indices, remote sensing and forest monitoring. *Geography Compass* 6, 513–532.
- IOC, 2024. [Intergovernmental oceanographic commission ; sea level monitoring station - le conquet, france \(LECY\)](#).
- Jankowska, E., Michel, L.N., Lepoint, G., Włodarska-Kowalczyk, M., 2019. Stabilizing effects of seagrass meadows on coastal water benthic food webs. *Journal of Experimental Marine Biology and Ecology* 510, 54–63.
- Kloos, S., Yuan, Y., Castelli, M., Menzel, A., 2021. Agricultural drought detection with MODIS based vegetation health indices in southeast germany. *Remote Sensing* 13, 3907.
- Louhaichi, M., Borman, M.M., Johnson, D.E., 2001. Spatially located platform and aerial photography for documentation of grazing impacts on wheat. *Geocarto International* 16, 65–70.
- Papathanasopoulou, E., Simis, S., Alikas, K., Ansper, A., Anttila, J., Barillé, A., Barillé, L., Brando, V., Bresciani, M., Bučas, M., others, 2019. Satellite-assisted monitoring of water quality to support the implementation of the water framework directive. EOMORES white paper.
- Perkins, S.E., Alexander, L.V., 2013. On the measurement of heat waves. *Journal of climate* 26, 4500–4517.
- Ramesh, C., Mohanraju, R., 2020. Seagrass ecosystems of andaman and nicobar islands: Status and future perspective. *Environmental & Earth Sciences Research Journal* 7.
- Rouse, J.W., Haas, R.H., Schell, J.A., Deering, D.W., others, 1974. Monitoring vegetation systems in the great plains with ERTS. *NASA Spec. Publ* 351, 309.
- Sawall, Y., Ito, M., Pansch, C., 2021. Chronically elevated sea surface temperatures revealed high susceptibility of the eelgrass *zostera marina* to winter and spring warming. *Limnology and Oceanography* 66, 4112–4124.
- Schlegel, R.W., Oliver, E.C., Wernberg, T., Smit, A.J., 2017. Nearshore and offshore co-occurrence of marine heatwaves and cold-spells. *Progress in Oceanography* 151, 189–205.
- Schlegel, R.W., Smit, A.J., 2018. heatwaveR: A central algorithm for the detection of heatwaves and cold-spells. *Journal of Open Source Software* 3, 821. <https://doi.org/10.21105/joss.00821>
- Scott, A.L., York, P.H., Duncan, C., Macreadie, P.I., Connolly, R.M., Ellis, M.T., Jarvis, J.C., Jinks, K.I., Marsh, H., Rasheed, M.A., 2018. The role of herbivory in structuring tropical seagrass ecosystem service delivery. *Frontiers in Plant Science* 9, 127.
- Sevgi, K., Leblebici, S., 2022. Bitkilerde ağır metal stresine verilen fizyolojik ve moleküler yanıtlar. *Journal of Anatolian Environmental and Animal Sciences* 7, 528–536.
- Shom, 2022. [Service hydrographique et océanographique de la marine ; références altimétriques maritimes: Ports de france métropolitaine et d'outre-mer, cotes du zéro hydrographique et niveaux caractéristiques de la marée](#). Shom, Brest, France.
- SHOM, 2021. [Service hydrographique et océanographique de la marine ; bathymétrie Litto3D® bretagne 2018-2021](#).
- Simpson, T.S., Wernberg, T., McDonald, J.I., 2016. Distribution and localised effects of the invasive ascidian *didemnum perlucidum* (monniot 1983) in an urban estuary. *PLoS One* 11, e0154201.
- Sousa, A.I., Silva, J.F. da, Azevedo, A., Lillebø, A.I., 2019. Blue carbon stock in *zostera noltei* meadows at ria de aveiro coastal lagoon (portugal) over a decade. *Scientific reports* 9, 14387.
- Thomsen, E., Herbeck, L.S., Viana, I.G., Jennerjahn, T.C., 2023. Meadow trophic status regulates the nitrogen filter function of tropical seagrasses in seasonally eutrophic coastal waters. *Limnology and Oceanography* 68, 1906–1919.
- Unsworth, R.K., Cullen-Unsworth, L.C., Jones, B.L., Lilley, R.J., 2022. The planetary role of seagrass conservation. *Science* 377, 609–613.

- Waycott, M., Duarte, C.M., Carruthers, T.J., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A.,  
Fourqurean, J.W., Heck Jr, K.L., Hughes, A.R., others, 2009. Accelerating loss of seagrasses across the  
globe threatens coastal ecosystems. *Proceedings of the national academy of sciences* 106, 12377–12381.
- Winters, G., Nelle, P., Fricke, B., Rauch, G., Reusch, T.B., 2011. Effects of a simulated heat wave on  
photophysiology and gene expression of high-and low-latitude populations of *zostera marina*. *Marine  
Ecology Progress Series* 435, 83–95.
- Zoffoli, M.L., Gernez, P., Godet, L., Peters, S., Oiry, S., Barillé, L., 2021. Decadal increase in the ecological  
status of a north-atlantic intertidal seagrass meadow observed with multi-mission satellite time-series.  
*Ecological Indicators* 130, 108033.
- Zoffoli, M.L., Gernez, P., Oiry, S., Godet, L., Dalloyau, S., Davies, B.F.R., Barillé, L., 2023. Remote sensing  
in seagrass ecology: Coupled dynamics between migratory herbivorous birds and intertidal meadows  
observed by satellite during four decades. *Remote Sensing in Ecology and Conservation* 9, 420–433.
- Zoffoli, M.L., Gernez, P., Rosa, P., Le Bris, A., Brando, V.E., Barillé, A.-L., Harin, N., Peters, S., Poser,  
K., Spaias, L., others, 2020. Sentinel-2 remote sensing of *zostera noltei*-dominated intertidal seagrass  
meadows. *Remote Sensing of Environment* 251, 112020.