

Seagrass mapping in two mudflats in the Auray River

About a rapid evolution of seagrasses

Simon Oiry^{a,*}

^a*Nantes Université, UR 2160, F-44000 Nantes, France, Institut des Substances et Organismes de la Mer, ISOMer, 2 chemin de la houssinière, Nantes, 44300*

Abstract

Maps of seagrass in two sites in the Auray River. These two sites were studied by Maxime Daviray during his PhD. Seagrass appeared very quickly during his PhD. This work aims to describe this rapid evolution of seagrasses.

Keywords: Remote Sensing, Sentinel-2, Seagrass

The data and scripts used for this work can be found [here](#).

1. Materiel & Methods

1.1. Seagrass mapping using Sentinel-2

To map the seagrass extent over time, the Sentinel-2 constellation has been used. Level-2 images, which are already orthorectified and atmospherically corrected using Sen2Cor, have been downloaded using the Copernicus Platform [1]. One low tide, cloud-free image per year, nearest to the period of maximum seagrass biomass at this latitude, has been used. The ICECREAM model (Davies et al., Accepted), a neural network classifier designed to identify and discriminate intertidal vegetation in Europe, has been applied to each Sentinel-2 image. Pixels of the Magnoliopsida class (seagrasses) have been isolated, and the Normalized Difference Vegetation Index (NDVI, [2]), a commonly used remote sensing biomass proxy for vegetation, has been employed.

References

- [1] Copernicus, [Copernicus open access hub](#), accessed: 2024-07-16 (2024).
URL <https://browser.dataspace.copernicus.eu/>

*Corresponding author
Email address: oirysimon@gmail.com (Simon Oiry)

- [2] J. W. Rouse, R. H. Haas, J. A. Schell, D. W. Deering, et al., Monitoring vegetation systems in the great plains with erts, NASA Spec. Publ 351 (1) (1974) 309.