

# Seagrass mapping in two mudflats in the Auray River

## About a rapid evolution of seagrasses

Simon Oiry<sup>a,\*</sup>

<sup>a</sup>*Nantes Université, UR 2160, F-44000 Nantes, France, Institut des Substances et Organismes de la Mer, ISOMer, 2 chemin de la houssinière, Nantes, 44300*

---

### Abstract

Maps of seagrass in two sites in the Auray River. These two sites were studied by Maxime Daviray during his PhD. Seagrass appeared very quickly during his PhD. This work aims to describe this rapid evolution of seagrasses.

*Keywords:* Remote Sensing, Sentinel-2, Seagrass

---

The data and scripts used for this work can be found [here](#).

## 1. Materiel & Methods

### 1.1. Seagrass mapping using Sentinel-2

To map the seagrass extent over time, the Sentinel-2 constellation has been used. Level-2 images, which are already orthorectified and atmospherically corrected using Sen2Cor, have been downloaded using the Copernicus Platform [1]. One low tide, cloud-free image per year, nearest to the period of maximum seagrass biomass at this latitude, has been used. The ICECREAM model (Davies et al., Accepted), a neural network classifier designed to identify and discriminate intertidal vegetation in Europe, has been applied to each Sentinel-2 image. Pixels of the Magnoliopsida class (seagrasses) have been isolated, and the Normalized Difference Vegetation Index (NDVI, [2]), a commonly used remote sensing biomass proxy for vegetation, has been employed. The equation of [3] have been used to transform NDVI values into Seagrass Percent Cover Equation 1. SPC values below 20%, corresponding to low biomass pixel, with a high risk of confusion with other vegetation classes, have been remove from the rest of the analysis.

$$SPC = 172.06 * NDVI - 22.18 \quad (1)$$

---

\*Corresponding author

Email address: [oirysimon@gmail.com](mailto:oirysimon@gmail.com) (Simon Oiry)

## References

- [1] Copernicus, [Copernicus open access hub](https://browser.dataspace.copernicus.eu/), accessed: 2024-07-16 (2024).  
URL <https://browser.dataspace.copernicus.eu/>
- [2] J. W. Rouse, R. H. Haas, J. A. Schell, D. W. Deering, et al., Monitoring vegetation systems in the great plains with erts, NASA Spec. Publ 351 (1) (1974) 309.
- [3] M. L. Zoffoli, P. Gernez, L. Godet, S. Peters, S. Oiry, L. Barillé, Decadal increase in the ecological status of a north-atlantic intertidal seagrass meadow observed with multi-mission satellite time-series, Ecological Indicators 130 (2021) 108033.