Pingala Series

Sumanth N R

1

2

CONTENTS

- 1 JEE 2019
- 2 Pingala Series 1
- 3 Power of the Z transform

Abstract—This manual provides a simple introduction to Transforms

1 JEE 2019

Let

$$a_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad n \ge 1 \tag{1.1}$$

$$b_n = a_{n-1} + a_{n+1}, \quad n \ge 2, \quad b_1 = 1$$
 (1.2)

Verify the following using a python code.

1.1

$$\sum_{k=1}^{n} a_k = a_{n+2} - 1, \quad n \ge 1$$
 (1.3)

1.2

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{10}{89} \tag{1.4}$$

1.3

$$b_n = \alpha^n + \beta^n, \quad n \ge 1 \tag{1.5}$$

1.4

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{8}{89} \tag{1.6}$$

Solution: The following code verifies all the equations

wget https://raw.githubusercontent.com/Sigma1084/EE3900/master/pingala/code/Ex1 verify.py

2 Pingala Series

2.1 The *one sided Z*-transform of x(n) is defined as

$$X^{+}(z) = \sum_{n=0}^{\infty} x(n)z^{-n}, \quad z \in \mathbb{C}$$
 (2.1)

2.2 The *Pingala* series is generated using the difference equation

$$x(n+2) = x(n+1) + x(n)$$

$$x(0) = x(1) = 1, n > 0$$
 (2.2)

Generate a stem plot for x(n).

Solution: The following code generates the plot

wget https://raw.githubusercontent.com/ Sigma1084/EE3900/master/pingala/code/ Ex2 pingala.py




Fig. 2.2: Plot of x(n)

2.3 Find $X^{+}(z)$.

Solution: Taking the *one-sided Z*-transform on both sides of (2.2)

$$Z^{+}[x(n+2)] = Z^{+}[x(n+1)] + Z^{+}[x(n)]$$

$$\Rightarrow z^{2}X^{+}(z) - z^{2} - z = zX^{+}(z) - z + X^{+}(z)$$

$$\Rightarrow (z^{2} - z - 1)X^{+}(z) = z^{2}$$

$$\Rightarrow X^{+}(z) = \frac{1}{1 - z^{-1} - z^{-2}}$$

$$X^{+}(z) = \frac{1}{(1 - \alpha z^{-1})(1 - \beta z^{-1})}, \quad |z| > |\alpha|$$
(2.3)

Here α and β are the roots of the characteristic equation of (2.2) and $|z| > |\alpha|$ is the region of convergence of $X^+(z)$.

(w.l.o.g, $|\alpha| > |\beta|$ is assumed)

2.4 Find x(n).

Solution: Expanding $X^+(z)$ in (2.3) using partial fractions, we get

$$X^{+}(z) = \frac{1}{(\alpha - \beta) z^{-1}} \left[\frac{1}{1 - \alpha z^{-1}} - \frac{1}{1 - \beta z^{-1}} \right]$$

$$= \frac{1}{(\alpha - \beta)} \sum_{n=0}^{\infty} (\alpha^{n} - \beta^{n}) z^{-n}$$

$$= \sum_{n=1}^{\infty} \frac{\alpha^{n} - \beta^{n}}{\alpha - \beta} z^{-n+1}$$

$$= \sum_{n=0}^{\infty} \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta} z^{-n}$$

$$x(n) = \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta} u(n) = a_{n+1} u(n)$$
 (2.4)

2.5 Sketch

$$y(n) = x(n-1) + x(n+1), \quad n \ge 0$$
 (2.5)

Solution: The following code generates the plot

wget https://raw.githubusercontent.com/ Sigma1084/EE3900/master/pingala/code/ Ex2_pingala.py




Fig. 2.5: Plot of y(n)

3 Power of the Z transform

3.1 Show that

$$\sum_{k=1}^{n} a_k = \sum_{k=0}^{n-1} x(n) = x(n) * u(n-1)$$
 (3.1)

3.2 Show that

$$a_{n+2} - 1, \quad n \ge 1$$
 (3.2)

can be expressed as

$$[x(n+1)-1]u(n)$$
 (3.3)

3.3 Show that

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{x(k)}{10^k} = \frac{1}{10} X^+ (10)$$
 (3.4)

3.4 Show that

$$\alpha^n + \beta^n, \quad n \ge 1 \tag{3.5}$$

can be expressed as

$$w(n) = (\alpha^{n+1} + \beta^{n+1})u(n)$$
 (3.6)

and find W(z).

3.5 Show that

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{y(k)}{10^k} = \frac{1}{10} Y^+ (10)$$
 (3.7)

3.6 Solve the JEE 2019 problem.

- 2.6 Find $Y^{+}(z)$.
- 2.7 Find y(n).