

# Derivations of aerodynamic equations

siim.alas

April 2021

## 1 Introduction

Bla.

## 2 Potential Flow Theory

This section contains derivations of equations relating to [potential flow theory](#), which describes [flow velocity](#)  $\vec{v}(\vec{r})$  at some point  $\vec{r}$  in space as the [gradient](#) of the [velocity potential](#)  $\phi(\vec{r})$  at that point. More succinctly, this is written as

$$\vec{v}(\vec{r}) = \nabla\phi(\vec{r}) \quad (1)$$

### 2.1 3D Source Panel Method

This subsection contains derivations of equations used in the 3D source panel method, which estimates the aerodynamic properties of a body in an [inviscid](#) flow.

The body is approximated as an array of [rectangular](#) panels, each of which is assumed to have constant [velocity potential](#) source (or sink) strength across its surface.

#### 2.1.1 The Effect of a Rectangular Source Panel on the Velocity Potential

The aforementioned rectangular panel is defined in 3D [Cartesian coordinates](#) by a [position vector](#)  $\vec{r}_0$  (pointing to the centre of the rectangle), two [orthonormal tangent vectors](#)  $\hat{u}$  ([parallel](#) to one side of the rectangle) and  $\hat{v}$  (parallel to the other side of the rectangle) as well as the dimensions  $a$  (the length of the side parallel to  $\hat{u}$ ) and  $b$  (the length of the side parallel to  $\hat{v}$ ).

The effect of a point source ( $Q > 0$ ) or sink ( $Q < 0$ ) at position  $\vec{r}_0$  on the velocity potential  $\phi(\vec{r})$  at position  $r$  is assumed to be

$$\phi(\vec{r}) = -\frac{Q}{4\pi|\vec{r} - \vec{r}_0|} \quad (2)$$

Modelling the source rectangle as consisting of an infinite series of such point sources, all of constant strength  $Q$ , yields the following [surface integral](#):

$$\phi(\vec{r}) = -\frac{Q}{4\pi} \iint_S \frac{1}{|\vec{r} - \vec{r}'|} dS = -\frac{Q}{4\pi} \iint_T \frac{1}{|\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)|} du dv \quad (3)$$

where  $\vec{r}' = \vec{r}_0 + \hat{u}u + \hat{v}v$ ,  $S = \{\vec{r}_0 + \hat{u}u + \hat{v}v | u, v \in T\}$ , and  $T = [-a/2; a/2] \times [-b/2; b/2]$ . Note that in the above surface integral  $|\frac{\partial \vec{r}'}{\partial u} \times \frac{\partial \vec{r}'}{\partial v}| = |\hat{u} \times \hat{v}| = 1$ , since  $\hat{u}$  and  $\hat{v}$  are orthonormal.

The denominator in the integrand can be simplified to

$$\begin{aligned} & \sqrt{|\hat{u}u + \hat{v}v|^2 - 2(\hat{u}u + \hat{v}v) \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - \vec{r}_0|^2} = \\ & = \sqrt{(\hat{u} \cdot \hat{u})u^2 + 2uv(\hat{u} \cdot \hat{v}) + (\hat{v} \cdot \hat{v})v^2 - 2(\hat{u}u + \hat{v}v) \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - \vec{r}_0|^2} = \\ & = \sqrt{u^2 + v^2 - 2(\hat{u}u + \hat{v}v) \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - \vec{r}_0|^2} = \\ & = \sqrt{(u - \hat{u} \cdot (\vec{r} - \vec{r}_0))^2 + (v - \hat{v} \cdot (\vec{r} - \vec{r}_0))^2 + k} \end{aligned}$$

where we defined  $k = |\vec{r} - \vec{r}_0|^2 - (\hat{u} \cdot (\vec{r} - \vec{r}_0))^2 - (\hat{v} \cdot (\vec{r} - \vec{r}_0))^2$  and used the fact that, by definition,  $\hat{u} \cdot \hat{u} = \hat{v} \cdot \hat{v} = 1$  and  $\hat{u} \cdot \hat{v} = 0$ .

Changing variables in integral 3 via  $u - \hat{u} \cdot (\vec{r} - \vec{r}_0) = iu'$  and  $v - \hat{v} \cdot (\vec{r} - \vec{r}_0) = iv'$ , leading to the [Jacobian determinant](#)

$$\det(J) = \det \left( \begin{bmatrix} \frac{\partial u}{\partial u'} & \frac{\partial u}{\partial v'} \\ \frac{\partial v}{\partial u'} & \frac{\partial v}{\partial v'} \end{bmatrix} \right) = \det \left( \begin{bmatrix} i & 0 \\ 0 & i \end{bmatrix} \right) = i^2 - 0 = -1$$

we write the integral as

$$\frac{Q}{4\pi} \iint_{T'} \frac{1}{\sqrt{k - u'^2 - v'^2}} du' dv'$$

where  $T' = [u'_1; u'_2] \times [v'_1; v'_2] = [-i(-a/2 - \hat{u} \cdot (\vec{r} - \vec{r}_0)); -i(a/2 - \hat{u} \cdot (\vec{r} - \vec{r}_0))] \times [-i(-b/2 - \hat{v} \cdot (\vec{r} - \vec{r}_0)); -i(b/2 - \hat{v} \cdot (\vec{r} - \vec{r}_0))]$ . Using the identity  $\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin(\frac{x}{a}) + C$ , we integrate in  $u'$  to get

$$\frac{Q}{4\pi} \int_{v'_1}^{v'_2} \left( \arcsin \left( \frac{u'_2}{\sqrt{k - v'^2}} \right) - \arcsin \left( \frac{u'_1}{\sqrt{k - v'^2}} \right) \right) dv'$$

The exponential definition  $\arcsin(x) = -i \ln(ix + \sqrt{1 - x^2})$  along with the identity  $\ln(a/b) = \ln(a) - \ln(b)$  allows us to rewrite the above integral as

$$-i \frac{Q}{4\pi} \int_{v'_1}^{v'_2} \left( \ln \left( iu'_2 + \sqrt{k - (u'_2)^2 - v'^2} \right) - \ln \left( iu'_1 + \sqrt{k - (u'_1)^2 - v'^2} \right) \right) dv'$$

To solve an integral in the form of

$$\int \ln \left( a + \sqrt{b - x^2} \right) dx$$

we first use integration by parts to get

$$x \ln \left( a + \sqrt{b - x^2} \right) - \int x \frac{d}{dx} \left( \ln \left( a + \sqrt{b - x^2} \right) \right) dx$$

Following just the above integral, we write it as

$$\begin{aligned} \int x \frac{d}{dx} \left( \ln \left( a + \sqrt{b - x^2} \right) \right) dx &= - \int \frac{x^2}{(a + \sqrt{b - x^2})\sqrt{b - x^2}} dx = \\ &= - \int \frac{x^2(a - \sqrt{b - x^2})}{(a^2 - b + x^2)\sqrt{b - x^2}} dx = \\ &= - \int \left( \frac{a^2 - b}{a^2 - b + x^2} - a \frac{a^2 - b}{(a^2 - b + x^2)\sqrt{b - x^2}} + \frac{a}{\sqrt{b - x^2}} - 1 \right) dx \end{aligned}$$

where we used a technique similar to [partial fraction decomposition](#) to transform the integral into a form where we can use the identities

$$\begin{aligned} \int \frac{dx}{a^2 + x^2} &= \frac{1}{a} \arctan \left( \frac{x}{a} \right) + C \\ \int \frac{dx}{(a^2 - b + x^2)\sqrt{b - x^2}} &= \frac{1}{a\sqrt{a^2 - b}} \arctan \left( \frac{ax}{\sqrt{a^2 - b}\sqrt{b - x^2}} \right) + C \\ \int \frac{dx}{\sqrt{a - x^2}} &= \arctan \left( \frac{x}{\sqrt{a - x^2}} \right) + C \end{aligned}$$

to solve the "decomposed" integral, resulting in

$$\begin{aligned} \int \ln \left( a + \sqrt{b - x^2} \right) dx &= x \ln \left( a + \sqrt{b - x^2} \right) + \sqrt{a^2 - b} \arctan \left( \frac{x}{\sqrt{a^2 - b}} \right) - \\ &\quad - \sqrt{a^2 - b} \arctan \left( \frac{ax}{\sqrt{a^2 - b}\sqrt{b - x^2}} \right) + a \arctan \left( \frac{x}{\sqrt{b - x^2}} \right) - x + C \end{aligned}$$

Tying this back into the original problem, we substitute in  $a_n = iu'_n$ ,  $b_n = k - (u'_n)^2$ , and  $x = v'$  and simplify to get

$$\begin{aligned}
& -i \frac{Q}{4\pi} \left( \left( v' \ln \left( iu' + \sqrt{k - u'^2 - v'^2} \right) + \sqrt{-k} \arctan \left( \frac{v'}{\sqrt{-k}} \right) - \right. \right. \\
& \quad \left. \left. - \sqrt{-k} \arctan \left( \frac{i u' v'}{\sqrt{-k} \sqrt{k - u'^2 - v'^2}} \right) + i u' \arctan \left( \frac{v'}{\sqrt{k - u'^2 - v'^2}} \right) - \right. \right. \\
& \quad \left. \left. - v' + C \right) \right|_{u'=u'_1}^{u'=u'_2} \bigg|_{v'=v'_1}^{v'=v'_2} = \\
& = -\frac{Q}{4\pi} \left( \left( i v' \ln \left( iu' + \sqrt{k - u'^2 - v'^2} \right) + \sqrt{k} \arctan \left( \frac{u' v'}{\sqrt{k} \sqrt{k - u'^2 - v'^2}} \right) + \right. \right. \\
& \quad \left. \left. - u' \arctan \left( \frac{v'}{\sqrt{k - u'^2 - v'^2}} \right) \right) \right|_{u'=u'_1}^{u'=u'_2} \bigg|_{v'=v'_1}^{v'=v'_2}
\end{aligned}$$

To emphasize the [symmetry](#) in  $u'$  and  $v'$ , we use the exponential definition  $\arctan(x) = \frac{i}{2} \ln \left( \frac{i+x}{i-x} \right)$  to rewrite the above expression as

$$\begin{aligned}
& -\frac{Q}{4\pi} \left( \left( i v' \ln \left( iu' + \sqrt{k - u'^2 - v'^2} \right) + \sqrt{k} \arctan \left( \frac{u' v'}{\sqrt{k} \sqrt{k - u'^2 - v'^2}} \right) + \right. \right. \\
& \quad \left. \left. + i u' \ln \left( i v' + \sqrt{k - u'^2 - v'^2} \right) \right) \right|_{u'=u'_1}^{u'=u'_2} \bigg|_{v'=v'_1}^{v'=v'_2}
\end{aligned}$$

where the  $\frac{i}{2} \ln(k - u'^2)$  term has been ignored since being invariant in  $v'$  means it will get cancelled in the outer  $v'$ -summation. Transitioning back to the original  $u$  and  $v$  variables, we get

$$\begin{aligned}
\phi(\vec{r}) = & -\frac{Q}{4\pi} \left( \left( (v - \hat{v} \cdot (\vec{r} - \vec{r}_0)) \ln (u - \hat{u} \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - (\vec{r}_0 - \hat{u}u + \hat{v}v)|) - \right. \right. \\
& \quad \left. \left. - \sqrt{k} \arctan \left( \frac{(u - \hat{u} \cdot (\vec{r} - \vec{r}_0))(v - \hat{v} \cdot (\vec{r} - \vec{r}_0))}{\sqrt{k} |\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)|} \right) + \right. \right. \\
& \quad \left. \left. + (u - \hat{u} \cdot (\vec{r} - \vec{r}_0)) \ln (v - \hat{v} \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)|) \right) \right|_{u=-\frac{a}{2}}^{u=\frac{a}{2}} \bigg|_{v=-\frac{b}{2}}^{v=\frac{b}{2}} \quad (4)
\end{aligned}$$

where, restating for convenience,  $k = |\vec{r} - \vec{r}_0|^2 - (\hat{u} \cdot (\vec{r} - \vec{r}_0))^2 - (\hat{v} \cdot (\vec{r} - \vec{r}_0))^2$ .

### 2.1.2 The Effect of a Rectangular Source Panel on Velocity

Given equations 1 and 3, finding the effect of a rectangular panel as defined above on the flow velocity simply entails taking the gradient of equation 3.

In the interest of brevity, we only compute the [partial derivative](#) with respect to  $r_i$  (the  $i$ -th component of  $\vec{r}$ ). Because the bounds of integration in [3](#) are constant in  $r_i$ , we can differentiate the integrand before integrating, resulting in

$$\begin{aligned}\frac{\partial}{\partial r_i}\phi(\vec{r}) &= -\frac{Q}{4\pi} \iint_T \frac{\partial}{\partial r_i} \left( \frac{1}{|\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)|} \right) dudv = \\ &= -\frac{Q}{4\pi} \iint_T \frac{(r_{0i} + u_i u + v_i v) - r_i}{|\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)|^3} dudv\end{aligned}$$

We can represent the [denominator](#) of the integrand as

$$\left( \sqrt{|\hat{u}u + \hat{v}v|^2 - 2(\hat{u}u + \hat{v}v) \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - \vec{r}_0|^2} \right)^3 = \left( \sqrt{u'^2 + v'^2 + k} \right)^3$$

where  $u' = u - \hat{u} \cdot (\vec{r} - \vec{r}_0)$ ,  $v' = v - \hat{v} \cdot (\vec{r} - \vec{r}_0)$  and  $k = |\vec{r} - \vec{r}_0|^2 - (\hat{u} \cdot (\vec{r} - \vec{r}_0))^2 - (\hat{v} \cdot (\vec{r} - \vec{r}_0))^2$ . Because the transition to the  $u'$  and  $v'$  coordinates is a simple [translation](#), its [Jacobian](#) is always 1, allowing us to rewrite the integral as

$$-\frac{Q}{4\pi} \iint_{T'} \frac{u_i u' + v_i v' - c_i}{(\sqrt{u'^2 + v'^2 + k})^3} du' dv'$$

where  $T' = [u'_1; u'_2] \times [v'_1; v'_2] = [-a/2 - \hat{u} \cdot (\vec{r} - \vec{r}_0); a/2 - \hat{u} \cdot (\vec{r} - \vec{r}_0)] \times [-b/2 - \hat{v} \cdot (\vec{r} - \vec{r}_0); b/2 - \hat{v} \cdot (\vec{r} - \vec{r}_0)]$  and  $c_i = r_i - r_{0i} - u_i(\hat{u} \cdot (\vec{r} - \vec{r}_0)) - v_i(\hat{v} \cdot (\vec{r} - \vec{r}_0))$ . Using the identities  $\int \frac{x dx}{(\sqrt{x^2 + a})^3} = -\frac{1}{\sqrt{x^2 + a}} + C$  and  $\int \frac{dx}{(\sqrt{x^2 + a})^3} = \frac{x}{a\sqrt{x^2 + a}} + C$  to integrate in  $u'$ , we get

$$\begin{aligned}& -\frac{Q}{4\pi} \int_{v'=v'_1}^{v'=v'_2} \left( -\frac{u_i}{\sqrt{u'^2 + v'^2 + k}} + \frac{v_i u' v' - c_i u'}{(v'^2 + k)\sqrt{u'^2 + v'^2 + k}} \right) \Big|_{u'=u'_1}^{u'=u'_2} dv' = \\ & -\frac{Q}{4\pi} \int_{v'=v'_1}^{v'=v'_2} \left( -u_i \frac{1}{\sqrt{u'^2 + v'^2 + k}} - v_i \frac{1}{1 - \left( \frac{\sqrt{u'^2 + v'^2 + k}}{u'} \right)^2} \cdot \frac{v'}{u' \sqrt{u'^2 + v'^2 + k}} - \right. \\ & \quad \left. - \frac{c_i}{\sqrt{k}} \frac{1}{1 + \left( \frac{u' v'}{\sqrt{k} \sqrt{u'^2 + v'^2 + k}} \right)^2} \cdot \frac{u'(u'^2 + k)}{\sqrt{k} (\sqrt{u'^2 + v'^2 + k})^3} \right) \Big|_{u'=u'_1}^{u'=u'_2} dv'\end{aligned}$$

We now use the identities  $\int \frac{dx}{\sqrt{x^2 + a^2}} = \ln \left( \frac{x + \sqrt{x^2 + a^2}}{a} \right) + C$ ,  $\frac{d}{dx} \text{artanh}(x) = \frac{1}{1-x^2}$ , and  $\frac{d}{dx} \arctan(x) = \frac{1}{1+x^2}$  to integrate in  $v'$ , leading to

$$\begin{aligned} \frac{Q}{4\pi} \left( \left( u_i \ln \left( \frac{v' + \sqrt{u'^2 + v'^2 + k}}{\sqrt{u'^2 + k}} \right) + v_i \operatorname{artanh} \left( \frac{\sqrt{u'^2 + v'^2 + k}}{u'} \right) + \right. \right. \\ \left. \left. + \frac{c_i}{\sqrt{k}} \arctan \left( \frac{u'v'}{\sqrt{k}\sqrt{u'^2 + v'^2 + k}} \right) \right) \right|_{u'=u'_1}^{u'=u'_2} \right) \bigg|_{v'=v'_1}^{v'=v'_2} \end{aligned}$$

Using the exponential definition  $\operatorname{artanh}(x) = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$  and noting that terms constant in either  $u'$  or  $v'$  will get cancelled, we rewrite this as

$$\begin{aligned} \frac{Q}{4\pi} \left( \left( u_i \ln \left( v' + \sqrt{u'^2 + v'^2 + k} \right) + v_i \ln \left( u' + \sqrt{u'^2 + v'^2 + k} \right) + \right. \right. \\ \left. \left. + \frac{c_i}{\sqrt{k}} \arctan \left( \frac{u'v'}{\sqrt{k}\sqrt{u'^2 + v'^2 + k}} \right) \right) \right|_{u'=u'_1}^{u'=u'_2} \right) \bigg|_{v'=v'_1}^{v'=v'_2} \end{aligned}$$

Transitioning back to the original  $u$  and  $v$  coordinates, we see trivially that

$$\begin{aligned} \vec{v}(\vec{r}) = \nabla \phi(\vec{r}) = \frac{Q}{4\pi} \left( \left( \hat{u} \ln \left( v - \hat{v} \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)| \right) + \right. \right. \\ \left. \left. + \hat{v} \ln \left( u - \hat{u} \cdot (\vec{r} - \vec{r}_0) + |\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)| \right) + \right. \right. \\ \left. \left. + \frac{\vec{c}}{\sqrt{k}} \arctan \left( \frac{(u - \hat{u} \cdot (\vec{r} - \vec{r}_0))(v - \hat{v} \cdot (\vec{r} - \vec{r}_0))}{\sqrt{k}|\vec{r} - (\vec{r}_0 + \hat{u}u + \hat{v}v)|} \right) \right) \right|_{u=u_1}^{u=u_2} \right) \bigg|_{v=v_1}^{v=v_2} \quad (5) \end{aligned}$$

where  $\vec{c} = \vec{r} - \vec{r}_0 - \hat{u}(\hat{u} \cdot (\vec{r} - \vec{r}_0)) - \hat{v}(\hat{v} \cdot (\vec{r} - \vec{r}_0))$  and, restating for convenience,  $k = |\vec{r} - \vec{r}_0|^2 - (\hat{u} \cdot (\vec{r} - \vec{r}_0))^2 - (\hat{v} \cdot (\vec{r} - \vec{r}_0))^2$ .