

RDD and diff-in-diff

Silje Synnøve Lyder Hermansen

08-12-2020

Regression discontinuity design (RDD)

Basic assumption

RDD assumes a running variable (x) with a cut point (c) beyond which treatment is assigned (D).

$$D_i = \begin{cases} 1 & \text{if } x_i \geq c \\ 0 & \text{if } x_i < c \end{cases} \quad (1)$$

Distinction

It has a flavor of logit or propensity scores, but there are some differences:

- ▶ **logit** : x (not y) is not latent and we know the cutpoint: Both are observed and included as a *predictors*.
- ▶ **matching** : we have no control/treatment group. However, we assume that units on either side of the treatment are increasingly similar as their x is similar.

⇒ *Supposes clear rules with little administrative discretion.*

Examples

Administrative data are perfect: You have some rule that kicks in at a specific threshold for otherwise almost identical observations.

- ▶ school test scores on school admission, restrictions on class size
- ▶ legal drinking age on alcohol related deaths
- ▶ election of candidates in close races

Two ways of understanding RDD

- ▶ **Individuals close to the threshold are interchangeable**

Two ways of understanding RDD

► **Individuals close to the threshold are interchangeable**

→ in a small window, you have a treatment and a control group.

Two ways of understanding RDD

- ▶ **Individuals close to the threshold are interchangeable**

→ in a small window, you have a treatment and a control group.

- ▶ **x is a bottleneck:** the relationship between D and Y is confounded by x , but all other confounders only influence Y through x .

Two ways of understanding RDD

- ▶ **Individuals close to the threshold are interchangeable**

→ in a small window, you have a treatment and a control group.

- ▶ **x is a bottleneck:** the relationship between D and Y is confounded by x , but all other confounders only influence Y through x .

→ conditioning on x is sufficient to isolate the causal effect.

Two ways of understanding RDD

- ▶ **Individuals close to the threshold are interchangeable**


→ in a small window, you have a treatment and a control group.

- ▶ **x is a bottleneck:** the relationship between D and Y is confounded by x , but all other confounders only influence Y through x .

→ conditioning on x is sufficient to isolate the causal effect.

Two ways of understanding RDD

X is a confounder
...so we only control for X




Two designs

We distinguish between two designs depending on how probable the treatment is:

- ▶ **sharp** RD: assignment is *deterministic*
- ▶ **fuzzy** RD: assignment is *probabilistic*

A visual representation



The basic model

The basic regression

We assume the relationship between x and y is linear and the treatment is deterministic

The basic regression

We assume the relationship between x and y is linear and the treatment is deterministic

$$y_i = \alpha + \rho \times D_i + \gamma \times x_1 + e_i \quad (2)$$

The basic regression

We assume the relationship between x and y is linear and the treatment is deterministic

$$y_i = \alpha + \rho \times D_i + \gamma \times x_1 + e_i \quad (2)$$

\Rightarrow *The treatment is reported by ρ*

The basic regression

We assume the relationship between x and y is linear and the treatment is deterministic

$$y_i = \alpha + \rho \times D_i + \gamma \times x_1 + e_i \quad (2)$$

\Rightarrow *The treatment is reported by ρ*

Do it in R

Check out alcohol related deaths (y) as a function of legal drinking age (D)

```
load("df_ch4.Rda")  
mod1 <- lm(alcohol ~ over21 + age,  
            df)
```

Is that all?

This is true on two conditions

Is that all?

This is true on two conditions

1. **the continuity assumption:** x must have a continuous effect on y

Is that all?

This is true on two conditions

1. **the continuity assumption:** x must have a continuous effect on y
2. **no omitted variable bias:** x must capture all influence on D .

The continuity assumption

The continuity assumption

Sometimes we may pick up a smooth non-linear change by dummy coding

The continuity assumption

Sometimes we may pick up a smooth non-linear change by dummy coding

... that's not a regression discontinuity.

Ensuring linear effect

We can obtain a linear effect in two ways:

Ensuring linear effect

We can obtain a linear effect in two ways:

- ▶ recode the x

Ensuring linear effect

We can obtain a linear effect in two ways:

- ▶ recode the $x \rightarrow$ compare with the recoding of y in GLMs.

Ensuring linear effect

We can obtain a linear effect in two ways:

- ▶ recode the $x \rightarrow$ compare with the recoding of y in GLMs.
- ▶ consider a sufficiently small window

Recode the x

We can create a curvilinear effect of x using polynomials (e.g.:)

Recode the x

We can create a curvilinear effect of x using polynomials (e.g.):

$$y_i = \alpha + \rho D_i + \gamma_1 x_i + \gamma_2 x_i^2 \quad (3)$$

Recode the x

We can create a curvilinear effect of x using polynomials (e.g.):

$$y_i = \alpha + \rho D_i + \gamma_1 x_i + \gamma_2 x_i^2 \quad (3)$$

```
df$age2 <- df$age^2
mod2 <- lm(alccohol ~ over21 + age + age2,
           df)
```


Recode the x: symmetric effect

We can create a curvilinear effect of x using polynomials (e.g.):

$$y_i = \alpha + \rho D_i + \gamma_1 x_i + \gamma_2 x_i^2 \quad (4)$$

```
df$age2 <- df$age^2
mod2 <- lm(alcokol ~ over21 + age + age2,
           df)
```

⇒ Here, x has a symmetrical effect on both sides of the treatment.

Recode the x: assymmetric effect

We can assume x has different effects on each side of the treatment

$$y_i = \alpha + \rho D_i + \gamma x_i + \delta x_i D_i \quad (5)$$

Recode the x: assymetric effect


We can assume x has different effects on each side of the treatment

$$y_i = \alpha + \rho D_i + \gamma x_i + \delta x_i D_i \quad (5)$$

⇒ we center the x on the cutpoint ($x_i - c$) → ρ still reports the change at the cutpoint.

```
mod3 <- lm(alcohol ~ over21 * age,  
           df)
```

Do it in R

Legal drinking age 21 ($x = 0$) and alcohol related deaths

Extrapolation

We do this to estimate the effect at the cutpoint (ρ)

Extrapolation

We do this to estimate the effect at the cutpoint (ρ)

- ▶ but we can also extrapolate y beyond the cutpoint with x :
 $\rho + \delta(x - c)$

Recode the x

- ▶ recoding the x is a **the parametric approach**.

Recode the x

- ▶ recoding the x is a **the parametric approach**.
- ▶ subsetting the data to tweak the window around the cutpoint is a **non-parametric approach**.

Bandwidth : the idea

If the span of x around c is sufficiently small, there is no problem with non-linearity

Bandwidth : the idea

If the span of x around c is sufficiently small, there is no problem with non-linearity

- ▶ There's a tradeoff between linearity and statistical power (we need sufficient N).

Bandwidth : how do we choose it?

We try out different bandwidths

Bandwidth : how do we choose it?

We try out different bandwidths

- ▶ We can do it by hand

Bandwidth : how do we choose it?

We try out different bandwidths

- ▶ We can do it by hand

```
mod5 <- lm(alcohol ~ over21 * age,  
           df[df$agecell > 20 & df$agecell < 22,])
```

Bandwidth : how do we choose it?

We try out different bandwidths

- ▶ We can do it by hand

```
mod5 <- lm(alcohol ~ over21 * age,  
           df[df$agecell > 20 & df$agecell < 22,])
```

⇒ *When you narrow down, do you get a weaker or stronger effect?*

Bandwidth : how do we choose it?

We try out different bandwidths

- ▶ We can do it by hand

Bandwidth : how do we choose it?

We try out different bandwidths

- ▶ We can do it by hand
- ▶ ...or we can make an algorithm do it:
 - ▶ run a local weighted regression line
 - ▶ bandwidth is estimated accordingly

⇒ *the point is to show robustness, not p-hack!*

Omitted variable bias

We want to make certain that

Omitted variable bias

We want to make certain that

- ▶ D has an effect on y :

Omitted variable bias

We want to make certain that

► D has an effect on y :

→ is there really a cutpoint? Try out placebos!

Omitted variable bias

We want to make certain that

- ▶ D has an effect on y :

→ is there really a cutpoint? Try out placebos!

- ▶ treatment was indeed assigned at the cutpoint:

Omitted variable bias

We want to make certain that

- ▶ D has an effect on y :

→ is there really a cutpoint? Try out placebos!

- ▶ treatment was indeed assigned at the cutpoint:

→ is there unnatural clustering around one side?

Omitted variable bias

We want to make certain that

- ▶ D has an effect on y :

→ is there really a cutpoint? Try out placebos!

- ▶ treatment was indeed assigned at the cutpoint:

→ is there unnatural clustering around one side?

- ▶ treatment has impact on outcome but not other pre-treatment covariates

Omitted variable bias

We want to make certain that

- ▶ D has an effect on y :

→ is there really a cutpoint? Try out placebos!

- ▶ treatment was indeed assigned at the cutpoint:

→ is there unnatural clustering around one side?

- ▶ treatment has impact on outcome but not other pre-treatment covariates

→ check for balance/is there a similar “bump” for covariates? (bad news)

Omitted variable bias

We want to make certain that

- ▶ D has an effect on y :

→ is there really a cutpoint? Try out placebos!

- ▶ treatment was indeed assigned at the cutpoint:

→ is there unnatural clustering around one side?

- ▶ treatment has impact on outcome but not other pre-treatment covariates

→ check for balance/is there a similar “bump” for covariates? (bad news)

Fuzzy RD

Fuzzy RD

Often the D increases the probability of a treatment, but we don't know!

⇒ This is a Instrumental Variable approach (more on Thursday)

In brief

x **has a unique effect on** D . I'm interested in the effect of \bar{x} on y , but x is completely endogenous:

In brief

x **has a unique effect on** D . I'm interested in the effect of \bar{x} on y , but x is completely endogenous:

$$\blacktriangleright y = \alpha + \beta_1 \bar{x} + \beta_2 x + e$$

In brief

x **has a unique effect on** D . I'm interested in the effect of \bar{x} on y , but x is completely endogenous:

►
$$y = \alpha + \beta_1 \bar{x} + \beta_2 x + e$$

I use treatment as an instrument. We do this in two steps

In brief

x **has a unique effect on** D . I'm interested in the effect of \bar{x} on y , but x is completely endogenous:

$$\blacktriangleright y = \alpha + \beta_1 \bar{x} + \beta_2 x + e$$

I use treatment as an instrument. We do this in two steps

$$\blacktriangleright \text{step 1: } \bar{x} = \alpha_1 + \phi D + \beta_1 x + e_1$$

In brief

x **has a unique effect on** D . I'm interested in the effect of \bar{x} on y , but x is completely endogenous:

► $y = \alpha + \beta_1 \bar{x} + \beta_2 x + e$

I use treatment as an instrument. We do this in two steps

► step 1: $\bar{x} = \alpha_1 + \phi D + \beta_1 x + e_1$

► step 2: $y = \alpha_2 + \gamma \tilde{x} + \beta_2 x + e_2$

In brief

x **has a unique effect on** D . I'm interested in the effect of \bar{x} on y , but x is completely endogenous:

$$\blacktriangleright y = \alpha + \beta_1 \bar{x} + \beta_2 x + e$$

I use treatment as an instrument. We do this in two steps

$$\blacktriangleright \text{step 1: } \bar{x} = \alpha_1 + \phi D + \beta_1 x + e_1$$

$$\blacktriangleright \text{step 2: } y = \alpha_2 + \gamma \tilde{x} + \beta_2 x + e_2$$

$\Rightarrow \gamma$ is the causal effect of D in a fuzzy design.

Differences-in-differences

Definition: Comparing two differences

Definition: Differences-in-differences

Treatment and control groups may differ in many ways (they are not randomly assigned)

Definition: Differences-in-differences

Treatment and control groups may differ in many ways (they are not randomly assigned)

- ▶ Pre-treatment: They move in parallel

Definition: Differences-in-differences

Treatment and control groups may differ in many ways (they are not randomly assigned)

- ▶ Pre-treatment: They move in parallel
- ▶ Post-treatment: They diverge

Definition: Differences-in-differences

Treatment and control groups may differ in many ways (they are not randomly assigned)

- ▶ Pre-treatment: They move in parallel
- ▶ Post-treatment: They diverge

⇒ *Treatment effect is that difference*

What differences?

Diff-in-diff is based on two comparisons

What differences?

Diff-in-diff is based on two comparisons

- ▶ the difference pre- and post treatment *within* each unit

What differences?

Diff-in-diff is based on two comparisons

- ▶ the difference pre- and post treatment *within* each unit
- ▶ the difference *between* the treatment and control groups

What differences?

Diff-in-diff is based on two comparisons

- ▶ the difference pre- and post treatment *within* each unit
- ▶ the difference *between* the treatment and control groups

⇒ *based on panel data (units are observed several times).*

Example: States' monetary policy and number of banks

Take the differences between number of banks in two districts in Mississippi

Example: States' monetary policy and number of banks

Take the differences between number of banks in two districts in Mississippi

- ▶ Treatment: District 6 provided money to banks, while district 8 did not.

Example: States' monetary policy and number of banks

Take the differences between number of banks in two districts in Mississippi

- ▶ Treatment: District 6 provided money to banks, while district 8 did not.
- ▶ After a year district 6 had 121 banks, while district 8 had 132

Example: States' monetary policy and number of banks

Take the differences between number of banks in two districts in Mississippi

- ▶ Treatment: District 6 provided money to banks, while district 8 did not.
- ▶ After a year district 6 had 121 banks, while district 8 had 132

⇒ *What was the treatment effect?*

Example: States' monetary policy and number of banks

The two districts started out differently

Example: States' monetary policy and number of banks

The two districts started out differently

- ▶ within-unit difference: Number of banks before and after the crisis in each district.

District 6: $\$121 - 135 = \$ -14$; District 8: $\$132 - 165 = \-33

Example: States' monetary policy and number of banks

The two districts started out differently

- ▶ within-unit difference: Number of banks before and after the crisis in each district.

District 6: $\$121 - 135 = \$ -14$; District 8: $\$132 - 165 = \-33

- ▶ between-unit differences: take the difference between the two.

$$-14 - (-33) = 19$$

Example: States' monetary policy and number of banks

The two districts started out differently

- ▶ within-unit difference: Number of banks before and after the crisis in each district.

District 6: $\$121 - 135 = \$ -14$; District 8: $\$132 - 165 = \-33

- ▶ between-unit differences: take the difference between the two.

$$-14 - (-33) = 19$$

\Rightarrow *Basically a 2-by-2 table*

How to do it?

Interaction effects

In a regression, these differences are represented by an interaction term between two dummies

Interaction effects

In a regression, these differences are represented by an interaction term between two dummies

$$y_i = \alpha + \beta_1 T_i + \beta_2 P + \beta_3 T_i P_i \quad (6)$$

Interaction effects

In a regression, these differences are represented by an interaction term between two dummies

$$y_i = \alpha + \beta_1 T_i + \beta_2 P + \beta_3 T_i P_i \quad (6)$$

- P represents post-treatment effect: differences *within* units

Interaction effects

In a regression, these differences are represented by an interaction term between two dummies

$$y_i = \alpha + \beta_1 T_i + \beta_2 P + \beta_3 T_i P_i \quad (6)$$

- ▶ P represents post-treatment effect: differences *within* units
- ▶ T represents the treatment group: differences *between* units

Interaction effects

In a regression, these differences are represented by an interaction term between two dummies

$$y_i = \alpha + \beta_1 T_i + \beta_2 P + \beta_3 T_i P_i \quad (6)$$

- ▶ P represents post-treatment effect: differences *within* units
- ▶ T represents the treatment group: differences *between* units
- ▶ β_3 is the causal effect

Data

Data requirements

- ▶ Requires panel data

Data

Data requirements

- ▶ Requires panel data → which means correcting the standard errors.
- ▶ Common panel types: state-year/administrative unit-time period; people over time . . .

⇒ *we want to know the trend before and after the break*

Another example: drinking age and death

Another example: drinking age and death

Does the legal drinking age has an effect on death rates among the young?

- ▶ y is number of deaths per 100 000
- ▶ P is post-treatment dummy
- ▶ T is dummies for states
- ▶ $trend$ is year dummies

Another example: step 1 → calculate differences

The authors have two tricks:

- ▶ Hardcode the interaction effect (dummy before/after treatment)
- ▶ They remove the intercept to retain all dummies

```
load("df2.Rda")  
##with intercept  
mod <- lm(mrate ~ legal +  
           state +  
           year_fct,  
           df)  
##without intercept; with all dummies  
mod <- lm(mrate ~ 0 +  
           legal +  
           state +  
           year_fct,  
           df)
```

Another example: step 2 → calculate errors

Calculate robust standard errors:

```
library(clusterSandwich)
```

```
## Warning: package 'clusterSandwich' was built under R version 4
```

```
## Registered S3 method overwritten by 'clusterSandwich':
```

```
##   method      from
```

```
##   bread.mlm sandwich
```

```
vcov <- vcovCR(mod, cluster = df[["state"]],  
               type = "CR2")
```

```
robust <- coef_test(mod, vcov = vcov)$SE
```

Another example: step 3 → interpretation

Display the results and interpret:

```
library(stargazer)
stargazer(mod, se = robust,
           omit = "state|year",
           type = "html",
           out = "regtable.html")
```

Another example: step 3 → interpretation

Table 1: Death rates among young as a function of legal drinking age

	<i>Dependent variable:</i>
	mrate
Legal drinking age (causal effect)	10.804** (4.479)
Observations	714
R ²	0.986
Adjusted R ²	0.985
Residual Std. Error	17.339 (df = 649)
F Statistic	726.005*** (df = 65; 649)
<i>Note:</i> * p<0.1; ** p<0.05; *** p<0.01	

⇒ *What did we find?*

The parallel trends assumption

Main assumption

Units can be different, but – absent treatment – they must follow the same trend (hence the panel data).

Main assumption

Units can be different, but – absent treatment – they must follow the same trend (hence the panel data).

- ▶ The regression assumes a counterfactual

Main assumption

Units can be different, but – absent treatment – they must follow the same trend (hence the panel data).

- ▶ The regression assumes a counterfactual → remember the extrapolation.

Main assumption: The way around

When we have several treated and control units they can follow .

Main assumption: The way around

When we have several treated and control units they can follow .

- ▶ individual trend lines. . .

Main assumption: The way around

When we have several treated and control units they can follow .

- ▶ individual trend lines. . .
- ▶ . . . that are modelled as deviations from one unique trend

Main assumption: The way around

When we have several treated and control units they can follow .

- ▶ individual trend lines. . .
- ▶ . . . that are modelled as deviations from one unique trend

Main assumption: The way around

When we have several treated and control units they can follow .

- ▶ individual trend lines...
- ▶ ... that are modelled as deviations from one unique trend

⇒ *We do that with an interaction effect!*

```
mod <- lm(mrate ~ 0 +  
          legal +  
          state *  
          year_fct,  
          df)
```

Last fix

If our units are in fact several units (say, populations in states)

Last fix

If our units are in fact several units (say, populations in states)

- ▶ we can use weights

Last fix

If our units are in fact several units (say, populations in states)

► we can use weights

⇒ *There's a tradeoff: treatment is at the unit level, statistical power at the subunit level.*