

Where are we?

Introduction

Interpretation

Two sources of variation in the data

Study technique

Interpretation

Silje Synnøve Lyder Hermansen

15 februar 2026

Where are we?

Where are we?

Where are we?

- ▶ week 1:
 - ▶ purpose of the course
 - ▶ R as an object oriented language
- ▶ week 2:
 - ▶ dialects in R
 - ▶ descriptive statistics:
 - ▶ measurement level and choice of descriptives
 - ▶ data exploration
- ▶ week 3: (this week)
 - ▶ linear regression (OLS)
 - ▶ interpretation
 - ▶ non-linear effects

Plan for the day

- ▶ lecture: uncertainty and interpretation of linear models
 - ▶ substantive interest: the size of the effect
 - ▶ statistical significance: sources of variation/uncertainty
 - ▶ R notebook 1: interpretation (Gelman and Hill, King et al)
- ▶ study technique: how to use AI/LLMs in this class
- ▶ Thursday:
 - ▶ implementation in R
 - ▶ R notebook 2: non-linear effects (Berry et al)

Introduction

Today's example

What is the effect of electoral systems on parliamentarians resource allocation?

- ▶ Members of the European Parliament (MEPs) sit together in one institution, but run for election under different rules
- ▶ expectation: more local investment among MEPs in candidate-centered systems (compared to party-centered systems), because of their need for a personal brand
- ▶ variables:
 - ▶ y : number of constituency-level assistants employed (metric)
 - ▶ x : candidate vs. party-centered systems (binary)

Two views on linear regression

Two views on linear regression

Linear regression summarizes how the average values of a numerical outcome variable vary over subpopulations defined by linear functions of predictors. (Gelman and Hill, 2007, ch 3)

- ▶ **comparison of means:** descriptive approach to regression; makes sense for categorical predictors
- ▶ **relationship between variables:** their correlation; more causal, makes sense for numerical predictors

A comparison of means: group means

Most obvious when my predictor is categorical

```
df %>%  
  group_by(OpenList) %>%  
  reframe("mean_y" = mean(LocalAssistants)) %>%  
  ungroup %>%  
  mutate(diff = mean_y - lag(mean_y))
```

```
## # A tibble: 2 x 3  
##   OpenList mean_y   diff  
##   <int>   <dbl> <dbl>  
## 1     0     2.47    NA  
## 2     1     3.42   0.949
```

- ▶ MEPs from *party-centered* systems employ on average 2.47 local assistants
- ▶ MEPs from *candidate-centered* systems employ on average 3.42 local assistants.
- ▶ The difference is 0.95

A comparison of means: regression

Most obvious when my predictor is categorical

```
#Estimate the equation
mod <- lm(LocalAssistants ~ OpenList,
          df)
#Summarize the results
summary(mod)
```

```
##
## Call:
## lm(formula = LocalAssistants ~ OpenList, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -3.42  -2.42  -0.47   1.53  36.08
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    2.468     0.161   15.35 < 2e-16 ***
## OpenList        0.949     0.234    4.05 5.7e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.2 on 737 degrees of freedom
## Multiple R-squared:  0.0218, Adjusted R-squared:  0.0204
## F-statistic: 16.4 on 1 and 737 DF, p-value: 5.68e-05
```

- ▶ MEPs from *party-centered* systems employ on average 2.47 local assistants
- ▶ The *difference* is 0.95.
- ▶ MEPs from *candidate-centered* systems employ on average 2.47 + 0.95 = 3.42 local assistants.

Relationship between variables: regression

More descriptive statistics

```
mod2 <- lm(LocalAssistants ~ OpenList + LaborCost,
            df)
```

```
summary(mod2)
```

```
##
## Call:
## lm(formula = LocalAssistants ~ OpenList + LaborCost, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -4.49  -1.94  -0.41   1.08   35.00
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   4.1266     0.2861   14.42 < 2e-16 ***
## OpenList      0.8288     0.2278    3.64 0.00029 ***
## LaborCost    -0.0702     0.0102   -6.91 1e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.1 on 736 degrees of freedom
## Multiple R-squared:  0.0814, Adjusted R-squared:  0.0789
## F-statistic: 32.6 on 2 and 736 DF,  p-value: 2.69e-14
```

- ▶ the relationship (correlation)
- ▶ net of other variable's influence (controlling for...)
- ▶ the precision (uncertainty)
- ▶ the shared variation (R^2)
- ▶ the remaining variation (residuals, σ^2)

Interpretation

Stages of interpretaion

- ▶ **hypothesis testing:** direction and significance
- ▶ **marginal effect:** the relative increase in your predictor wo/accounting for the value of other preditors.
- ▶ **prediction:** fill in the equation for all predictors and calculate the predicted effect
- ▶ **first difference:** fill in the equation for two *scenarios* and calculate the difference in y
- ▶ **effect plot:** fill in the equation for all scenarios relevant to your predictor

⇒ *as we move to GLMs, the importance of stages 3-6 becomes important*

Hypothesis testing

Hypothesis testing

Hypotheses are mostly about direction and significance

```
summary(mod2)
```

```
##
## Call:
## lm(formula = LocalAssistants ~ OpenList + LaborCost, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -4.49  -1.94  -0.41   1.08   35.00
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   4.1266     0.2861   14.42 < 2e-16 ***
## OpenList      0.8288     0.2278    3.64 0.00029 ***
## LaborCost    -0.0702     0.0102   -6.91 1e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.1 on 736 degrees of freedom
## Multiple R-squared:  0.0814, Adjusted R-squared:  0.0789
## F-statistic: 32.6 on 2 and 736 DF, p-value: 2.69e-14
```

► **direction:** MEPs from candidate-centered systems have on average more local assistants on their payroll

► **significance:** this is unlikely to be random

⇒ ... *but what is the substantive effect?*

Marginal effect: change in x

Marginal effect: change in x

The relative (marginal) increase in your predictor (difference in means)

- ▶ without accounting for the value of other predictors
 - ▶ important once we move to GLMs
- ▶ regression is the estimation of an equation

$$y = \alpha + \beta x$$

- ▶ marginal effects focus on βx
 - ▶ β : from the model (you estimated it)
 - ▶ x : from the data (you pick it)

Marginal effect: example of change in x

The relative (marginal) increase in your predictor (difference in means) without accounting for the value of other predictors.

► interpretation:

```
summary(mod2)
```

```
##
## Call:
## lm(formula = LocalAssistants ~ OpenList + LaborCost, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -4.49  -1.94  -0.41   1.08  35.00
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   4.1266     0.2861   14.42 < 2e-16 ***
## OpenList      0.8288     0.2278    3.64 0.00029 ***
## LaborCost    -0.0702     0.0102   -6.91 1e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.1 on 736 degrees of freedom
## Multiple R-squared:  0.0814, Adjusted R-squared:  0.0789
## F-statistic: 32.6 on 2 and 736 DF, p-value: 2.69e-14
```

- β : “when x increases with one unit, y increases with β units”
- x : when labor cost increases with *one* unit (x , here 1000 euros), the average number of assistants decreases by 0.07

⇒ *but is this what we want to know?*

Partial scenario: set values for x

Find an increment (change) in x that makes sense for your story

```
##Summary of x
summary(df$LaborCost)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
##         4      10      26      23      31      41
```

```
##Find two typical values
summary(df$LaborCost)[c(4,5)]
```

```
##      Mean 3rd Qu.
##         23      31
```

```
## E.g. change from mean to 3rd quartile
summary(df$LaborCost)[c(4,5)] %>% diff
```

```
## 3rd Qu.
##      8.6
```

- ▶ univariate statistics / data exploration helps you find interesting changes in x
- ▶ calculate βx by filling in a realistic *change* in x .
- ▶ 8580 euro increase (increase by 8.58) corresponds to a 0.6 decrease in assistants ($\beta x = -0.07 \times 8.58$).

\Rightarrow use the univariate statistics to find an interesting increments

Prediction: fill in all x's

Prediction: fill in all x's

We estimated an equation with the help of our data

$$Y_i = \alpha + \beta_1 x_i + \beta_2 x_i$$

data (observed)

- ▶ variables: X and Y
- ▶ observations: i is a counter for the observations, refers to the i^{th} observation. $i \dots N$

parameters (estimated)

- ▶ α intercept, the value of Y when $X == 0$
- ▶ β slope, the increase in Y when X increases by one unit

We make predictions by filling in data points for that equation

$$Y_i = 4.13 + 0.83 \times OpenList + -0.07 \times LaborCost$$

If all x's were 1:

$$4.89 = 4.13 + 0.83 \times 1 + -0.07 \times 1$$

Why prediction?

- ▶ data description: “out-of-sample”
 - ▶ forecasting: e. g. election
 - ▶ machine learning: e.g create a new variable
- ▶ model statistics: “in-sample”
 - ▶ compare observed and predicted y
- ▶ interpretation:
 - ▶ set scenarios (fill in x)
 - ▶ predict outcomes (using β)

Creating one full scenario

You create a predicted scenario when you fill in values for *all* the predictors (x).

In R:

```
##Create variables
x1 = 1; x2 = 22

# or a data frame
scenario <- data.frame(
  OpenList = 1,
  LaborCost = 22)

# extract coefficients and apply to new data
predict(mod2, newdata = scenario)
```

```
## 1
## 3.4
```

By hand:

$$Y_i = \alpha + \beta_1 \text{OpenList} + \beta_2 \text{LaborCost}$$

$$Y_i = \alpha + \beta_1 \times 1 + \beta_2 \times 22$$

$$3.41 = 4.13 + 0.83 \times 1 + -0.07 \times 22$$

⇒ MEPs from candidate-centered electoral systems with average labor cost, are predicted to have – on average – a local staff of 3.41 people.

When would you be interested in full scenarios

When we use prediction for interpretation, we are interested in three metrics:

- ▶ two asymmetric scenarios: describe two typical value constellations (Ward and Ahlquist, ch 3)
- ▶ first difference: the difference in y between two predicted scenarios
- ▶ effect plots: the predicted y , as x increases, holding all other x constant.

First difference

First difference

First difference compares the predicted outcomes of two scenarios where one x changes, holding all other predictors constant

- ▶ first difference: difference between the two
- ▶ marginal effect vs first difference:
 - ▶ linear effects: marginal effect with partial scenario is the same as first difference
 - ▶ non-linear effects: the two are different

How to calculate a first difference

You create *two* scenarios and calculate the difference in y between the two

In R:

```
x1 = c(0, 1); x2 = 22

# or data frame
scenario <- data.frame(OpenList = c(0, 1),
                      LaborCost = 22)

#Predict both
predict(mod2, scenario)
```

```
##      1      2
## 2.6 3.4
```

```
#Take the difference
predict(mod2, scenario) %>% diff
```

```
##      2
## 0.83
```

By hand:

$$Y_i = \alpha + \beta_{OpenList_{1:2}} + \beta_{LaborCost}$$

$$\text{scenario 1: } 2.58 = 4.13 + 0.83 \times 0 + -0.07 \times 22$$

$$\text{scenario 1: } 3.41 = 4.13 + 0.83 \times 1 + -0.07 \times 22$$

$$\text{First difference: } 0.83 = 2.58 - 3.41$$

\Rightarrow The first difference can be calculated for any two scenarios of your choice!

Effect plot

Effect plot

Effect plots allow us to visualize our effects

- ▶ choice depends on the measurement level of x
 - ▶ categorical predictor: a tilted coefficient plot
 - ▶ numeric predictor: effectplot

Prediction using many scenarios

You create a bunch of scenarios covering the entire range of x

In R:

```
#Scenario
scenario <- data.frame(OpenList = c(0),
                       LaborCost = min(df$LaborCost): max(df$LaborCost))
#Inspect the first three scenarios
scenario[1:3,]
```

```
##   OpenList LaborCost
## 1         0        3.8
## 2         0        4.8
## 3         0        5.8
```


```
#Predict
scenario <- scenario %>% mutate(preds = predict(mod2, newdata = scenario))
scenario[1:3, ]
```

```
##   OpenList LaborCost preds
## 1         0        3.8  3.9
## 2         0        4.8  3.8
## 3         0        5.8  3.7
```

Visualize

Visualize by plotting your x against your predicted y

```
scenario %>%  
ggplot +  
  geom_line(aes(x = LaborCost,  
                y = preds))
```




Visualization for metric predictors

Metric predictors are best visualized with an effect plot (a line).

```
eff <- ggpredict(mod2,  
                 terms = "LaborCost")
```

```
eff %>% plot
```



Visualization: categorical predictor

Categorical predictors are best visualized with a titled coefficient plot



Two sources of variation in the data

Two sources of variation in the data

But are these effects statistically significant?

- ▶ **Fundamental uncertainty:** The natural randomness in outcomes, even if the true parameters were known (Captured by residual variance).
- ▶ **Estimation uncertainty:** How precisely are the coefficients estimated? (Captured by the variance-covariance matrix)

⇒ *the uncertainty of your predictions depend on both*

Fundamental uncertainty

Fundamental uncertainty

$$Y_i = \alpha + \beta X1_i + \beta X2_i + \sigma^2$$

data (observed)

- ▶ variables: X and Y
- ▶ observations: i is a counter for the observations, refers to the i^{th} observation. $i \dots N$

parameters (estimated)

- ▶ α intercept, the value of Y when $X == 0$
- ▶ β slope, the increase in Y when X increases by one unit
- ▶ σ^2 variance in the error term; $\sqrt{\sigma^2}$ = standard deviation

Let's rewrite

Any model in this class

$$Y \sim g(\theta, \sigma^2)$$

$$\theta = \alpha + \beta X_i$$

- ▶ $g()$: probability distribution (the link between parameters and outcomes)
- ▶ θ : central tendency (systematic component)
- ▶ σ^2 : spread around that systematic component

The normal (linear) model

$$Y_i \sim N(\mu_i, \sigma^2)$$

$$\mu_i = \alpha + \beta X_i$$

- ▶ $N()$: the normal distribution (identity link; linear; OLS)
- ▶ μ : mean predicted value
- ▶ σ^2 : residual variance (variation of Y around μ)


⇒ *focus on the distribution / link function*

What are the residuals?

We are always wrong in our predictions, but how wrong are we (in-sample)?

- ▶ **residuals:** difference between expected and observed outcomes ($Y_i - \mu_i$)
- ▶ a variable: one error / distance per observation in the data

```
df <- df %>% mutate(  
  #Predict in sample  
  preds = predict(mod2, newdata = .),  
  #Difference between expected and observed  
  residuals = LocalAssistants - preds  
)
```



How to describe the residuals?

We describe the residuals by their spread (standard deviation/residual standard error)

```
mean(df$residuals)
```

```
## [1] -9.8e-15
```

- ▶ mean: with an unbiased estimator, their average is 0

```
sd(df$residuals)
```

```
## [1] 3.1
```

- ▶ standard deviation: but their spread can be more or less high
- ▶ here, the average distance from their mean is a staff size of 3.08 local assistants.

⇒ *residual standard error*

Where is it reported?

```
summary(mod2)
```

```
##
## Call:
## lm(formula = LocalAssistants ~ OpenList + LaborCost, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -4.49  -1.94  -0.41   1.08   35.00
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   4.1266     0.2861   14.42 < 2e-16 ***
## OpenList       0.8288     0.2278    3.64 0.00029 ***
## LaborCost     -0.0702     0.0102   -6.91 1e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.1 on 736 degrees of freedom
## Multiple R-squared:  0.0814, Adjusted R-squared:  0.0789
## F-statistic: 32.6 on 2 and 736 DF, p-value: 2.69e-14
```

```
summary(mod2)$sigma
```

```
## [1] 3.1
```

- ▶ we may have predicted an average staff size of 3.86 per MEP in candidate-centered systems
- ▶ ... but individual MEPs typically deviate by 3.08 staffers
- ▶ from the variance, we can calculate the standard deviation, and thus prediction intervals around the mean prediction.

Conclusion: fundamental error

- ▶ important for predictions and model statistics
- ▶ not really for the uncertainty of the estimation of our effect

Estimation uncertainty

Estimation uncertainty

- ▶ most research is about the *effect of* x on y
- ▶ so, we're interested in the uncertainty of β

The central limit theorem and sampling

A fiction: the assumptions underpinning the uncertainty of the parameters

- ▶ assumption that data is a sample from a population
- ▶ we *could* sample many times
- ▶ we calculate the same parameter (e.g. mean, differences in means. . .) in each sample
- ▶ they will vary, but will follow a *normal distribution*

⇒ *each parameter is a distribution with a mean and a standard deviation*

Standard errors

```
summary(mod2)
```

```
##
## Call:
## lm(formula = LocalAssistants ~ OpenList + LaborCost, data = df)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -4.49  -1.94  -0.41   1.08   35.00
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)   4.1266     0.2861   14.42 < 2e-16 ***
## OpenList       0.8288     0.2278    3.64 0.00029 ***
## LaborCost     -0.0702     0.0102   -6.91 1e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 3.1 on 736 degrees of freedom
## Multiple R-squared:  0.0814, Adjusted R-squared:  0.0789
## F-statistic: 32.6 on 2 and 736 DF,  p-value: 2.69e-14
```

- ▶ mean: average of all the differences in means between the two groups of MEPs: 0.95
- ▶ spread: the standard deviation of this distribution is 0.23

⇒ *a standard error is the standard deviation of a hypothetical distribution (parameters)*

Colinearities

Colinearities

Regression parameters may be correlated

```
mat <-vcov(mod2)
mat
```

```
##              (Intercept) OpenList LaborCost
## (Intercept)      0.0819 -0.02846  -0.00244
## OpenList         -0.0285  0.05191   0.00018
## LaborCost        -0.0024  0.00018   0.00010
```

- ▶ reported in the *variance-covariance matrix*
- ▶ diagonal: the variance of the parameter.
 - ▶ variance in effect of electoral system: $\sigma^2 = 0.05$
 - ▶ standard error in effect of electoral system: $\sqrt{\sigma^2} = 0.23$
- ▶ off-diagonal: the covariance of the parameters
 - ▶ low correlation between labor cost and electoral system

Estimate

King et al. (2000) make two points

- ▶ find interesting scenarios when you interpret
- ▶ estimate the uncertainty for the scenarios including
 - ▶ standard error (diagonal)
 - ▶ covariance (off-diagonal)

⇒ *the correlation between variables may mean higher or lower uncertainty than only using the standard error*

Simulation

They do this using simulation

- ▶ set scenario for all predictors
- ▶ draw from the distribution of parameters
- ▶ make prediction
- ▶ repeat many times
- ▶ extract the information and report
 - ▶ mean
 - ▶ median
 - ▶ mode
 - ▶ standard deviation
 - ▶ plot the distribution!

Our class

We will see two ways of doing this in R

- ▶ `ggeffects` package: simulates scenarios for us and can be plotted seamlessly → *effect plots, coefplots and point predictions*
- ▶ `MASS` package: the “manual” simulation from a multivariate normal distribution using the variance-covariance matrix. → *entire vector of simulations; for other plots/purposes*

Study technique

For this class

- ▶ learn by doing!
 - ▶ all readings include R examples; code along!
 - ▶ my R notebooks
 - ▶ then play around with the concepts; also with your own data/former exams
- ▶ dialogue with AI (ChatGPT, Claude)

What to ask and not to ask chat for?

R codes

- ▶ dont ask for complex codes
 - ▶ requires quirey competence on your end
 - ▶ you don't learn
- ▶ ask it to annotate your scripts
 - ▶ explain what each line means
 - ▶ dissect all code chunks you find and ask

What to ask and not to ask chat for?

Statistics

- ▶ don't ask for a summary of the reading
 - ▶ it's not necessarily what we will focus on
 - ▶ you don't learn
- ▶ ask for definitions
 - ▶ ask it to define key concepts you don't understand while you read
 - ▶ rephrase definitions and ask it this is a good understanding
- ▶ match with your readings
 - ▶ upload the PDF and ask specific questions
 - ▶ ask for examples, possibly with R codes
- ▶ interpretation
 - ▶ copy-paste your model output and ask for an explainer
 - ▶ use descriptive statistics to find interesting scenarios, ask it to help you find a plain English intuitive sentence