

# Convex Optimization

Stephen Boyd   Lieven Vandenberghe

Revised slides by Stephen Boyd, Lieven Vandenberghe, and Parth Nobel

# 1. Introduction

# Outline

Mathematical optimization

Convex optimization

## Optimization problem

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & g_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- ▶  $x \in \mathbf{R}^n$  is (vector) variable to be chosen ( $n$  scalar variables  $x_1, \dots, x_n$ )
- ▶  $f_0$  is the **objective function**, to be minimized
- ▶  $f_1, \dots, f_m$  are the **inequality constraint functions**
- ▶  $g_1, \dots, g_p$  are the **equality constraint functions**
  
- ▶ variations: maximize objective, multiple objectives, ...

## Finding good (or best) actions

- ▶  $x$  represents some **action**, *e.g.*,
  - trades in a portfolio
  - airplane control surface deflections
  - schedule or assignment
  - resource allocation
- ▶ constraints limit actions or impose conditions on outcome
- ▶ the smaller the objective  $f_0(x)$ , the better
  - total cost (or negative profit)
  - deviation from desired or target outcome
  - risk
  - fuel use

## Finding good models

- ▶  $x$  represents the **parameters** in a model
- ▶ constraints impose requirements on model parameters (*e.g.*, nonnegativity)
- ▶ objective  $f_0(x)$  is sum of two terms:
  - a prediction error (or loss) on some observed data
  - a (regularization) term that penalizes model complexity

## Worst-case analysis (pessimization)

- ▶ variables are actions or parameters out of our control (and possibly under the control of an adversary)
- ▶ constraints limit the possible values of the parameters
- ▶ minimizing  $-f_0(x)$  finds **worst possible parameter values**
  
- ▶ if the worst possible value of  $f_0(x)$  is tolerable, you're OK
- ▶ it's good to know what the worst possible scenario can be

## Optimization-based models

- ▶ model an entity as taking actions that solve an optimization problem
  - an individual makes choices that maximize expected utility
  - an organism acts to maximize its reproductive success
  - reaction rates in a cell maximize growth
  - currents in a circuit minimize total power
- ▶ (except the last) these are **very crude** models
- ▶ and yet, they often work very well



## Basic use model for mathematical optimization

- ▶ instead of saying how to choose (action, model)  $x$
- ▶ you articulate what you want (by stating the problem)
- ▶ then let an algorithm decide on (action, model)  $x$

## Can you solve it?

- ▶ generally, no
- ▶ but you can try to solve it approximately, and it often doesn't matter
- ▶ the exception: **convex optimization**
  - includes linear programming (LP), quadratic programming (QP), many others
  - we can solve these problems reliably and efficiently
  - come up in many applications across many fields

## Nonlinear optimization

traditional techniques for general nonconvex problems involve compromises

### **local optimization methods** (nonlinear programming)

- ▶ find a point that minimizes  $f_0$  among feasible points near it
- ▶ can handle large problems, *e.g.*, neural network training
- ▶ require initial guess, and often, algorithm parameter tuning
- ▶ provide no information about how suboptimal the point found is

### **global optimization methods**

- ▶ find the (global) solution
- ▶ worst-case complexity grows exponentially with problem size
- ▶ often based on solving convex subproblems

# Outline

Mathematical optimization

Convex optimization

# Convex optimization

convex optimization problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

- ▶ variable  $x \in \mathbf{R}^n$
- ▶ equality constraints are linear
- ▶  $f_0, \dots, f_m$  are **convex**: for  $\theta \in [0, 1]$ ,

$$f_i(\theta x + (1 - \theta)y) \leq \theta f_i(x) + (1 - \theta)f_i(y)$$

*i.e.*,  $f_i$  have nonnegative (upward) curvature

## When is an optimization problem hard to solve?

- ▶ classical view:
  - linear (zero curvature) is easy
  - nonlinear (nonzero curvature) is hard
- ▶ the classical view is **wrong**
- ▶ the correct view:
  - convex (nonnegative curvature) is easy
  - nonconvex (negative curvature) is hard

## Solving convex optimization problems

- ▶ many different algorithms (that run on many platforms)
  - interior-point methods for up to 10000s of variables
  - first-order methods for larger problems
  - do not require initial point, babysitting, or tuning
- ▶ can develop and deploy quickly using modeling languages such as CVXPY
- ▶ solvers are reliable, so can be embedded
- ▶ code generation yields real-time solvers that execute in milliseconds (e.g., on Falcon 9 and Heavy for landing)

## Modeling languages for convex optimization

- ▶ domain specific languages (DSLs) for convex optimization
  - describe problem in high level language, close to the math
  - can automatically transform problem to standard form, then solve
- ▶ enables rapid prototyping
- ▶ it's now much easier to develop an optimization-based application
- ▶ ideal for teaching and research (can do a lot with short scripts)
- ▶ gets close to the basic idea: **say what you want, not how to get it**



## CVXPY example: non-negative least squares

math:

$$\begin{array}{ll}\text{minimize} & \|Ax - b\|_2^2 \\ \text{subject to} & x \geq 0\end{array}$$

- ▶ variable is  $x$
- ▶  $A, b$  given
- ▶  $x \geq 0$  means  $x_1 \geq 0, \dots, x_n \geq 0$

CVXPY code:

```
import cvxpy as cp

A, b = ...

x = cp.Variable(n)
obj = cp.norm2(A @ x - b)**2
constr = [x >= 0]
prob = cp.Problem(cp.Minimize(obj), constr)
prob.solve()
```

# Brief history of convex optimization

- ▶ **theory (convex analysis):** 1900–1970

- ▶ **algorithms**

- 1947: simplex algorithm for linear programming (Dantzig)
- 1960s: early interior-point methods (Fiacco & McCormick, Dikin, ...)
- 1970s: ellipsoid method and other subgradient methods
- 1980s & 90s: interior-point methods (Karmarkar, Nesterov & Nemirovski)
- since 2000s: many methods for large-scale convex optimization

- ▶ **applications**

- before 1990: mostly in operations research, a few in engineering
- since 1990: many applications in engineering (control, signal processing, communications, circuit design, ...)
- since 2000s: machine learning and statistics, finance

# Summary

convex optimization problems

- ▶ are optimization problems of a special form
- ▶ arise in many applications
- ▶ can be solved effectively
- ▶ are easy to specify using DSLs

## 2. Convex sets

# Outline

Some standard convex sets


Operations that preserve convexity

Generalized inequalities

Separating and supporting hyperplanes

## Affine set

**line** through  $x_1, x_2$ : all points of form  $x = \theta x_1 + (1 - \theta)x_2$ , with  $\theta \in \mathbf{R}$



**affine set:** contains the line through any two distinct points in the set

**example:** solution set of linear equations  $\{x \mid Ax = b\}$

(conversely, every affine set can be expressed as solution set of system of linear equations)


## Convex set

**line segment** between  $x_1$  and  $x_2$ : all points of form  $x = \theta x_1 + (1 - \theta)x_2$ , with  $0 \leq \theta \leq 1$

**convex set**: contains line segment between any two points in the set

$$x_1, x_2 \in C, \quad 0 \leq \theta \leq 1 \quad \implies \quad \theta x_1 + (1 - \theta)x_2 \in C$$

**examples** (one convex, two nonconvex sets)





## Convex combination and convex hull

**convex combination** of  $x_1, \dots, x_k$ : any point  $x$  of the form

$$x = \theta_1 x_1 + \theta_2 x_2 + \dots + \theta_k x_k$$

with  $\theta_1 + \dots + \theta_k = 1$ ,  $\theta_i \geq 0$

**convex hull**  $\text{conv } S$ : set of all convex combinations of points in  $S$






## Convex cone

**conic (nonnegative) combination** of  $x_1$  and  $x_2$ : any point of the form

$$x = \theta_1 x_1 + \theta_2 x_2$$


with  $\theta_1 \geq 0, \theta_2 \geq 0$




**convex cone**: set that contains all conic combinations of points in the set

## Hyperplanes and halfspaces

**hyperplane:** set of the form  $\{x \mid a^T x = b\}$ , with  $a \neq 0$



**halfspace:** set of the form  $\{x \mid a^T x \leq b\}$ , with  $a \neq 0$



- ▶  $a$  is the normal vector
- ▶ hyperplanes are affine and convex; halfspaces are convex

## Euclidean balls and ellipsoids


**(Euclidean) ball** with center  $x_c$  and radius  $r$ :

$$B(x_c, r) = \{x \mid \|x - x_c\|_2 \leq r\} = \{x_c + ru \mid \|u\|_2 \leq 1\}$$

**ellipsoid:** set of the form

$$\{x \mid (x - x_c)^T P^{-1} (x - x_c) \leq 1\}$$

with  $P \in \mathbf{S}_{++}^n$  (i.e.,  $P$  symmetric positive definite)



another representation:  $\{x_c + Au \mid \|u\|_2 \leq 1\}$  with  $A$  square and nonsingular


## Norm balls and norm cones

- ▶ **norm:** a function  $\|\cdot\|$  that satisfies
  - $\|x\| \geq 0$ ;  $\|x\| = 0$  if and only if  $x = 0$
  - $\|tx\| = |t| \|x\|$  for  $t \in \mathbf{R}$
  - $\|x + y\| \leq \|x\| + \|y\|$
- ▶ notation:  $\|\cdot\|$  is general (unspecified) norm;  $\|\cdot\|_{\text{symb}}$  is particular norm
- ▶ **norm ball** with center  $x_c$  and radius  $r$ :  $\{x \mid \|x - x_c\| \leq r\}$
- ▶ **norm cone:**  $\{(x, t) \mid \|x\| \leq t\}$
- ▶ norm balls and cones are convex

Euclidean norm cone

$$\{(x, t) \mid \|x\|_2 \leq t\} \subset \mathbf{R}^{n+1}$$

is called **second-order cone**




## Polyhedra

- **polyhedron** is solution set of finitely many linear inequalities and equalities

$$\{x \mid Ax \leq b, Cx = d\}$$

( $A \in \mathbf{R}^{m \times n}$ ,  $C \in \mathbf{R}^{p \times n}$ ,  $\leq$  is componentwise inequality)

- intersection of finite number of halfspaces and hyperplanes
- example with no equality constraints;  $a_i^T$  are rows of  $A$



## Positive semidefinite cone


notation:

- ▶  $\mathbf{S}^n$  is set of symmetric  $n \times n$  matrices
- ▶  $\mathbf{S}_+^n = \{X \in \mathbf{S}^n \mid X \geq 0\}$ : positive semidefinite (symmetric)  $n \times n$  matrices

$$X \in \mathbf{S}_+^n \iff z^T X z \geq 0 \text{ for all } z$$

- ▶  $\mathbf{S}_+^n$  is a convex cone, the **positive semidefinite cone**
- ▶  $\mathbf{S}_{++}^n = \{X \in \mathbf{S}^n \mid X > 0\}$ : positive definite (symmetric)  $n \times n$  matrices

example:  $\begin{bmatrix} x & y \\ y & z \end{bmatrix} \in \mathbf{S}_+^2$



# Outline

Some standard convex sets

Operations that preserve convexity

Generalized inequalities

Separating and supporting hyperplanes

## Showing a set is convex

methods for establishing convexity of a set  $C$



1. apply definition: show  $x_1, x_2 \in C, 0 \leq \theta \leq 1 \implies \theta x_1 + (1 - \theta)x_2 \in C$ 
  - recommended only for **very simple** sets
2. use convex functions (next lecture)
3. show that  $C$  is obtained from simple convex sets (hyperplanes, halfspaces, norm balls, ...) by operations that preserve convexity
  - intersection
  - affine mapping
  - perspective mapping
  - linear-fractional mapping

you'll mostly use methods 2 and 3



## Intersection

- ▶ the intersection of (any number of) convex sets is convex
- ▶ **example:**
  - $S = \{x \in \mathbf{R}^m \mid |p(t)| \leq 1 \text{ for } |t| \leq \pi/3\}$ , with  $p(t) = x_1 \cos t + \cdots + x_m \cos mt$
  - write  $S = \bigcap_{|t| \leq \pi/3} \{x \mid |p(t)| \leq 1\}$ , i.e., an intersection of (convex) slabs
- ▶ picture for  $m = 2$ :



## Affine mappings

- ▶ suppose  $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$  is affine, i.e.,  $f(x) = Ax + b$  with  $A \in \mathbf{R}^{m \times n}$ ,  $b \in \mathbf{R}^m$
- ▶ the **image** of a convex set under  $f$  is convex

$$S \subseteq \mathbf{R}^n \text{ convex} \implies f(S) = \{f(x) \mid x \in S\} \text{ convex}$$

- ▶ the **inverse image**  $f^{-1}(C)$  of a convex set under  $f$  is convex

$$C \subseteq \mathbf{R}^m \text{ convex} \implies f^{-1}(C) = \{x \in \mathbf{R}^n \mid f(x) \in C\} \text{ convex}$$

## Examples

- ▶ scaling, translation:  $aS + b = \{ax + b \mid x \in S\}$ ,  $a, b \in \mathbf{R}$
- ▶ projection onto some coordinates:  $\{x \mid (x, y) \in S\}$
- ▶ if  $S \subseteq \mathbf{R}^n$  is convex and  $c \in \mathbf{R}^n$ ,  $c^T S = \{c^T x \mid x \in S\}$  is an interval
- ▶ solution set of **linear matrix inequality**  $\{x \mid x_1 A_1 + \cdots + x_m A_m \leq B\}$  with  $A_i, B \in \mathbf{S}^p$
- ▶ hyperbolic cone  $\{x \mid x^T P x \leq (c^T x)^2, c^T x \geq 0\}$  with  $P \in \mathbf{S}_+^n$

## Perspective and linear-fractional function

- ▶ **perspective function**  $P : \mathbf{R}^{n+1} \rightarrow \mathbf{R}^n$ :

$$P(x, t) = x/t, \quad \text{dom } P = \{(x, t) \mid t > 0\}$$

- ▶ images and inverse images of convex sets under perspective are convex



- ▶ **linear-fractional function**  $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$ :

$$f(x) = \frac{Ax + b}{c^T x + d}, \quad \text{dom } f = \{x \mid c^T x + d > 0\}$$

- ▶ images and inverse images of convex sets under linear-fractional functions are convex

## Linear-fractional function example

$$f(x) = \frac{1}{x_1 + x_2 + 1}x$$



# Outline

Some standard convex sets

Operations that preserve convexity

Generalized inequalities

Separating and supporting hyperplanes

## Proper cones

a convex cone  $K \subseteq \mathbf{R}^n$  is a **proper cone** if

- ▶  $K$  is closed (contains its boundary)
- ▶  $K$  is solid (has nonempty interior)
- ▶  $K$  is pointed (contains no line)

### examples

- ▶ nonnegative orthant  $K = \mathbf{R}_+^n = \{x \in \mathbf{R}^n \mid x_i \geq 0, i = 1, \dots, n\}$
- ▶ positive semidefinite cone  $K = \mathbf{S}_+^n$
- ▶ nonnegative polynomials on  $[0, 1]$ :

$$K = \{x \in \mathbf{R}^n \mid x_1 + x_2 t + x_3 t^2 + \dots + x_n t^{n-1} \geq 0 \text{ for } t \in [0, 1]\}$$

## Generalized inequality

- ▶ (nonstrict and strict) **generalized inequality** defined by a proper cone  $K$ :

$$x \preceq_K y \iff y - x \in K, \quad x \prec_K y \iff y - x \in \text{int } K$$

- ▶ **examples**

- componentwise inequality ( $K = \mathbf{R}_+^n$ ):  $x \preceq_{\mathbf{R}_+^n} y \iff x_i \leq y_i, \quad i = 1, \dots, n$
- matrix inequality ( $K = \mathbf{S}_+^n$ ):  $X \preceq_{\mathbf{S}_+^n} Y \iff Y - X$  positive semidefinite

these two types are so common that we drop the subscript in  $\preceq_K$

- ▶ many properties of  $\preceq_K$  are similar to  $\leq$  on  $\mathbf{R}$ , e.g.,

$$x \preceq_K y, \quad u \preceq_K v \implies x + u \preceq_K y + v$$



# Outline

Some standard convex sets

Operations that preserve convexity


Generalized inequalities

Separating and supporting hyperplanes

## Separating hyperplane theorem

- ▶ if  $C$  and  $D$  are nonempty disjoint (i.e.,  $C \cap D = \emptyset$ ) convex sets, there exist  $a \neq 0$ ,  $b$  s.t.


$$a^T x \leq b \text{ for } x \in C, \quad a^T x \geq b \text{ for } x \in D$$



- ▶ the hyperplane  $\{x \mid a^T x = b\}$  **separates**  $C$  and  $D$
- ▶ strict separation requires additional assumptions (e.g.,  $C$  is closed,  $D$  is a singleton)

## Supporting hyperplane theorem

- ▶ suppose  $x_0$  is a boundary point of set  $C \subset \mathbf{R}^n$
- ▶ **supporting hyperplane** to  $C$  at  $x_0$  has form  $\{x \mid a^T x = a^T x_0\}$ , where  $a \neq 0$  and  $a^T x \leq a^T x_0$  for all  $x \in C$



- ▶ **supporting hyperplane theorem:** if  $C$  is convex, then there exists a supporting hyperplane at every boundary point of  $C$

### 3. Convex functions

# Outline

Convex functions

Operations that preserve convexity

Constructive convex analysis


Perspective and conjugate

Quasiconvexity

## Definition

- ▶  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex if  $\mathbf{dom} f$  is a convex set and for all  $x, y \in \mathbf{dom} f$ ,  $0 \leq \theta \leq 1$ ,

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$$



- ▶  $f$  is concave if  $-f$  is convex
- ▶  $f$  is strictly convex if  $\mathbf{dom} f$  is convex and for  $x, y \in \mathbf{dom} f$ ,  $x \neq y$ ,  $0 < \theta < 1$ ,

$$f(\theta x + (1 - \theta)y) < \theta f(x) + (1 - \theta)f(y)$$

## Examples on $\mathbf{R}$

convex functions:

- ▶ affine:  $ax + b$  on  $\mathbf{R}$ , for any  $a, b \in \mathbf{R}$
- ▶ exponential:  $e^{ax}$ , for any  $a \in \mathbf{R}$
- ▶ powers:  $x^\alpha$  on  $\mathbf{R}_{++}$ , for  $\alpha \geq 1$  or  $\alpha \leq 0$
- ▶ powers of absolute value:  $|x|^p$  on  $\mathbf{R}$ , for  $p \geq 1$
- ▶ positive part (relu):  $\max\{0, x\}$

concave functions:

- ▶ affine:  $ax + b$  on  $\mathbf{R}$ , for any  $a, b \in \mathbf{R}$
- ▶ powers:  $x^\alpha$  on  $\mathbf{R}_{++}$ , for  $0 \leq \alpha \leq 1$
- ▶ logarithm:  $\log x$  on  $\mathbf{R}_{++}$
- ▶ entropy:  $-x \log x$  on  $\mathbf{R}_{++}$
- ▶ negative part:  $\min\{0, x\}$

## Examples on $\mathbf{R}^n$

convex functions:

- ▶ affine functions:  $f(x) = a^T x + b$
- ▶ any norm, e.g., the  $\ell_p$  norms
  - $\|x\|_p = (|x_1|^p + \cdots + |x_n|^p)^{1/p}$  for  $p \geq 1$
  - $\|x\|_\infty = \max\{|x_1|, \dots, |x_n|\}$
- ▶ sum of squares:  $\|x\|_2^2 = x_1^2 + \cdots + x_n^2$
- ▶ max function:  $\max(x) = \max\{x_1, x_2, \dots, x_n\}$
- ▶ softmax or log-sum-exp function:  $\log(\exp x_1 + \cdots + \exp x_n)$



## Examples on $\mathbf{R}^{m \times n}$

- ▶  $X \in \mathbf{R}^{m \times n}$  ( $m \times n$  matrices) is the variable
- ▶ general affine function has form

$$f(X) = \mathbf{tr}(A^T X) + b = \sum_{i=1}^m \sum_{j=1}^n A_{ij} X_{ij} + b$$

for some  $A \in \mathbf{R}^{m \times n}$ ,  $b \in \mathbf{R}$

- ▶ spectral norm (maximum singular value) is convex

$$f(X) = \|X\|_2 = \sigma_{\max}(X) = (\lambda_{\max}(X^T X))^{1/2}$$

- ▶ log-determinant: for  $X \in \mathbf{S}_{++}^n$ ,  $f(X) = \log \det X$  is concave

## Extended-value extension

- ▶ suppose  $f$  is convex on  $\mathbf{R}^n$ , with domain  $\mathbf{dom} f$
- ▶ its extended-value extension  $\tilde{f}$  is function  $\tilde{f} : \mathbf{R}^n \rightarrow \mathbf{R} \cup \{\infty\}$

$$\tilde{f}(x) = \begin{cases} f(x) & x \in \mathbf{dom} f \\ \infty & x \notin \mathbf{dom} f \end{cases}$$

- ▶ often simplifies notation; for example, the condition

$$0 \leq \theta \leq 1 \implies \tilde{f}(\theta x + (1 - \theta)y) \leq \theta \tilde{f}(x) + (1 - \theta) \tilde{f}(y)$$

(as an inequality in  $\mathbf{R} \cup \{\infty\}$ ), means the same as the two conditions

- $\mathbf{dom} f$  is convex
- $x, y \in \mathbf{dom} f, 0 \leq \theta \leq 1 \implies f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$

## Restriction of a convex function to a line

- ▶  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex if and only if the function  $g : \mathbf{R} \rightarrow \mathbf{R}$ ,

$$g(t) = f(x + tv), \quad \mathbf{dom} \, g = \{t \mid x + tv \in \mathbf{dom} \, f\}$$

is convex (in  $t$ ) for any  $x \in \mathbf{dom} \, f$ ,  $v \in \mathbf{R}^n$

- ▶ can check convexity of  $f$  by checking convexity of functions of one variable

## Example

- ▶  $f : \mathbf{S}^n \rightarrow \mathbf{R}$  with  $f(X) = \log \det X$ ,  $\text{dom } f = \mathbf{S}_{++}^n$
- ▶ consider line in  $\mathbf{S}^n$  given by  $X + tV$ ,  $X \in \mathbf{S}_{++}^n$ ,  $V \in \mathbf{S}^n$ ,  $t \in \mathbf{R}$

$$\begin{aligned} g(t) &= \log \det(X + tV) \\ &= \log \det \left( X^{1/2} \left( I + tX^{-1/2} V X^{-1/2} \right) X^{1/2} \right) \\ &= \log \det X + \log \det \left( I + tX^{-1/2} V X^{-1/2} \right) \\ &= \log \det X + \sum_{i=1}^n \log(1 + t\lambda_i) \end{aligned}$$

where  $\lambda_i$  are the eigenvalues of  $X^{-1/2} V X^{-1/2}$

- ▶  $g$  is concave in  $t$  (for any choice of  $X \in \mathbf{S}_{++}^n$ ,  $V \in \mathbf{S}^n$ ); hence  $f$  is concave

## First-order condition

- ▶  $f$  is **differentiable** if  $\text{dom } f$  is open and the gradient


$$\nabla f(x) = \left( \frac{\partial f(x)}{\partial x_1}, \frac{\partial f(x)}{\partial x_2}, \dots, \frac{\partial f(x)}{\partial x_n} \right) \in \mathbf{R}^n$$

exists at each  $x \in \text{dom } f$

- ▶ **1st-order condition:** differentiable  $f$  with convex domain is convex if and only if

$$f(y) \geq f(x) + \nabla f(x)^T (y - x) \quad \text{for all } x, y \in \text{dom } f$$

- ▶ first order Taylor approximation of convex  $f$  is a **global underestimator** of  $f$



## Second-order conditions

- ▶  $f$  is **twice differentiable** if  $\text{dom } f$  is open and the Hessian  $\nabla^2 f(x) \in \mathbf{S}^n$ ,

$$\nabla^2 f(x)_{ij} = \frac{\partial^2 f(x)}{\partial x_i \partial x_j}, \quad i, j = 1, \dots, n,$$

exists at each  $x \in \text{dom } f$

- ▶ **2nd-order conditions:** for twice differentiable  $f$  with convex domain
  - $f$  is convex if and only if  $\nabla^2 f(x) \succeq 0$  for all  $x \in \text{dom } f$
  - if  $\nabla^2 f(x) \succ 0$  for all  $x \in \text{dom } f$ , then  $f$  is strictly convex

## Examples

- **quadratic function:**  $f(x) = (1/2)x^T Px + q^T x + r$  (with  $P \in \mathbf{S}^n$ )

$$\nabla f(x) = Px + q, \quad \nabla^2 f(x) = P$$

convex if  $P \succeq 0$  (concave if  $P \preceq 0$ )

- **least-squares objective:**  $f(x) = \|Ax - b\|_2^2$


$$\nabla f(x) = 2A^T(Ax - b), \quad \nabla^2 f(x) = 2A^T A$$

convex (for any  $A$ )

- **quadratic-over-linear:**  $f(x, y) = x^2/y, y > 0$

$$\nabla^2 f(x, y) = \frac{2}{y^3} \begin{bmatrix} y \\ -x \end{bmatrix} \begin{bmatrix} y \\ -x \end{bmatrix}^T \succeq 0$$

convex for  $y > 0$



## More examples

- **log-sum-exp:**  $f(x) = \log \sum_{k=1}^n \exp x_k$  is convex

$$\nabla^2 f(x) = \frac{1}{\mathbf{1}^T z} \mathbf{diag}(z) - \frac{1}{(\mathbf{1}^T z)^2} z z^T \quad (z_k = \exp x_k)$$

- to show  $\nabla^2 f(x) \succeq 0$ , we must verify that  $v^T \nabla^2 f(x) v \geq 0$  for all  $v$ :

$$v^T \nabla^2 f(x) v = \frac{(\sum_k z_k v_k^2)(\sum_k z_k) - (\sum_k v_k z_k)^2}{(\sum_k z_k)^2} \geq 0$$


since  $(\sum_k v_k z_k)^2 \leq (\sum_k z_k v_k^2)(\sum_k z_k)$  (from Cauchy-Schwarz inequality)

- **geometric mean:**  $f(x) = (\prod_{k=1}^n x_k)^{1/n}$  on  $\mathbf{R}_{++}^n$  is concave (similar proof as above)



## Epigraph and sublevel set

- ▶  $\alpha$ -**sublevel set** of  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is  $C_\alpha = \{x \in \mathbf{dom} f \mid f(x) \leq \alpha\}$
- ▶ sublevel sets of convex functions are convex sets (but converse is false)
- ▶ **epigraph** of  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is  $\mathbf{epi} f = \{(x, t) \in \mathbf{R}^{n+1} \mid x \in \mathbf{dom} f, f(x) \leq t\}$



- ▶  $f$  is convex if and only if  $\mathbf{epi} f$  is a convex set

## Jensen's inequality

- ▶ **basic inequality:** if  $f$  is convex, then for  $x, y \in \mathbf{dom} f$ ,  $0 \leq \theta \leq 1$ ,

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y)$$

- ▶ **extension:** if  $f$  is convex and  $z$  is a random variable on  $\mathbf{dom} f$ ,

$$f(\mathbf{E} z) \leq \mathbf{E} f(z)$$

- ▶ basic inequality is special case with discrete distribution

$$\mathbf{prob}(z = x) = \theta, \quad \mathbf{prob}(z = y) = 1 - \theta$$


## Example: log-normal random variable

- ▶ suppose  $X \sim \mathcal{N}(\mu, \sigma^2)$
- ▶ with  $f(u) = \exp u$ ,  $Y = f(X)$  is log-normal
- ▶ we have  $\mathbf{E}f(X) = \exp(\mu + \sigma^2/2)$
- ▶ Jensen's inequality is

$$f(\mathbf{E}X) = \exp \mu \leq \mathbf{E}f(X) = \exp(\mu + \sigma^2/2)$$

which indeed holds since  $\exp \sigma^2/2 > 1$

## Example: log-normal random variable



# Outline

Convex functions

Operations that preserve convexity

Constructive convex analysis

Perspective and conjugate

Quasiconvexity

## Showing a function is convex

methods for establishing convexity of a function  $f$

1. verify definition (often simplified by restricting to a line)
2. for twice differentiable functions, show  $\nabla^2 f(x) \succeq 0$ 
  - recommended only for **very simple** functions
3. show that  $f$  is obtained from simple convex functions by operations that preserve convexity
  - nonnegative weighted sum
  - composition with affine function
  - pointwise maximum and supremum
  - composition
  - minimization
  - perspective

you'll mostly use methods 2 and 3

## Nonnegative scaling, sum, and integral

- ▶ **nonnegative multiple:**  $\alpha f$  is convex if  $f$  is convex,  $\alpha \geq 0$
- ▶ **sum:**  $f_1 + f_2$  convex if  $f_1, f_2$  convex
- ▶ **infinite sum:** if  $f_1, f_2, \dots$  are convex functions, infinite sum  $\sum_{i=1}^{\infty} f_i$  is convex
- ▶ **integral:** if  $f(x, \alpha)$  is convex in  $x$  for each  $\alpha \in \mathcal{A}$ , then  $\int_{\alpha \in \mathcal{A}} f(x, \alpha) d\alpha$  is convex
- ▶ there are analogous rules for concave functions

## Composition with affine function

**(pre-)composition with affine function:**  $f(Ax + b)$  is convex if  $f$  is convex

### examples

- ▶ log barrier for linear inequalities

$$f(x) = - \sum_{i=1}^m \log(b_i - a_i^T x), \quad \text{dom } f = \{x \mid a_i^T x < b_i, i = 1, \dots, m\}$$

- ▶ norm approximation error:  $f(x) = \|Ax - b\|$  (any norm)



## Pointwise maximum

if  $f_1, \dots, f_m$  are convex, then  $f(x) = \max\{f_1(x), \dots, f_m(x)\}$  is convex

### examples

- ▶ piecewise-linear function:  $f(x) = \max_{i=1, \dots, m} (a_i^T x + b_i)$
- ▶ sum of  $r$  largest components of  $x \in \mathbf{R}^n$ :

$$f(x) = x_{[1]} + x_{[2]} + \dots + x_{[r]}$$

( $x_{[i]}$  is  $i$ th largest component of  $x$ )

proof:  $f(x) = \max\{x_{i_1} + x_{i_2} + \dots + x_{i_r} \mid 1 \leq i_1 < i_2 < \dots < i_r \leq n\}$

## Pointwise supremum

if  $f(x, y)$  is convex in  $x$  for each  $y \in \mathcal{A}$ , then  $g(x) = \sup_{y \in \mathcal{A}} f(x, y)$  is convex

### examples

- ▶ distance to farthest point in a set  $C$ :  $f(x) = \sup_{y \in C} \|x - y\|$
- ▶ maximum eigenvalue of symmetric matrix: for  $X \in \mathbf{S}^n$ ,  $\lambda_{\max}(X) = \sup_{\|y\|_2=1} y^T X y$  is convex
- ▶ support function of a set  $C$ :  $S_C(x) = \sup_{y \in C} y^T x$  is convex

## Partial minimization

- ▶ the function  $g(x) = \inf_{y \in C} f(x, y)$  is called the **partial minimization** of  $f$  (w.r.t.  $y$ )
- ▶ if  $f(x, y)$  is convex in  $(x, y)$  and  $C$  is a convex set, then partial minimization  $g$  is convex

### examples

- ▶  $f(x, y) = x^T A x + 2x^T B y + y^T C y$  with

$$\begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \succeq 0, \quad C \succ 0$$

minimizing over  $y$  gives  $g(x) = \inf_y f(x, y) = x^T (A - B C^{-1} B^T) x$   
 $g$  is convex, hence Schur complement  $A - B C^{-1} B^T \succeq 0$

- ▶ distance to a set: **dist** $(x, S) = \inf_{y \in S} \|x - y\|$  is convex if  $S$  is convex

## Composition with scalar functions

- ▶ composition of  $g : \mathbf{R}^n \rightarrow \mathbf{R}$  and  $h : \mathbf{R} \rightarrow \mathbf{R}$  is  $f(x) = h(g(x))$  (written as  $f = h \circ g$ )
- ▶ composition  $f$  is convex if
  - $g$  convex,  $h$  convex,  $\tilde{h}$  nondecreasing
  - or  $g$  concave,  $h$  convex,  $\tilde{h}$  nonincreasing(monotonicity must hold for extended-value extension  $\tilde{h}$ )
- ▶ proof (for  $n = 1$ , differentiable  $g, h$ )

$$f''(x) = h''(g(x))g'(x)^2 + h'(g(x))g''(x)$$

### examples

- ▶  $f(x) = \exp g(x)$  is convex if  $g$  is convex
- ▶  $f(x) = 1/g(x)$  is convex if  $g$  is concave and positive

## General composition rule

- ▶ composition of  $g : \mathbf{R}^n \rightarrow \mathbf{R}^k$  and  $h : \mathbf{R}^k \rightarrow \mathbf{R}$  is  $f(x) = h(g(x)) = h(g_1(x), g_2(x), \dots, g_k(x))$
- ▶  $f$  is convex if  $h$  is convex and for each  $i$  one of the following holds
  - $g_i$  convex,  $\tilde{h}$  nondecreasing in its  $i$ th argument
  - $g_i$  concave,  $\tilde{h}$  nonincreasing in its  $i$ th argument
  - $g_i$  affine
- ▶ you will use this composition rule **constantly** throughout this course
- ▶ you need to commit this rule to memory

## Examples

- ▶  $\log \sum_{i=1}^m \exp g_i(x)$  is convex if  $g_i$  are convex
- ▶  $f(x) = p(x)^2/q(x)$  is convex if
  - $p$  is nonnegative and convex
  - $q$  is positive and concave
- ▶ composition rule subsumes others, *e.g.*,
  - $\alpha f$  is convex if  $f$  is, and  $\alpha \geq 0$
  - sum of convex (concave) functions is convex (concave)
  - max of convex functions is convex
  - min of concave functions is concave

# Outline

Convex functions

Operations that preserve convexity

**Constructive convex analysis**

Perspective and conjugate

Quasiconvexity

## Constructive convexity verification

- ▶ start with function  $f$  given as **expression**
- ▶ build parse tree for expression
  - leaves are variables or constants
  - nodes are functions of child expressions
- ▶ use composition rule to tag subexpressions as convex, concave, affine, or none
- ▶ if root node is labeled convex (concave), then  $f$  is convex (concave)
- ▶ extension: tag sign of each expression, and use sign-dependent monotonicity
  
- ▶ this is sufficient to show  $f$  is convex (concave), but not necessary
- ▶ this method for checking convexity (concavity) is readily automated



## Example

the function


$$f(x, y) = \frac{(x - y)^2}{1 - \max(x, y)}, \quad x < 1, \quad y < 1$$

is convex

constructive analysis:

- ▶ (leaves)  $x$ ,  $y$ , and 1 are affine
- ▶  $\max(x, y)$  is convex;  $x - y$  is affine
- ▶  $1 - \max(x, y)$  is concave
- ▶ function  $u^2/v$  is convex, monotone decreasing in  $v$  for  $v > 0$
- ▶  $f$  is composition of  $u^2/v$  with  $u = x - y$ ,  $v = 1 - \max(x, y)$ , hence convex

## Example (from dcp.stanford.edu)



## Disciplined convex programming

in **disciplined convex programming** (DCP) users construct convex and concave functions as expressions using constructive convex analysis

- ▶ expressions formed from
  - **variables**,
  - **constants**,
  - and **atomic functions** from a library
- ▶ atomic functions have known convexity, monotonicity, and sign properties
- ▶ all subexpressions match general composition rule
- ▶ a valid DCP function is
  - convex-by-construction
  - ‘syntactically’ convex (can be checked ‘locally’)
- ▶ convexity depends only on attributes of atomic functions, not their meanings
  - e.g., could swap  $\sqrt{\cdot}$  and  $\sqrt[4]{\cdot}$ , or  $\exp \cdot$  and  $(\cdot)_+$ , since their attributes match

## CVXPY example

$$\frac{(x-y)^2}{1-\max(x,y)}, \quad x < 1, \quad y < 1$$

```
import cvxpy as cp
x = cp.Variable()
y = cp.Variable()
expr = cp.quad_over_lin(x - y, 1 - cp.maximum(x, y))
expr.curvature # Convex
expr.sign # Positive
expr.is_dcp() # True
```

(atom `quad_over_lin(u,v)` includes domain constraint `v>0`)

## DCP is only sufficient

- ▶ consider convex function  $f(x) = \sqrt{1+x^2}$
- ▶ expression `f1 = cp.sqrt(1+cp.square(x))` is **not** DCP
- ▶ expression `f2 = cp.norm2([1,x])` **is** DCP
- ▶ CVXPY will not recognize `f1` as convex, even though it represents a convex function

# Outline

Convex functions

Operations that preserve convexity

Constructive convex analysis

**Perspective and conjugate**

Quasiconvexity

## Perspective

- ▶ the **perspective** of a function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is the function  $g : \mathbf{R}^n \times \mathbf{R} \rightarrow \mathbf{R}$ ,

$$g(x, t) = tf(x/t), \quad \text{dom } g = \{(x, t) \mid x/t \in \text{dom } f, t > 0\}$$


- ▶  $g$  is convex if  $f$  is convex

### examples

- ▶  $f(x) = x^T x$  is convex; so  $g(x, t) = x^T x/t$  is convex for  $t > 0$
- ▶  $f(x) = -\log x$  is convex; so relative entropy  $g(x, t) = t \log t - t \log x$  is convex on  $\mathbf{R}_{++}^2$

## Conjugate function

- ▶ the **conjugate** of a function  $f$  is  $f^*(y) = \sup_{x \in \text{dom}_f} (y^T x - f(x))$



- ▶  $f^*$  is convex (even if  $f$  is not)
- ▶ will be useful in chapter 5



## Examples

- ▶ negative logarithm  $f(x) = -\log x$

$$f^*(y) = \sup_{x>0} (xy + \log x) = \begin{cases} -1 - \log(-y) & y < 0 \\ \infty & \text{otherwise} \end{cases}$$

- ▶ strictly convex quadratic,  $f(x) = (1/2)x^T Qx$  with  $Q \in \mathbf{S}_{++}^n$

$$f^*(y) = \sup_x (y^T x - (1/2)x^T Qx) = \frac{1}{2}y^T Q^{-1}y$$

# Outline

Convex functions

Operations that preserve convexity

Constructive convex analysis

Perspective and conjugate


Quasiconvexity

## Quasiconvex functions

- $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is **quasiconvex** if  $\text{dom} f$  is convex and the sublevel sets

$$S_\alpha = \{x \in \text{dom} f \mid f(x) \leq \alpha\}$$

are convex for all  $\alpha$



- $f$  is **quasiconcave** if  $-f$  is quasiconvex
- $f$  is **quasilinear** if it is quasiconvex and quasiconcave

## Examples

- ▶  $\sqrt{|x|}$  is quasiconvex on  $\mathbf{R}$
- ▶  $\text{ceil}(x) = \inf\{z \in \mathbf{Z} \mid z \geq x\}$  is quasilinear
- ▶  $\log x$  is quasilinear on  $\mathbf{R}_{++}$
- ▶  $f(x_1, x_2) = x_1 x_2$  is quasiconcave on  $\mathbf{R}_{++}^2$
- ▶ linear-fractional function

$$f(x) = \frac{a^T x + b}{c^T x + d}, \quad \text{dom } f = \{x \mid c^T x + d > 0\}$$

is quasilinear

## Example: Internal rate of return

- ▶ cash flow  $x = (x_0, \dots, x_n)$ ;  $x_i$  is payment in period  $i$  (to us if  $x_i > 0$ )
- ▶ we assume  $x_0 < 0$  (i.e., an initial investment) and  $x_0 + x_1 + \dots + x_n > 0$
- ▶ **net present value** (NPV) of cash flow  $x$ , for interest rate  $r$ , is  $PV(x, r) = \sum_{i=0}^n (1+r)^{-i} x_i$
- ▶ **internal rate of return** (IRR) is smallest interest rate for which  $PV(x, r) = 0$ :

$$IRR(x) = \inf\{r \geq 0 \mid PV(x, r) = 0\}$$

- ▶ IRR is quasiconcave: superlevel set is intersection of open halfspaces

$$IRR(x) \geq R \iff \sum_{i=0}^n (1+r)^{-i} x_i > 0 \text{ for } 0 \leq r < R$$


## Properties of quasiconvex functions

- ▶ **modified Jensen inequality:** for quasiconvex  $f$

$$0 \leq \theta \leq 1 \implies f(\theta x + (1 - \theta)y) \leq \max\{f(x), f(y)\}$$

- ▶ **first-order condition:** differentiable  $f$  with convex domain is quasiconvex if and only if

$$f(y) \leq f(x) \implies \nabla f(x)^T (y - x) \leq 0$$



- ▶ **sum** of quasiconvex functions is not necessarily quasiconvex

## 4. Convex optimization problems

# Outline

## Optimization problems

Some standard convex problems

Transforming problems

Disciplined convex programming

Geometric programming

Quasiconvex optimization

Multicriterion optimization



## Optimization problem in standard form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- ▶  $x \in \mathbf{R}^n$  is the optimization variable
- ▶  $f_0 : \mathbf{R}^n \rightarrow \mathbf{R}$  is the objective or cost function
- ▶  $f_i : \mathbf{R}^n \rightarrow \mathbf{R}, i = 1, \dots, m$ , are the inequality constraint functions
- ▶  $h_i : \mathbf{R}^n \rightarrow \mathbf{R}$  are the equality constraint functions


## Feasible and optimal points

- ▶  $x \in \mathbf{R}^n$  is **feasible** if  $x \in \text{dom} f_0$  and it satisfies the constraints
- ▶ **optimal value** is  $p^\star = \inf\{f_0(x) \mid f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p\}$
- ▶  $p^\star = \infty$  if problem is infeasible
- ▶  $p^\star = -\infty$  if problem is **unbounded below**
- ▶ a feasible  $x$  is **optimal** if  $f_0(x) = p^\star$
- ▶  $X_{\text{opt}}$  is the set of optimal points

## Locally optimal points

$x$  is **locally optimal** if there is an  $R > 0$  such that  $x$  is optimal for


$$\begin{array}{ll}\text{minimize (over } z) & f_0(z) \\ \text{subject to} & f_i(z) \leq 0, \quad i = 1, \dots, m, \quad h_i(z) = 0, \quad i = 1, \dots, p \\ & \|z - x\|_2 \leq R\end{array}$$




## Examples

examples with  $n = 1$ ,  $m = p = 0$


- ▶  $f_0(x) = 1/x$ ,  $\text{dom } f_0 = \mathbf{R}_{++}$ :  $p^\star = 0$ , no optimal point
- ▶  $f_0(x) = -\log x$ ,  $\text{dom } f_0 = \mathbf{R}_{++}$ :  $p^\star = -\infty$
- ▶  $f_0(x) = x \log x$ ,  $\text{dom } f_0 = \mathbf{R}_{++}$ :  $p^\star = -1/e$ ,  $x = 1/e$  is optimal
- ▶  $f_0(x) = x^3 - 3x$ :  $p^\star = -\infty$ ,  $x = 1$  is locally optimal




$$f_0(x) = 1/x$$



$$f_0(x) = -\log x$$



$$f_0(x) = x \log x$$



$$f_0(x) = x^3 - 3x$$

## Implicit and explicit constraints

standard form optimization problem has **implicit constraint**

$$x \in \mathcal{D} = \bigcap_{i=0}^m \text{dom } f_i \cap \bigcap_{i=1}^p \text{dom } h_i,$$

- ▶ we call  $\mathcal{D}$  the **domain** of the problem
- ▶ the constraints  $f_i(x) \leq 0$ ,  $h_i(x) = 0$  are the **explicit constraints**
- ▶ a problem is **unconstrained** if it has no explicit constraints ( $m = p = 0$ )

**example:**

$$\text{minimize } f_0(x) = -\sum_{i=1}^k \log(b_i - a_i^T x)$$

is an unconstrained problem with implicit constraints  $a_i^T x < b_i$

## Feasibility problem

$$\begin{array}{ll}\text{find} & x \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

can be considered a special case of the general problem with  $f_0(x) = 0$ :

$$\begin{array}{ll}\text{minimize} & 0 \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- ▶  $p^\star = 0$  if constraints are feasible; any feasible  $x$  is optimal
- ▶  $p^\star = \infty$  if constraints are infeasible

## Standard form convex optimization problem

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & a_i^T x = b_i, \quad i = 1, \dots, p\end{array}$$

- ▶ objective and inequality constraints  $f_0, f_1, \dots, f_m$  are convex
- ▶ equality constraints are affine, often written as  $Ax = b$
- ▶ feasible and optimal sets of a convex optimization problem are convex
- ▶ problem is **quasiconvex** if  $f_0$  is quasiconvex,  $f_1, \dots, f_m$  are convex,  $h_1, \dots, h_p$  are affine

## Example

- ▶ standard form problem

$$\begin{array}{ll}\text{minimize} & f_0(x) = x_1^2 + x_2^2 \\ \text{subject to} & f_1(x) = x_1/(1+x_2^2) \leq 0 \\ & h_1(x) = (x_1+x_2)^2 = 0\end{array}$$

- ▶  $f_0$  is convex; feasible set  $\{(x_1, x_2) \mid x_1 = -x_2 \leq 0\}$  is convex
- ▶ not a convex problem (by our definition) since  $f_1$  is not convex,  $h_1$  is not affine
- ▶ equivalent (but not identical) to the convex problem

$$\begin{array}{ll}\text{minimize} & x_1^2 + x_2^2 \\ \text{subject to} & x_1 \leq 0 \\ & x_1 + x_2 = 0\end{array}$$



## Local and global optima

any locally optimal point of a convex problem is (globally) optimal

**proof:**

- ▶ suppose  $x$  is locally optimal, but there exists a feasible  $y$  with  $f_0(y) < f_0(x)$
- ▶  $x$  locally optimal means there is an  $R > 0$  such that


$$z \text{ feasible, } \|z - x\|_2 \leq R \implies f_0(z) \geq f_0(x)$$

- ▶ consider  $z = \theta y + (1 - \theta)x$  with  $\theta = R/(2\|y - x\|_2)$
- ▶  $\|y - x\|_2 > R$ , so  $0 < \theta < 1/2$
- ▶  $z$  is a convex combination of two feasible points, hence also feasible
- ▶  $\|z - x\|_2 = R/2$  and  $f_0(z) \leq \theta f_0(y) + (1 - \theta)f_0(x) < f_0(x)$ , which contradicts our assumption that  $x$  is locally optimal

## Optimality criterion for differentiable $f_0$

- $x$  is optimal for a convex problem if and only if it is feasible and

$$\nabla f_0(x)^T (y - x) \geq 0 \text{ for all feasible } y$$



- if nonzero,  $\nabla f_0(x)$  defines a supporting hyperplane to feasible set  $X$  at  $x$

## Examples

- ▶ **unconstrained problem:**  $x$  minimizes  $f_0(x)$  if and only if  $\nabla f_0(x) = 0$
- ▶ **equality constrained problem:**  $x$  minimizes  $f_0(x)$  subject to  $Ax = b$  if and only if there exists a  $\nu$  such that

$$Ax = b, \quad \nabla f_0(x) + A^T \nu = 0$$

- ▶ **minimization over nonnegative orthant:**  $x$  minimizes  $f_0(x)$  over  $\mathbf{R}_+^n$  if and only if

$$x \geq 0, \quad \begin{cases} \nabla f_0(x)_i \geq 0 & x_i = 0 \\ \nabla f_0(x)_i = 0 & x_i > 0 \end{cases}$$

# Outline

Optimization problems

Some standard convex problems

Transforming problems

Disciplined convex programming

Geometric programming


Quasiconvex optimization

Multicriterion optimization

## Linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x + d \\ \text{subject to} & Gx \leq h \\ & Ax = b\end{array}$$

- ▶ convex problem with affine objective and constraint functions
- ▶ feasible set is a polyhedron



## Example: Diet problem

- ▶ choose nonnegative quantities  $x_1, \dots, x_n$  of  $n$  foods
- ▶ one unit of food  $j$  costs  $c_j$  and contains amount  $A_{ij}$  of nutrient  $i$
- ▶ healthy diet requires nutrient  $i$  in quantity at least  $b_i$
- ▶ to find cheapest healthy diet, solve

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \geq b, \quad x \geq 0\end{array}$$

- ▶ express in standard LP form as

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & \begin{bmatrix} -A \\ -I \end{bmatrix} x \leq \begin{bmatrix} -b \\ 0 \end{bmatrix}\end{array}$$

## Example: Piecewise-linear minimization

- ▶ minimize convex piecewise-linear function  $f_0(x) = \max_{i=1,\dots,m}(a_i^T x + b_i)$ ,  $x \in \mathbf{R}^n$
- ▶ equivalent to LP


$$\begin{array}{ll}\text{minimize} & t \\ \text{subject to} & a_i^T x + b_i \leq t, \quad i = 1, \dots, m\end{array}$$

with variables  $x \in \mathbf{R}^n$ ,  $t \in \mathbf{R}$

- ▶ constraints describe  $\text{epi } f_0$

## Example: Chebyshev center of a polyhedron

**Chebyshev center** of  $\mathcal{P} = \{x \mid a_i^T x \leq b_i, i = 1, \dots, m\}$  is center of largest inscribed ball  $\mathcal{B} = \{x_c + u \mid \|u\|_2 \leq r\}$



- ▶  $a_i^T x \leq b_i$  for all  $x \in \mathcal{B}$  if and only if

$$\sup\{a_i^T (x_c + u) \mid \|u\|_2 \leq r\} = a_i^T x_c + r\|a_i\|_2 \leq b_i$$

- ▶ hence,  $x_c, r$  can be determined by solving LP with variables  $x_c, r$


$$\begin{array}{ll} \text{maximize} & r \\ \text{subject to} & a_i^T x_c + r\|a_i\|_2 \leq b_i, \quad i = 1, \dots, m \end{array}$$



## Quadratic program (QP)

$$\begin{array}{ll}\text{minimize} & (1/2)x^T Px + q^T x + r \\ \text{subject to} & Gx \leq h \\ & Ax = b\end{array}$$

- ▶  $P \in \mathbf{S}_+^n$ , so objective is convex quadratic
- ▶ minimize a convex quadratic function over a polyhedron



## Example: Least squares

- ▶ **least squares** problem: minimize  $\|Ax - b\|_2^2$
- ▶ analytical solution  $x^\star = A^\dagger b$  ( $A^\dagger$  is pseudo-inverse)
- ▶ can add linear constraints, *e.g.*,
  - $x \geq 0$  (**nonnegative least squares**)
  - $x_1 \leq x_2 \leq \dots \leq x_n$  (**isotonic regression**)

## Example: Linear program with random cost

- ▶ LP with random cost  $c$ , with mean  $\bar{c}$  and covariance  $\Sigma$
- ▶ hence, LP objective  $c^T x$  is random variable with mean  $\bar{c}^T x$  and variance  $x^T \Sigma x$
- ▶ **risk-averse** problem:

$$\begin{array}{ll}\text{minimize} & \mathbf{E} c^T x + \gamma \mathbf{var}(c^T x) \\ \text{subject to} & Gx \leq h, \quad Ax = b\end{array}$$

- ▶  $\gamma > 0$  is **risk aversion parameter**; controls the trade-off between expected cost and variance (risk)
- ▶ express as QP

$$\begin{array}{ll}\text{minimize} & \bar{c}^T x + \gamma x^T \Sigma x \\ \text{subject to} & Gx \leq h, \quad Ax = b\end{array}$$

## Quadratically constrained quadratic program (QCQP)

$$\begin{array}{ll}\text{minimize} & (1/2)x^T P_0 x + q_0^T x + r_0 \\ \text{subject to} & (1/2)x^T P_i x + q_i^T x + r_i \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

- ▶  $P_i \in \mathbf{S}_+^n$ ; objective and constraints are convex quadratic
- ▶ if  $P_1, \dots, P_m \in \mathbf{S}_{++}^n$ , feasible region is intersection of  $m$  ellipsoids and an affine set

## Second-order cone programming

$$\begin{array}{ll}\text{minimize} & f^T x \\ \text{subject to} & \|A_i x + b_i\|_2 \leq c_i^T x + d_i, \quad i = 1, \dots, m \\ & Fx = g\end{array}$$

$$(A_i \in \mathbf{R}^{n_i \times n}, F \in \mathbf{R}^{p \times n})$$

- ▶ inequalities are called second-order cone (SOC) constraints:

$$(A_i x + b_i, c_i^T x + d_i) \in \text{second-order cone in } \mathbf{R}^{n_i+1}$$

- ▶ for  $n_i = 0$ , reduces to an LP; if  $c_i = 0$ , reduces to a QCQP
- ▶ more general than QCQP and LP

## Example: Robust linear programming

suppose constraint vectors  $a_i$  are uncertain in the LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i, \quad i = 1, \dots, m,\end{array}$$

two common approaches to handling uncertainty

- ▶ **deterministic worst-case:** constraints must hold for all  $a_i \in \mathcal{E}_i$  (uncertainty ellipsoids)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i \text{ for all } a_i \in \mathcal{E}_i, \quad i = 1, \dots, m,\end{array}$$

- ▶ **stochastic:**  $a_i$  is random variable; constraints must hold with probability  $\eta$

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & \mathbf{prob}(a_i^T x \leq b_i) \geq \eta, \quad i = 1, \dots, m\end{array}$$

## Deterministic worst-case approach

- uncertainty ellipsoids are  $\mathcal{E}_i = \{\bar{a}_i + P_i u \mid \|u\|_2 \leq 1\}$ , ( $\bar{a}_i \in \mathbf{R}^n$ ,  $P_i \in \mathbf{R}^{n \times n}$ )
- center of  $\mathcal{E}_i$  is  $\bar{a}_i$ ; semi-axes determined by singular values/vectors of  $P_i$
- robust LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i \quad \forall a_i \in \mathcal{E}_i, \quad i = 1, \dots, m\end{array}$$

- equivalent to SOCP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & \bar{a}_i^T x + \|P_i^T x\|_2 \leq b_i, \quad i = 1, \dots, m\end{array}$$

(follows from  $\sup_{\|u\|_2 \leq 1} (\bar{a}_i + P_i u)^T x = \bar{a}_i^T x + \|P_i^T x\|_2$ )

## Stochastic approach

- ▶ assume  $a_i \sim \mathcal{N}(\bar{a}_i, \Sigma_i)$
- ▶  $a_i^T x \sim \mathcal{N}(\bar{a}_i^T x, x^T \Sigma_i x)$ , so

$$\mathbf{prob}(a_i^T x \leq b_i) = \Phi\left(\frac{b_i - \bar{a}_i^T x}{\|\Sigma_i^{1/2} x\|_2}\right)$$

where  $\Phi(u) = (1/\sqrt{2\pi}) \int_{-\infty}^u e^{-t^2/2} dt$  is  $\mathcal{N}(0, 1)$  CDF

- ▶  $\mathbf{prob}(a_i^T x \leq b_i) \geq \eta$  can be expressed as  $\bar{a}_i^T x + \Phi^{-1}(\eta) \|\Sigma_i^{1/2} x\|_2 \leq b_i$
- ▶ for  $\eta \geq 1/2$ , robust LP equivalent to SOCP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & \bar{a}_i^T x + \Phi^{-1}(\eta) \|\Sigma_i^{1/2} x\|_2 \leq b_i, \quad i = 1, \dots, m \end{array}$$



## Conic form problem

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Fx + g \preceq_K 0 \\ & Ax = b\end{array}$$

- ▶ constraint  $Fx + g \preceq_K 0$  involves a generalized inequality with respect to a proper cone  $K$
- ▶ linear programming is a conic form problem with  $K = \mathbf{R}_+^m$
- ▶ as with standard convex problem
  - feasible and optimal sets are convex
  - any local optimum is global

## Semidefinite program (SDP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & x_1 F_1 + x_2 F_2 + \cdots + x_n F_n + G \preceq 0 \\ & Ax = b\end{array}$$

with  $F_i, G \in \mathbf{S}^k$

- ▶ inequality constraint is called **linear matrix inequality** (LMI)
- ▶ includes problems with multiple LMI constraints: for example,

$$x_1 \hat{F}_1 + \cdots + x_n \hat{F}_n + \hat{G} \preceq 0, \quad x_1 \tilde{F}_1 + \cdots + x_n \tilde{F}_n + \tilde{G} \preceq 0$$

is equivalent to single LMI

$$x_1 \begin{bmatrix} \hat{F}_1 & 0 \\ 0 & \tilde{F}_1 \end{bmatrix} + x_2 \begin{bmatrix} \hat{F}_2 & 0 \\ 0 & \tilde{F}_2 \end{bmatrix} + \cdots + x_n \begin{bmatrix} \hat{F}_n & 0 \\ 0 & \tilde{F}_n \end{bmatrix} + \begin{bmatrix} \hat{G} & 0 \\ 0 & \tilde{G} \end{bmatrix} \preceq 0$$

## Example: Matrix norm minimization

$$\text{minimize} \quad \|A(x)\|_2 = (\lambda_{\max}(A(x)^T A(x)))^{1/2}$$

where  $A(x) = A_0 + x_1 A_1 + \cdots + x_n A_n$  (with given  $A_i \in \mathbf{R}^{p \times q}$ )

equivalent SDP

$$\begin{array}{ll} \text{minimize} & t \\ \text{subject to} & \begin{bmatrix} tI & A(x) \\ A(x)^T & tI \end{bmatrix} \succeq 0 \end{array}$$

- ▶ variables  $x \in \mathbf{R}^n, t \in \mathbf{R}$
- ▶ constraint follows from

$$\begin{aligned} \|A\|_2 \leq t & \iff A^T A \leq t^2 I, \quad t \geq 0 \\ & \iff \begin{bmatrix} tI & A \\ A^T & tI \end{bmatrix} \succeq 0 \end{aligned}$$

# Outline

Optimization problems

Some standard convex problems

Transforming problems

Disciplined convex programming

Geometric programming

Quasiconvex optimization

Multicriterion optimization

## Change of variables

- ▶  $\phi : \mathbf{R}^n \rightarrow \mathbf{R}^n$  is one-to-one with  $\phi(\text{dom } \phi) \supseteq \mathcal{D}$
- ▶ consider (possibly non-convex) problem

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

- ▶ change variables to  $z$  with  $x = \phi(z)$
- ▶ can solve equivalent problem

$$\begin{array}{ll}\text{minimize} & \tilde{f}_0(z) \\ \text{subject to} & \tilde{f}_i(z) \leq 0, \quad i = 1, \dots, m \\ & \tilde{h}_i(z) = 0, \quad i = 1, \dots, p\end{array}$$

where  $\tilde{f}_i(z) = f_i(\phi(z))$  and  $\tilde{h}_i(z) = h_i(\phi(z))$

- ▶ recover original optimal point as  $x^\star = \phi(z^\star)$

## Example

- ▶ **non-convex** problem

$$\begin{array}{ll}\text{minimize} & x_1/x_2 + x_3/x_1 \\ \text{subject to} & x_2/x_3 + x_1 \leq 1\end{array}$$

with implicit constraint  $x > 0$

- ▶ change variables using  $x = \phi(z) = \exp z$  to get

$$\begin{array}{ll}\text{minimize} & \exp(z_1 - z_2) + \exp(z_3 - z_1) \\ \text{subject to} & \exp(z_2 - z_3) + \exp(z_1) \leq 1\end{array}$$

which is **convex**

## Transformation of objective and constraint functions

suppose

- ▶  $\phi_0$  is monotone increasing
- ▶  $\psi_i(u) \leq 0$  if and only if  $u \leq 0$ ,  $i = 1, \dots, m$
- ▶  $\varphi_i(u) = 0$  if and only if  $u = 0$ ,  $i = 1, \dots, p$

standard form optimization problem is equivalent to

$$\begin{array}{ll}\text{minimize} & \phi_0(f_0(x)) \\ \text{subject to} & \psi_i(f_i(x)) \leq 0, \quad i = 1, \dots, m \\ & \varphi_i(h_i(x)) = 0, \quad i = 1, \dots, p\end{array}$$

example: minimizing  $\|Ax - b\|$  is equivalent to minimizing  $\|Ax - b\|^2$

## Converting maximization to minimization

- ▶ suppose  $\phi_0$  is monotone decreasing
- ▶ the maximization problem

$$\begin{array}{ll}\text{maximize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

is equivalent to the minimization problem

$$\begin{array}{ll}\text{minimize} & \phi_0(f_0(x)) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

### ▶ examples:

- $\phi_0(u) = -u$  transforms maximizing a concave function to minimizing a convex function
- $\phi_0(u) = 1/u$  transforms maximizing a concave positive function to minimizing a convex function



## Eliminating equality constraints

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

is equivalent to

$$\begin{array}{ll}\text{minimize (over } z) & f_0(Fz + x_0) \\ \text{subject to} & f_i(Fz + x_0) \leq 0, \quad i = 1, \dots, m\end{array}$$

where  $F$  and  $x_0$  are such that  $Ax = b \iff x = Fz + x_0$  for some  $z$

## Introducing equality constraints

$$\begin{array}{ll}\text{minimize} & f_0(A_0x + b_0) \\ \text{subject to} & f_i(A_ix + b_i) \leq 0, \quad i = 1, \dots, m\end{array}$$

is equivalent to

$$\begin{array}{ll}\text{minimize (over } x, y_i) & f_0(y_0) \\ \text{subject to} & f_i(y_i) \leq 0, \quad i = 1, \dots, m \\ & y_i = A_ix + b_i, \quad i = 0, 1, \dots, m\end{array}$$

## Introducing slack variables for linear inequalities

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & a_i^T x \leq b_i, \quad i = 1, \dots, m\end{array}$$

is equivalent to

$$\begin{array}{ll}\text{minimize (over } x, s) & f_0(x) \\ \text{subject to} & a_i^T x + s_i = b_i, \quad i = 1, \dots, m \\ & s_i \geq 0, \quad i = 1, \dots, m\end{array}$$

## Epigraph form

standard form convex problem is equivalent to

$$\begin{array}{ll}\text{minimize (over } x, t) & t \\ \text{subject to} & f_0(x) - t \leq 0 \\ & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

## Minimizing over some variables

$$\begin{array}{ll}\text{minimize} & f_0(x_1, x_2) \\ \text{subject to} & f_i(x_1) \leq 0, \quad i = 1, \dots, m\end{array}$$

is equivalent to

$$\begin{array}{ll}\text{minimize} & \tilde{f}_0(x_1) \\ \text{subject to} & f_i(x_1) \leq 0, \quad i = 1, \dots, m\end{array}$$

where  $\tilde{f}_0(x_1) = \inf_{x_2} f_0(x_1, x_2)$

## LP and SOCP as SDP

### LP and equivalent SDP

$$\begin{array}{ll} \text{LP:} & \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \end{array} \end{array} \qquad \begin{array}{ll} \text{SDP:} & \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & \mathbf{diag}(Ax - b) \leq 0 \end{array} \end{array}$$

(note different interpretation of generalized inequalities  $\leq$  in LP and SDP)

### SOCP and equivalent SDP

$$\begin{array}{ll} \text{SOCP:} & \begin{array}{ll} \text{minimize} & f^T x \\ \text{subject to} & \|A_i x + b_i\|_2 \leq c_i^T x + d_i, \quad i = 1, \dots, m \end{array} \end{array}$$

$$\begin{array}{ll} \text{SDP:} & \begin{array}{ll} \text{minimize} & f^T x \\ \text{subject to} & \begin{bmatrix} (c_i^T x + d_i)I & A_i x + b_i \\ (A_i x + b_i)^T & c_i^T x + d_i \end{bmatrix} \succeq 0, \quad i = 1, \dots, m \end{array} \end{array}$$

## Convex relaxation

- ▶ start with **nonconvex problem**: minimize  $h(x)$  subject to  $x \in C$
- ▶ find convex function  $\hat{h}$  with  $\hat{h}(x) \leq h(x)$  for all  $x \in \text{dom } h$  (i.e., a pointwise lower bound on  $h$ )
- ▶ find set  $\hat{C} \supseteq C$  (e.g.,  $\hat{C} = \text{conv } C$ ) described by linear equalities and convex inequalities

$$\hat{C} = \{x \mid f_i(x) \leq 0, \ i = 1, \dots, m, \ f_m(x) \leq 0, \ Ax = b\}$$

- ▶ convex problem

$$\begin{array}{ll} \text{minimize} & \hat{h}(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m, \quad Ax = b \end{array}$$

is a **convex relaxation** of the original problem

- ▶ optimal value of relaxation is lower bound on optimal value of original problem

## Example: Boolean LP

- ▶ **mixed integer linear program (MILP):**

$$\begin{array}{ll}\text{minimize} & c^T(x, z) \\ \text{subject to} & F(x, z) \leq g, \quad A(x, z) = b, \quad z \in \{0, 1\}^q\end{array}$$

with variables  $x \in \mathbf{R}^n, z \in \mathbf{R}^q$

- ▶  $z_i$  are called **Boolean variables**
- ▶ this problem is in general hard to solve
- ▶ **LP relaxation:** replace  $z \in \{0, 1\}^q$  with  $z \in [0, 1]^q$
- ▶ optimal value of relaxation LP is lower bound on MILP
- ▶ can use as heuristic for approximately solving MILP, e.g., **relax and round**



# Outline

Optimization problems

Some standard convex problems

Transforming problems

**Disciplined convex programming**

Geometric programming

Quasiconvex optimization

Multicriterion optimization

## Disciplined convex program

- ▶ specify objective as
  - minimize {scalar convex expression}, or
  - maximize {scalar concave expression}
- ▶ specify constraints as
  - {convex expression}  $\leq$  {concave expression} or
  - {concave expression}  $\geq$  {convex expression} or
  - {affine expression}  $=$  {affine expression}
- ▶ curvature of expressions are DCP certified, *i.e.*, follow composition rule
- ▶ DCP-compliant problems can be automatically transformed to standard forms, then solved

## CVXPY example

math:

$$\begin{array}{ll}\text{minimize} & \|x\|_1 \\ \text{subject to} & Ax = b \\ & \|x\|_\infty \leq 1\end{array}$$

- ▶  $x$  is the variable
- ▶  $A, b$  are given

CVXPY code:


```
import cvxpy as cp

A, b = ...

x = cp.Variable(n)
obj = cp.norm(x, 1)
constr = [
    A @ x == b,
    cp.norm(x, 'inf') <= 1,
]
prob = cp.Problem(cp.Minimize(obj), constr)
prob.solve()
```

## How CVXPY works

- ▶ starts with your optimization problem  $\mathcal{P}_1$
- ▶ finds a sequence of equivalent problems  $\mathcal{P}_2, \dots, \mathcal{P}_N$
- ▶ final problem  $\mathcal{P}_N$  matches a standard form (e.g., LP, QP, SOCP, or SDP)
- ▶ calls a specialized solver on  $\mathcal{P}_N$
- ▶ retrieves solution of original problem by reversing the transformations



# Outline

Optimization problems

Some standard convex problems

Transforming problems

Disciplined convex programming

**Geometric programming**

Quasiconvex optimization

Multicriterion optimization

## Geometric programming

- **monomial function:**

$$f(x) = cx_1^{a_1} x_2^{a_2} \cdots x_n^{a_n}, \quad \text{dom } f = \mathbf{R}_{++}^n$$

with  $c > 0$ ; exponent  $a_i$  can be any real number

- **posynomial function:** sum of monomials

$$f(x) = \sum_{k=1}^K c_k x_1^{a_{1k}} x_2^{a_{2k}} \cdots x_n^{a_{nk}}, \quad \text{dom } f = \mathbf{R}_{++}^n$$

- **geometric program (GP)**

$$\begin{array}{ll} \text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 1, \quad i = 1, \dots, m \\ & h_i(x) = 1, \quad i = 1, \dots, p \end{array}$$

with  $f_i$  posynomial,  $h_i$  monomial

## Geometric program in convex form

- ▶ change variables to  $y_i = \log x_i$ , and take logarithm of cost, constraints
- ▶ monomial  $f(x) = cx_1^{a_1} \cdots x_n^{a_n}$  transforms to

$$\log f(e^{y_1}, \dots, e^{y_n}) = a^T y + b \quad (b = \log c)$$

- ▶ posynomial  $f(x) = \sum_{k=1}^K c_k x_1^{a_{1k}} x_2^{a_{2k}} \cdots x_n^{a_{nk}}$  transforms to

$$\log f(e^{y_1}, \dots, e^{y_n}) = \log \left( \sum_{k=1}^K e^{a_k^T y + b_k} \right) \quad (b_k = \log c_k)$$

- ▶ geometric program transforms to convex problem

$$\begin{aligned} & \text{minimize} && \log \left( \sum_{k=1}^K \exp(a_{0k}^T y + b_{0k}) \right) \\ & \text{subject to} && \log \left( \sum_{k=1}^K \exp(a_{ik}^T y + b_{ik}) \right) \leq 0, \quad i = 1, \dots, m \\ & && Gy + d = 0 \end{aligned}$$

## Examples: Frobenius norm diagonal scaling

- ▶ we seek diagonal matrix  $D = \mathbf{diag}(d)$ ,  $d > 0$ , to minimize  $\|DMD^{-1}\|_F^2$
- ▶ express as

$$\|DMD^{-1}\|_F^2 = \sum_{i,j=1}^n \left( DMD^{-1} \right)_{ij}^2 = \sum_{i,j=1}^n M_{ij}^2 d_i^2 / d_j^2$$

- ▶ a posynomial in  $d$  (with exponents 0, 2, and  $-2$ )
- ▶ in convex form, with  $y = \log d$ ,

$$\log \|DMD^{-1}\|_F^2 = \log \left( \sum_{i,j=1}^n \exp \left( 2(y_i - y_j + \log |M_{ij}|) \right) \right)$$



# Outline

Optimization problems

Some standard convex problems

Transforming problems

Disciplined convex programming

Geometric programming

**Quasiconvex optimization**


Multicriterion optimization

## Quasiconvex optimization

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

with  $f_0 : \mathbf{R}^n \rightarrow \mathbf{R}$  quasiconvex,  $f_1, \dots, f_m$  convex

can have locally optimal points that are not (globally) optimal



## Linear-fractional program

- ▶ linear-fractional program

$$\begin{array}{ll}\text{minimize} & (c^T x + d)/(e^T x + f) \\ \text{subject to} & Gx \leq h, \quad Ax = b\end{array}$$

with variable  $x$  and implicit constraint  $e^T x + f > 0$

- ▶ equivalent to the LP (with variables  $y, z$ )

$$\begin{array}{ll}\text{minimize} & c^T y + dz \\ \text{subject to} & Gy \leq hz, \quad Ay = bz \\ & e^T y + fz = 1, \quad z \geq 0\end{array}$$

- ▶ recover  $x^\star = y^\star / z^\star$

## Von Neumann model of a growing economy

- ▶  $x, x^+ \in \mathbf{R}_{++}^n$ : activity levels of  $n$  economic sectors, in current and next period
- ▶  $(Ax)_i$ : amount of good  $i$  produced in current period
- ▶  $(Bx^+)_i$ : amount of good  $i$  consumed in next period
- ▶  $Bx^+ \leq Ax$ : goods consumed next period no more than produced this period
- ▶  $x_i^+/x_i$ : growth rate of sector  $i$
- ▶ allocate activity to maximize growth rate of slowest growing sector

$$\begin{array}{ll} \text{maximize (over } x, x^+) & \min_{i=1, \dots, n} x_i^+/x_i \\ \text{subject to} & x^+ \geq 0, \quad Bx^+ \leq Ax \end{array}$$

- ▶ a quasiconvex problem with variables  $x, x^+$

## Convex representation of sublevel sets

- ▶ if  $f_0$  is quasiconvex, there exists a family of functions  $\phi_t$  such that:
  - $\phi_t(x)$  is convex in  $x$  for fixed  $t$
  - $t$ -sublevel set of  $f_0$  is 0-sublevel set of  $\phi_t$ , i.e.,  $f_0(x) \leq t \iff \phi_t(x) \leq 0$

### example:

- ▶  $f_0(x) = p(x)/q(x)$ , with  $p$  convex and nonnegative,  $q$  concave and positive
- ▶ take  $\phi_t(x) = p(x) - tq(x)$ : for  $t \geq 0$ ,
  - $\phi_t$  convex in  $x$
  - $p(x)/q(x) \leq t$  if and only if  $\phi_t(x) \leq 0$

## Bisection method for quasiconvex optimization

- ▶ for fixed  $t$ , consider convex feasibility problem

$$\phi_t(x) \leq 0, \quad f_i(x) \leq 0, \quad i = 1, \dots, m, \quad Ax = b \quad (1)$$

if feasible, we can conclude that  $t \geq p^\star$ ; if infeasible,  $t \leq p^\star$

- ▶ bisection method:

---

**given**  $l \leq p^\star, u \geq p^\star$ , tolerance  $\epsilon > 0$ .

**repeat**

1.  $t := (l + u)/2$ .
2. Solve the convex feasibility problem (1).
3. **if** (1) is feasible,  $u := t$ ; **else**  $l := t$ .

**until**  $u - l \leq \epsilon$ .

---

- ▶ requires exactly  $\lceil \log_2((u - l)/\epsilon) \rceil$  iterations

# Outline

Optimization problems

Some standard convex problems

Transforming problems

Disciplined convex programming

Geometric programming

Quasiconvex optimization

**Multicriterion optimization**

## Multicriterion optimization

- ▶ **multicriterion** or **multi-objective** problem:

$$\begin{array}{ll}\text{minimize} & f_0(x) = (F_1(x), \dots, F_q(x)) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m, \quad Ax = b\end{array}$$

- ▶ objective is the **vector**  $f_0(x) \in \mathbf{R}^q$
- ▶  $q$  different objectives  $F_1, \dots, F_q$ ; roughly speaking we want all  $F_i$ 's to be small
- ▶ feasible  $x^\star$  is **optimal** if  $y$  feasible  $\implies f_0(x^\star) \leq f_0(y)$
- ▶ this means that  $x^\star$  simultaneously minimizes each  $F_i$ ; the objectives are **noncompeting**
- ▶ not surprisingly, this doesn't happen very often





## Pareto optimality

- ▶ feasible  $x$  **dominates** another feasible  $\tilde{x}$  if  $f_0(x) \leq f_0(\tilde{x})$  and for at least one  $i$ ,  $F_i(x) < F_i(\tilde{x})$
- ▶ *i.e.*,  $x$  meets  $\tilde{x}$  on all objectives, and beats it on at least one
  
- ▶ feasible  $x^{\text{po}}$  is **Pareto optimal** if it is not dominated by any feasible point
- ▶ can be expressed as:  $y$  feasible,  $f_0(y) \leq f_0(x^{\text{po}}) \implies f_0(x^{\text{po}}) = f_0(y)$
  
- ▶ there are typically many Pareto optimal points
- ▶ for  $q = 2$ , set of Pareto optimal objective values is the **optimal trade-off curve**
- ▶ for  $q = 3$ , set of Pareto optimal objective values is the **optimal trade-off surface**

## Optimal and Pareto optimal points


set of achievable objective values  $\mathcal{O} = \{f_0(x) \mid x \text{ feasible}\}$

- ▶ feasible  $x$  is **optimal** if  $f_0(x)$  is the minimum value of  $\mathcal{O}$
- ▶ feasible  $x$  is **Pareto optimal** if  $f_0(x)$  is a minimal value of  $\mathcal{O}$



## Regularized least-squares



- ▶ minimize  $(\|Ax - b\|_2^2, \|x\|_2^2)$  (first objective is loss; second is regularization)
- ▶ example with  $A \in \mathbf{R}^{100 \times 10}$ ; heavy line shows Pareto optimal points



## Risk return trade-off in portfolio optimization

- ▶ variable  $x \in \mathbf{R}^n$  is investment portfolio, with  $x_i$  fraction invested in asset  $i$
- ▶  $\bar{p} \in \mathbf{R}^n$  is mean,  $\Sigma$  is covariance of asset returns
- ▶ portfolio return has mean  $\bar{p}^T x$ , variance  $x^T \Sigma x$
- ▶ minimize  $(-\bar{p}^T x, x^T \Sigma x)$ , subject to  $\mathbf{1}^T x = 1, x \geq 0$
- ▶ Pareto optimal portfolios trace out optimal risk-return curve

## Example




## Scalarization

- ▶ **scalarization** combines the multiple objectives into one (scalar) objective
- ▶ a standard method for finding Pareto optimal points
- ▶ choose  $\lambda > 0$  and solve scalar problem

$$\begin{array}{ll}\text{minimize} & \lambda^T f_0(x) = \lambda_1 F_1(x) + \cdots + \lambda_q F_q(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m, \quad h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$


- ▶  $\lambda_i$  are relative weights on the objectives
- ▶ if  $x$  is optimal for scalar problem, then it is Pareto-optimal for multicriterion problem
- ▶ for convex problems, can find (almost) all Pareto optimal points by varying  $\lambda > 0$

## Example



## Example: Regularized least-squares

- ▶ regularized least-squares problem: minimize  $(\|Ax - b\|_2^2, \|x\|_2^2)$
- ▶ take  $\lambda = (1, \gamma)$  with  $\gamma > 0$ , and minimize  $\|Ax - b\|_2^2 + \gamma \|x\|_2^2$





## Example: Risk-return trade-off

- ▶ risk-return trade-off: minimize  $(-\bar{p}^T x, x^T \Sigma x)$  subject to  $\mathbf{1}^T x = 1, x \geq 0$
- ▶ with  $\lambda = (1, \gamma)$  we obtain scalarized problem

$$\begin{array}{ll}\text{minimize} & -\bar{p}^T x + \gamma x^T \Sigma x \\ \text{subject to} & \mathbf{1}^T x = 1, \quad x \geq 0\end{array}$$

- ▶ objective is negative **risk-adjusted return**,  $\bar{p}^T x - \gamma x^T \Sigma x$
- ▶  $\gamma$  is called the **risk-aversion parameter**

## 5. Duality

# Outline

Lagrangian and dual function

Lagrange dual problem

KKT conditions

Sensitivity analysis

Problem reformulations

Theorems of alternatives

## Lagrangian

- ▶ **standard form problem** (not necessarily convex)

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

variable  $x \in \mathbf{R}^n$ , domain  $\mathcal{D}$ , optimal value  $p^\star$

- ▶ **Lagrangian:**  $L : \mathbf{R}^n \times \mathbf{R}^m \times \mathbf{R}^p \rightarrow \mathbf{R}$ , with  $\text{dom } L = \mathcal{D} \times \mathbf{R}^m \times \mathbf{R}^p$ ,

$$L(x, \lambda, \nu) = f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x)$$

- weighted sum of objective and constraint functions
- $\lambda_i$  is **Lagrange multiplier** associated with  $f_i(x) \leq 0$
- $\nu_i$  is Lagrange multiplier associated with  $h_i(x) = 0$

## Lagrange dual function

- ▶ **Lagrange dual function:**  $g : \mathbf{R}^m \times \mathbf{R}^p \rightarrow \mathbf{R}$ ,

$$g(\lambda, \nu) = \inf_{x \in \mathcal{D}} L(x, \lambda, \nu) = \inf_{x \in \mathcal{D}} \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x) \right)$$

- ▶  $g$  is concave, can be  $-\infty$  for some  $\lambda, \nu$
- ▶ **lower bound property:** if  $\lambda \geq 0$ , then  $g(\lambda, \nu) \leq p^\star$
- ▶ proof: if  $\tilde{x}$  is feasible and  $\lambda \geq 0$ , then

$$f_0(\tilde{x}) \geq L(\tilde{x}, \lambda, \nu) \geq \inf_{x \in \mathcal{D}} L(x, \lambda, \nu) = g(\lambda, \nu)$$

minimizing over all feasible  $\tilde{x}$  gives  $p^\star \geq g(\lambda, \nu)$

## Least-norm solution of linear equations

$$\begin{array}{ll}\text{minimize} & x^T x \\ \text{subject to} & Ax = b\end{array}$$

- ▶ Lagrangian is  $L(x, v) = x^T x + v^T (Ax - b)$
- ▶ to minimize  $L$  over  $x$ , set gradient equal to zero:

$$\nabla_x L(x, v) = 2x + A^T v = 0 \implies x = -(1/2)A^T v$$

- ▶ plug  $x$  into  $L$  to obtain

$$g(v) = L((-1/2)A^T v, v) = -\frac{1}{4}v^T A A^T v - b^T v$$

- ▶ lower bound property:  $p^\star \geq -(1/4)v^T A A^T v - b^T v$  for all  $v$

## Standard form LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b, \quad x \geq 0\end{array}$$

- ▶ Lagrangian is

$$L(x, \lambda, \nu) = c^T x + \nu^T (Ax - b) - \lambda^T x = -b^T \nu + (c + A^T \nu - \lambda)^T x$$

- ▶  $L$  is affine in  $x$ , so

$$g(\lambda, \nu) = \inf_x L(x, \lambda, \nu) = \begin{cases} -b^T \nu & A^T \nu - \lambda + c = 0 \\ -\infty & \text{otherwise} \end{cases}$$

- ▶  $g$  is linear on affine domain  $\{(\lambda, \nu) \mid A^T \nu - \lambda + c = 0\}$ , hence concave
- ▶ lower bound property:  $p^\star \geq -b^T \nu$  if  $A^T \nu + c \geq 0$

## Equality constrained norm minimization

$$\begin{array}{ll}\text{minimize} & \|x\| \\ \text{subject to} & Ax = b\end{array}$$

- dual function is

$$g(v) = \inf_x (\|x\| - v^T Ax + b^T v) = \begin{cases} b^T v & \|A^T v\|_* \leq 1 \\ -\infty & \text{otherwise} \end{cases}$$

where  $\|v\|_* = \sup_{\|u\| \leq 1} u^T v$  is dual norm of  $\|\cdot\|$

- lower bound property:  $p^* \geq b^T v$  if  $\|A^T v\|_* \leq 1$



## Two-way partitioning

$$\begin{array}{ll}\text{minimize} & x^T W x \\ \text{subject to} & x_i^2 = 1, \quad i = 1, \dots, n\end{array}$$

- ▶ a nonconvex problem; feasible set contains  $2^n$  discrete points
- ▶ interpretation: partition  $\{1, \dots, n\}$  in two sets encoded as  $x_i = 1$  and  $x_i = -1$
- ▶  $W_{ij}$  is cost of assigning  $i, j$  to the same set;  $-W_{ij}$  is cost of assigning to different sets
- ▶ dual function is

$$g(\nu) = \inf_x \left( x^T W x + \sum_i \nu_i (x_i^2 - 1) \right) = \inf_x x^T (W + \mathbf{diag}(\nu)) x - \mathbf{1}^T \nu = \begin{cases} -\mathbf{1}^T \nu & W + \mathbf{diag}(\nu) \geq 0 \\ -\infty & \text{otherwise} \end{cases}$$

- ▶ lower bound property:  $p^\star \geq -\mathbf{1}^T \nu$  if  $W + \mathbf{diag}(\nu) \geq 0$

## Lagrange dual and conjugate function

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & Ax \leq b, \quad Cx = d\end{array}$$

- dual function

$$\begin{aligned}g(\lambda, \nu) &= \inf_{x \in \text{dom } f_0} \left( f_0(x) + (A^T \lambda + C^T \nu)^T x - b^T \lambda - d^T \nu \right) \\ &= -f_0^*(-A^T \lambda - C^T \nu) - b^T \lambda - d^T \nu\end{aligned}$$

where  $f^*(y) = \sup_{x \in \text{dom } f} (y^T x - f(x))$  is conjugate of  $f_0$

- simplifies derivation of dual if conjugate of  $f_0$  is known
- **example: entropy maximization**

$$f_0(x) = \sum_{i=1}^n x_i \log x_i, \quad f_0^*(y) = \sum_{i=1}^n e^{y_i - 1}$$

# Outline

Lagrangian and dual function

Lagrange dual problem

KKT conditions

Sensitivity analysis

Problem reformulations

Theorems of alternatives

# The Lagrange dual problem

(Lagrange) **dual problem**

$$\begin{array}{ll}\text{maximize} & g(\lambda, \nu) \\ \text{subject to} & \lambda \geq 0\end{array}$$

- ▶ finds best lower bound on  $p^\star$ , obtained from Lagrange dual function
- ▶ a convex optimization problem, even if original **primal** problem is not
- ▶ dual optimal value denoted  $d^\star$
- ▶  $\lambda, \nu$  are dual feasible if  $\lambda \geq 0, (\lambda, \nu) \in \mathbf{dom} \, g$
- ▶ often simplified by making implicit constraint  $(\lambda, \nu) \in \mathbf{dom} \, g$  explicit

## Example: standard form LP

(see slide 5.5)

- ▶ primal standard form LP:

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- ▶ dual problem is

$$\begin{array}{ll}\text{maximize} & g(\lambda, \nu) \\ \text{subject to} & \lambda \geq 0\end{array}$$

with  $g(\lambda, \nu) = -b^T \nu$  if  $A^T \nu - \lambda + c = 0$ ,  $-\infty$  otherwise

- ▶ make implicit constraint explicit, and eliminate  $\lambda$  to obtain (transformed) dual problem

$$\begin{array}{ll}\text{maximize} & -b^T \nu \\ \text{subject to} & A^T \nu + c \geq 0\end{array}$$

## Weak and strong duality

**weak duality:**  $d^\star \leq p^\star$

- ▶ always holds (for convex and nonconvex problems)
- ▶ can be used to find nontrivial lower bounds for difficult problems, e.g., solving the SDP

$$\begin{array}{ll}\text{maximize} & -\mathbf{1}^T \boldsymbol{\nu} \\ \text{subject to} & W + \mathbf{diag}(\boldsymbol{\nu}) \succeq 0\end{array}$$

gives a lower bound for the two-way partitioning problem on page 5.7

**strong duality:**  $d^\star = p^\star$

- ▶ does not hold in general
- ▶ (usually) holds for convex problems
- ▶ conditions that guarantee strong duality in convex problems are called **constraint qualifications**

## Slater's constraint qualification

strong duality holds for a convex problem

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

if it is **strictly feasible**, *i.e.*, there is an  $x \in \mathbf{int} \mathcal{D}$  with  $f_i(x) < 0, i = 1, \dots, m, Ax = b$

- ▶ also guarantees that the dual optimum is attained (if  $p^\star > -\infty$ )
- ▶ can be sharpened: *e.g.*,
  - can replace  $\mathbf{int} \mathcal{D}$  with  $\mathbf{relint} \mathcal{D}$  (interior relative to affine hull)
  - affine inequalities do not need to hold with strict inequality
- ▶ there are many other types of constraint qualifications

## Inequality form LP

### primal problem

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

### dual function

$$g(\lambda) = \inf_x \left( (c + A^T \lambda)^T x - b^T \lambda \right) = \begin{cases} -b^T \lambda & A^T \lambda + c = 0 \\ -\infty & \text{otherwise} \end{cases}$$

### dual problem

$$\begin{array}{ll}\text{maximize} & -b^T \lambda \\ \text{subject to} & A^T \lambda + c = 0, \quad \lambda \geq 0\end{array}$$

- ▶ from the sharpened Slater's condition:  $p^\star = d^\star$  if the primal problem is feasible
- ▶ in fact,  $p^\star = d^\star$  except when primal and dual are both infeasible



## Quadratic program

**primal problem** (assume  $P \in \mathbf{S}_{++}^n$ )

$$\begin{array}{ll}\text{minimize} & x^T P x \\ \text{subject to} & A x \leq b\end{array}$$

**dual function**

$$g(\lambda) = \inf_x \left( x^T P x + \lambda^T (A x - b) \right) = -\frac{1}{4} \lambda^T A P^{-1} A^T \lambda - b^T \lambda$$



**dual problem**

$$\begin{array}{ll}\text{maximize} & -(1/4) \lambda^T A P^{-1} A^T \lambda - b^T \lambda \\ \text{subject to} & \lambda \geq 0\end{array}$$

- ▶ from the sharpened Slater's condition:  $p^\star = d^\star$  if the primal problem is feasible
- ▶ in fact,  $p^\star = d^\star$  always

## Geometric interpretation


- ▶ for simplicity, consider problem with one constraint  $f_1(x) \leq 0$
- ▶  $\mathcal{G} = \{(f_1(x), f_0(x)) \mid x \in \mathcal{D}\}$  is set of achievable (constraint, objective) values
- ▶ **interpretation of dual function:**  $g(\lambda) = \inf_{(u,t) \in \mathcal{G}} (t + \lambda u)$



- ▶  $\lambda u + t = g(\lambda)$  is (non-vertical) supporting hyperplane to  $\mathcal{G}$
- ▶ hyperplane intersects  $t$ -axis at  $t = g(\lambda)$

## Epigraph variation

- ▶ same with  $\mathcal{G}$  replaced with  $\mathcal{A} = \{(u, t) \mid f_1(x) \leq u, f_0(x) \leq t \text{ for some } x \in \mathcal{D}\}$



- ▶ strong duality holds if there is a non-vertical supporting hyperplane to  $\mathcal{A}$  at  $(0, p^*)$
- ▶ for convex problem,  $\mathcal{A}$  is convex, hence has supporting hyperplane at  $(0, p^*)$
- ▶ Slater's condition: if there exist  $(\tilde{u}, \tilde{t}) \in \mathcal{A}$  with  $\tilde{u} < 0$ , then supporting hyperplane at  $(0, p^*)$  must be non-vertical

# Outline

Lagrangian and dual function

Lagrange dual problem

**KKT conditions**

Sensitivity analysis

Problem reformulations

Theorems of alternatives

## Complementary slackness

- ▶ assume strong duality holds,  $x^\star$  is primal optimal,  $(\lambda^\star, \nu^\star)$  is dual optimal

$$\begin{aligned} f_0(x^\star) = g(\lambda^\star, \nu^\star) &= \inf_x \left( f_0(x) + \sum_{i=1}^m \lambda_i^\star f_i(x) + \sum_{i=1}^p \nu_i^\star h_i(x) \right) \\ &\leq f_0(x^\star) + \sum_{i=1}^m \lambda_i^\star f_i(x^\star) + \sum_{i=1}^p \nu_i^\star h_i(x^\star) \\ &\leq f_0(x^\star) \end{aligned}$$

- ▶ hence, the two inequalities hold with equality
- ▶  $x^\star$  minimizes  $L(x, \lambda^\star, \nu^\star)$
- ▶  $\lambda_i^\star f_i(x^\star) = 0$  for  $i = 1, \dots, m$  (known as **complementary slackness**):

$$\lambda_i^\star > 0 \implies f_i(x^\star) = 0, \quad f_i(x^\star) < 0 \implies \lambda_i^\star = 0$$

## Karush-Kuhn-Tucker (KKT) conditions

the **KKT conditions** (for a problem with differentiable  $f_i, h_i$ ) are

1. primal constraints:  $f_i(x) \leq 0, i = 1, \dots, m, h_i(x) = 0, i = 1, \dots, p$
2. dual constraints:  $\lambda \geq 0$
3. complementary slackness:  $\lambda_i f_i(x) = 0, i = 1, \dots, m$
4. gradient of Lagrangian with respect to  $x$  vanishes:

$$\nabla f_0(x) + \sum_{i=1}^m \lambda_i \nabla f_i(x) + \sum_{i=1}^p \nu_i \nabla h_i(x) = 0$$

if strong duality holds and  $x, \lambda, \nu$  are optimal, they satisfy the KKT conditions

## KKT conditions for convex problem

if  $\tilde{x}$ ,  $\tilde{\lambda}$ ,  $\tilde{v}$  satisfy KKT for a convex problem, then they are optimal:

- ▶ from complementary slackness:  $f_0(\tilde{x}) = L(\tilde{x}, \tilde{\lambda}, \tilde{v})$
- ▶ from 4th condition (and convexity):  $g(\tilde{\lambda}, \tilde{v}) = L(\tilde{x}, \tilde{\lambda}, \tilde{v})$

hence,  $f_0(\tilde{x}) = g(\tilde{\lambda}, \tilde{v})$

if Slater's condition is satisfied, then

*$x$  is optimal if and only if there exist  $\lambda$ ,  $v$  that satisfy KKT conditions*

- ▶ recall that Slater implies strong duality, and dual optimum is attained
- ▶ generalizes optimality condition  $\nabla f_0(x) = 0$  for unconstrained problem

# Outline

Lagrangian and dual function

Lagrange dual problem

KKT conditions

**Sensitivity analysis**

Problem reformulations

Theorems of alternatives



## Perturbation and sensitivity analysis

### (unperturbed) optimization problem and its dual

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & h_i(x) = 0, \quad i = 1, \dots, p\end{array}$$

$$\begin{array}{ll}\text{maximize} & g(\lambda, v) \\ \text{subject to} & \lambda \geq 0\end{array}$$

### perturbed problem and its dual

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq u_i, \quad i = 1, \dots, m \\ & h_i(x) = v_i, \quad i = 1, \dots, p\end{array}$$

$$\begin{array}{ll}\text{maximize} & g(\lambda, v) - u^T \lambda - v^T v \\ \text{subject to} & \lambda \geq 0\end{array}$$

- ▶  $x$  is primal variable;  $u, v$  are parameters
- ▶  $p^\star(u, v)$  is optimal value as a function of  $u, v$
- ▶  $p^\star(0, 0)$  is optimal value of unperturbed problem

## Global sensitivity via duality

- ▶ assume strong duality holds for unperturbed problem, with  $\lambda^\star$ ,  $v^\star$  dual optimal
- ▶ apply weak duality to perturbed problem:

$$p^\star(u, v) \geq g(\lambda^\star, v^\star) - u^T \lambda^\star - v^T v^\star = p^\star(0, 0) - u^T \lambda^\star - v^T v^\star$$

- ▶ **implications**

- if  $\lambda_i^\star$  large:  $p^\star$  increases greatly if we tighten constraint  $i$  ( $u_i < 0$ )
- if  $\lambda_i^\star$  small:  $p^\star$  does not decrease much if we loosen constraint  $i$  ( $u_i > 0$ )
- if  $v_i^\star$  large and positive:  $p^\star$  increases greatly if we take  $v_i < 0$
- if  $v_i^\star$  large and negative:  $p^\star$  increases greatly if we take  $v_i > 0$
- if  $v_i^\star$  small and positive:  $p^\star$  does not decrease much if we take  $v_i > 0$
- if  $v_i^\star$  small and negative:  $p^\star$  does not decrease much if we take  $v_i < 0$

## Local sensitivity via duality

if (in addition)  $p^*(u, v)$  is differentiable at  $(0, 0)$ , then


$$\lambda_i^* = -\frac{\partial p^*(0, 0)}{\partial u_i}, \quad v_i^* = -\frac{\partial p^*(0, 0)}{\partial v_i}$$

proof (for  $\lambda_i^*$ ): from global sensitivity result,

$$\frac{\partial p^*(0, 0)}{\partial u_i} = \lim_{t \searrow 0} \frac{p^*(te_i, 0) - p^*(0, 0)}{t} \geq -\lambda_i^* \quad \frac{\partial p^*(0, 0)}{\partial u_i} = \lim_{t \nearrow 0} \frac{p^*(te_i, 0) - p^*(0, 0)}{t} \leq -\lambda_i^*$$

hence, equality

$p^*(u)$  for a problem with one (inequality) constraint:



# Outline

Lagrangian and dual function

Lagrange dual problem

KKT conditions

Sensitivity analysis

**Problem reformulations**

Theorems of alternatives

## Duality and problem reformulations

- ▶ equivalent formulations of a problem can lead to very different duals
- ▶ reformulating primal problem can be useful when dual is difficult to derive, or uninteresting

### common reformulations

- ▶ introduce new variables and equality constraints
- ▶ make explicit constraints implicit or vice-versa
- ▶ transform objective or constraint functions, e.g., replace  $f_0(x)$  by  $\phi(f_0(x))$  with  $\phi$  convex, increasing

## Introducing new variables and equality constraints

- ▶ unconstrained problem: minimize  $f_0(Ax + b)$
- ▶ dual function is constant:  $g = \inf_x L(x) = \inf_x f_0(Ax + b) = p^\star$
- ▶ we have strong duality, but dual is quite useless
- ▶ introduce new variable  $y$  and equality constraints  $y = Ax + b$

$$\begin{array}{ll}\text{minimize} & f_0(y) \\ \text{subject to} & Ax + b - y = 0\end{array}$$

- ▶ dual of reformulated problem is

$$\begin{array}{ll}\text{maximize} & b^T \nu - f_0^*(\nu) \\ \text{subject to} & A^T \nu = 0\end{array}$$

- ▶ a nontrivial, useful dual (assuming the conjugate  $f_0^*$  is easy to express)

## Example: Norm approximation

- ▶ minimize  $\|Ax - b\|$
- ▶ reformulate as minimize  $\|y\|$  subject to  $y = Ax - b$
- ▶ recall conjugate of general norm:

$$\|z\|^* = \begin{cases} 0 & \|z\|_* \leq 1 \\ \infty & \text{otherwise} \end{cases}$$

- ▶ dual of (reformulated) norm approximation problem:

$$\begin{array}{ll} \text{maximize} & b^T v \\ \text{subject to} & A^T v = 0, \quad \|v\|_* \leq 1 \end{array}$$

# Outline

Lagrangian and dual function

Lagrange dual problem

KKT conditions

Sensitivity analysis

Problem reformulations

**Theorems of alternatives**



## Theorems of alternatives

- ▶ consider two systems of inequality and equality constraints
- ▶ called **weak alternatives** if no more than one system is feasible
- ▶ called **strong alternatives** if exactly one of them is feasible
- ▶ examples: for any  $a \in \mathbf{R}$ , with variable  $x \in \mathbf{R}$ ,
  - $x > a$  and  $x \leq a - 1$  are weak alternatives
  - $x > a$  and  $x \leq a$  are strong alternatives
- ▶ a **theorem of alternatives** states that two inequality systems are (weak or strong) alternatives
- ▶ can be considered the extension of duality to feasibility problems

## Feasibility problems

- ▶ consider system of (not necessarily convex) inequalities and equalities

$$f_i(x) \leq 0, \quad i = 1, \dots, m, \quad h_i(x) = 0, \quad i = 1, \dots, p$$

- ▶ express as **feasibility problem**

$$\begin{array}{ll} \text{minimize} & 0 \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m, \\ & h_i(x) = 0, \quad i = 1, \dots, p \end{array}$$

- ▶ if system is feasible,  $p^\star = 0$ ; if not,  $p^\star = \infty$

## Duality for feasibility problems

- ▶ dual function of feasibility problem is  $g(\lambda, \nu) = \inf_x \left( \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x) \right)$
- ▶ for  $\lambda \geq 0$ , we have  $g(\lambda, \nu) \leq p^\star$
- ▶ it follows that feasibility of the inequality system

$$\lambda \geq 0, \quad g(\lambda, \nu) > 0$$

implies the original system is infeasible

- ▶ so this is a weak alternative to original system
- ▶ it is strong if  $f_i$  convex,  $h_i$  affine, and a constraint qualification holds
- ▶  $g$  is positive homogeneous so we can write alternative system as

$$\lambda \geq 0, \quad g(\lambda, \nu) \geq 1$$

## Example: Nonnegative solution of linear equations

- ▶ consider system

$$Ax = b, \quad x \geq 0$$

- ▶ dual function is  $g(\lambda, \nu) = \begin{cases} -b^T \nu & A^T \nu = \lambda \\ -\infty & \text{otherwise} \end{cases}$

- ▶ can express strong alternative of  $Ax = b, x \geq 0$  as

$$A^T \nu \geq 0, \quad b^T \nu \leq -1$$

(we can replace  $b^T \nu \leq -1$  with  $b^T \nu = -1$ )

## Farkas' lemma

- Farkas' lemma:

$$Ax \leq 0, \quad c^T x < 0 \quad \text{and} \quad A^T y + c = 0, \quad y \geq 0$$

are strong alternatives

- proof: use (strong) duality for (feasible) LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq 0 \end{array}$$

## Investment arbitrage

- ▶ we invest  $x_j$  in each of  $n$  assets  $1, \dots, n$  with prices  $p_1, \dots, p_n$
- ▶ our initial cost is  $p^T x$
- ▶ at the end of the investment period there are only  $m$  possible outcomes  $i = 1, \dots, m$
- ▶  $V_{ij}$  is the **payoff** or final value of asset  $j$  in outcome  $i$
- ▶ first investment is risk-free (cash):  $p_1 = 1$  and  $V_{i1} = 1$  for all  $i$
  
- ▶ **arbitrage** means there is  $x$  with  $p^T x < 0$ ,  $Vx \geq 0$
- ▶ arbitrage means we receive money up front, and our investment cannot lose
- ▶ standard assumption in economics: the prices are such that **there is no arbitrage**

## Absence of arbitrage

- ▶ by Farkas' lemma, there is no arbitrage  $\iff$  there exists  $y \in \mathbf{R}_+^m$  with  $V^T y = p$
- ▶ since first column of  $V$  is  $\mathbf{1}$ , we have  $\mathbf{1}^T y = 1$
- ▶  $y$  is interpreted as a **risk-neutral probability** on the outcomes  $1, \dots, m$
- ▶  $V^T y$  are the expected values of the payoffs under the risk-neutral probability
- ▶ interpretation of  $V^T y = p$ :  
*asset prices equal their expected payoff under the risk-neutral probability*

- ▶ **arbitrage theorem**: there is no arbitrage  $\iff$  there exists a risk-neutral probability distribution under which each asset price is its expected payoff

## Example

$$V = \begin{bmatrix} 1.0 & 0.5 & 0.0 \\ 1.0 & 0.8 & 0.0 \\ 1.0 & 1.0 & 1.0 \\ 1.0 & 1.3 & 4.0 \end{bmatrix}, \quad p = \begin{bmatrix} 1.0 \\ 0.9 \\ 0.3 \end{bmatrix}, \quad \tilde{p} = \begin{bmatrix} 1.0 \\ 0.8 \\ 0.7 \end{bmatrix}$$

- ▶ with prices  $p$ , there is an arbitrage

$$x = \begin{bmatrix} 6.2 \\ -7.7 \\ 1.5 \end{bmatrix}, \quad p^T x = -0.2, \quad \mathbf{1}^T x = 0, \quad Vx = \begin{bmatrix} 2.35 \\ 0.04 \\ 0.00 \\ 2.19 \end{bmatrix}$$

- ▶ with prices  $\tilde{p}$ , there is no arbitrage, with risk-neutral probability

$$y = \begin{bmatrix} 0.36 \\ 0.27 \\ 0.26 \\ 0.11 \end{bmatrix}, \quad V^T y = \begin{bmatrix} 1.0 \\ 0.8 \\ 0.7 \end{bmatrix}$$



## 6. Approximation and fitting

# Outline

Norm and penalty approximation

Regularized approximation

Robust approximation

## Norm approximation

- ▶ minimize  $\|Ax - b\|$ , with  $A \in \mathbf{R}^{m \times n}$ ,  $m \geq n$ ,  $\|\cdot\|$  is any norm
- ▶ **approximation**:  $Ax^\star$  is the best approximation of  $b$  by a linear combination of columns of  $A$
- ▶ **geometric**:  $Ax^\star$  is point in  $\mathcal{R}(A)$  closest to  $b$  (in norm  $\|\cdot\|$ )
- ▶ **estimation**: linear measurement model  $y = Ax + v$ 
  - measurement  $y$ ,  $v$  is measurement error,  $x$  is to be estimated
  - implausibility of  $v$  is  $\|v\|$
  - given  $y = b$ , most plausible  $x$  is  $x^\star$
- ▶ **optimal design**:  $x$  are design variables (input),  $Ax$  is result (output)
  - $x^\star$  is design that best approximates desired result  $b$  (in norm  $\|\cdot\|$ )

## Examples

► Euclidean approximation ( $\|\cdot\|_2$ )

- solution  $x^\star = A^\dagger b$

► Chebyshev or minimax approximation ( $\|\cdot\|_\infty$ )

- can be solved via LP

$$\begin{array}{ll}\text{minimize} & t \\ \text{subject to} & -t\mathbf{1} \leq Ax - b \leq t\mathbf{1}\end{array}$$

► sum of absolute residuals approximation ( $\|\cdot\|_1$ )

- can be solved via LP

$$\begin{array}{ll}\text{minimize} & \mathbf{1}^T y \\ \text{subject to} & -y \leq Ax - b \leq y\end{array}$$

## Penalty function approximation

$$\begin{array}{ll}\text{minimize} & \phi(r_1) + \cdots + \phi(r_m) \\ \text{subject to} & r = Ax - b\end{array}$$

( $A \in \mathbf{R}^{m \times n}$ ,  $\phi : \mathbf{R} \rightarrow \mathbf{R}$  is a convex penalty function)


### examples

- ▶ quadratic:  $\phi(u) = u^2$
- ▶ deadzone-linear with width  $a$ :

$$\phi(u) = \max\{0, |u| - a\}$$

- ▶ log-barrier with limit  $a$ :

$$\phi(u) = \begin{cases} -a^2 \log(1 - (u/a)^2) & |u| < a \\ \infty & \text{otherwise} \end{cases}$$



## Example: histograms of residuals


$A \in \mathbf{R}^{100 \times 30}$ ; shape of penalty function affects distribution of residuals

absolute value  $\phi(u) = |u|$

square  $\phi(u) = u^2$


deadzone  $\phi(u) = \max\{0, |u| - 0.5\}$

log-barrier  $\phi(u) = -\log(1 - u^2)$




## Huber penalty function

$$\phi_{\text{hub}}(u) = \begin{cases} u^2 & |u| \leq M \\ M(2|u| - M) & |u| > M \end{cases}$$



- ▶ linear growth for large  $u$  makes approximation less sensitive to outliers
- ▶ called a **robust penalty**

## Example



- ▶ 42 points (circles)  $t_i, y_i$ , with two outliers
- ▶ affine function  $f(t) = \alpha + \beta t$  fit using quadratic (dashed) and Huber (solid) penalty



## Least-norm problems

- ▶ least-norm problem:

$$\begin{array}{ll}\text{minimize} & \|x\| \\ \text{subject to} & Ax = b,\end{array}$$

with  $A \in \mathbf{R}^{m \times n}$ ,  $m \leq n$ ,  $\|\cdot\|$  is any norm

- ▶ **geometric:**  $x^\star$  is smallest point in solution set  $\{x \mid Ax = b\}$
- ▶ **estimation:**
  - $b = Ax$  are (perfect) measurements of  $x$
  - $\|x\|$  is implausibility of  $x$
  - $x^\star$  is most plausible estimate consistent with measurements
- ▶ **design:**  $x$  are design variables (inputs);  $b$  are required results (outputs)
  - $x^\star$  is smallest ('most efficient') design that satisfies requirements

## Examples

- ▶ least Euclidean norm ( $\|\cdot\|_2$ )
  - solution  $x = A^\dagger b$  (assuming  $b \in \mathcal{R}(A)$ )

- ▶ least sum of absolute values ( $\|\cdot\|_1$ )

- can be solved via LP

$$\begin{array}{ll}\text{minimize} & \mathbf{1}^T y \\ \text{subject to} & -y \leq x \leq y, \quad Ax = b\end{array}$$

- tends to yield sparse  $x^\star$

# Outline

Norm and penalty approximation

Regularized approximation

Robust approximation

## Regularized approximation

- ▶ a bi-objective problem:

$$\text{minimize (w.r.t. } \mathbf{R}_+^2) \quad (\|Ax - b\|, \|x\|)$$

- ▶  $A \in \mathbf{R}^{m \times n}$ , norms on  $\mathbf{R}^m$  and  $\mathbf{R}^n$  can be different
- ▶ interpretation: find good approximation  $Ax \approx b$  with small  $x$
- ▶ **estimation:** linear measurement model  $y = Ax + v$ , with prior knowledge that  $\|x\|$  is small
- ▶ **optimal design:** small  $x$  is cheaper or more efficient, or the linear model  $y = Ax$  is only valid for small  $x$
- ▶ **robust approximation:** good approximation  $Ax \approx b$  with small  $x$  is less sensitive to errors in  $A$  than good approximation with large  $x$

## Scalarized problem

- ▶ minimize  $\|Ax - b\| + \gamma\|x\|$
- ▶ solution for  $\gamma > 0$  traces out optimal trade-off curve
- ▶ other common method: minimize  $\|Ax - b\|^2 + \delta\|x\|^2$  with  $\delta > 0$
- ▶ with  $\|\cdot\|_2$ , called **Tikhonov regularization** or **ridge regression**

$$\text{minimize} \quad \|Ax - b\|_2^2 + \delta\|x\|_2^2$$

- ▶ can be solved as a least-squares problem

$$\text{minimize} \quad \left\| \begin{bmatrix} A \\ \sqrt{\delta}I \end{bmatrix} x - \begin{bmatrix} b \\ 0 \end{bmatrix} \right\|_2^2$$

with solution  $x^\star = (A^T A + \delta I)^{-1} A^T b$

## Optimal input design

- ▶ **linear dynamical system** (or **convolution system**) with impulse response  $h$ :

$$y(t) = \sum_{\tau=0}^t h(\tau)u(t-\tau), \quad t = 0, 1, \dots, N$$

- ▶ **input design problem:** multicriterion problem with 3 objectives







- tracking error with desired output  $y_{\text{des}}$ :  $J_{\text{track}} = \sum_{t=0}^N (y(t) - y_{\text{des}}(t))^2$
- input variation:  $J_{\text{der}} = \sum_{t=0}^{N-1} (u(t+1) - u(t))^2$
- input magnitude:  $J_{\text{mag}} = \sum_{t=0}^N u(t)^2$

track desired output using a small and slowly varying input signal

- ▶ **regularized least-squares formulation:** minimize  $J_{\text{track}} + \delta J_{\text{der}} + \eta J_{\text{mag}}$ 
  - for fixed  $\delta, \eta$ , a least-squares problem in  $u(0), \dots, u(N)$

## Example

- ▶ minimize  $J_{\text{track}} + \delta J_{\text{der}} + \eta J_{\text{mag}}$
- ▶ (top)  $\delta = 0$ , small  $\eta$ ; (middle)  $\delta = 0$ , larger  $\eta$ ; (bottom) large  $\delta$



## Signal reconstruction

- bi-objective problem:

$$\text{minimize (w.r.t. } \mathbf{R}_+^2) \quad (\|\hat{x} - x_{\text{cor}}\|_2, \phi(\hat{x}))$$


- $x \in \mathbf{R}^n$  is unknown signal
- $x_{\text{cor}} = x + v$  is (known) corrupted version of  $x$ , with additive noise  $v$
- variable  $\hat{x}$  (reconstructed signal) is estimate of  $x$
- $\phi : \mathbf{R}^n \rightarrow \mathbf{R}$  is regularization function or smoothing objective

- **examples:**


- quadratic smoothing,  $\phi_{\text{quad}}(\hat{x}) = \sum_{i=1}^{n-1} (\hat{x}_{i+1} - \hat{x}_i)^2$
- total variation smoothing,  $\phi_{\text{tv}}(\hat{x}) = \sum_{i=1}^{n-1} |\hat{x}_{i+1} - \hat{x}_i|$



## Quadratic smoothing example




original signal  $x$  and noisy signal  $x_{\text{cor}}$




three solutions on trade-off curve  
 $\|\hat{x} - x_{\text{cor}}\|_2$  versus  $\phi_{\text{quad}}(\hat{x})$

## Reconstructing a signal with sharp transitions



original signal  $x$  and noisy signal  $x_{\text{cor}}$




three solutions on trade-off curve


$\|\hat{x} - x_{\text{cor}}\|_2$  versus  $\phi_{\text{quad}}(\hat{x})$

- quadratic smoothing smooths out noise **and** sharp transitions in signal

## Total variation reconstruction



original signal  $x$  and noisy signal  $x_{\text{cor}}$



three solutions on trade-off curve

$$\|\hat{x} - x_{\text{cor}}\|_2 \text{ versus } \phi_{\text{tv}}(\hat{x})$$

- ▶ total variation smoothing preserves sharp transitions in signal

# Outline

Norm and penalty approximation

Regularized approximation

Robust approximation

## Robust approximation


- ▶ minimize  $\|Ax - b\|$  with uncertain  $A$
- ▶ two approaches:
  - **stochastic**: assume  $A$  is random, minimize  $\mathbf{E} \|Ax - b\|$
  - **worst-case**: set  $\mathcal{A}$  of possible values of  $A$ , minimize  $\sup_{A \in \mathcal{A}} \|Ax - b\|$
- ▶ tractable only in special cases (certain norms  $\|\cdot\|$ , distributions, sets  $\mathcal{A}$ )

## Example

$$A(u) = A_0 + uA_1, u \in [-1, 1]$$

- ▶  $x_{\text{nom}}$  minimizes  $\|A_0x - b\|_2^2$
- ▶  $x_{\text{stoch}}$  minimizes  $\mathbf{E} \|A(u)x - b\|_2^2$   
with  $u$  uniform on  $[-1, 1]$
- ▶  $x_{\text{wc}}$  minimizes  $\sup_{-1 \leq u \leq 1} \|A(u)x - b\|_2^2$

plot shows  $r(u) = \|A(u)x - b\|_2$  versus  $u$



## Stochastic robust least-squares

- ▶  $A = \bar{A} + U$ ,  $U$  random,  $\mathbf{E} U = 0$ ,  $\mathbf{E} U^T U = P$
- ▶ stochastic least-squares problem: minimize  $\mathbf{E} \|(\bar{A} + U)x - b\|_2^2$
- ▶ explicit expression for objective:

$$\begin{aligned}\mathbf{E} \|Ax - b\|_2^2 &= \mathbf{E} \|\bar{A}x - b + Ux\|_2^2 \\ &= \|\bar{A}x - b\|_2^2 + \mathbf{E} x^T U^T U x \\ &= \|\bar{A}x - b\|_2^2 + x^T P x\end{aligned}$$

- ▶ hence, robust least-squares problem is equivalent to: minimize  $\|\bar{A}x - b\|_2^2 + \|P^{1/2}x\|_2^2$
- ▶ for  $P = \delta I$ , get Tikhonov regularized problem: minimize  $\|\bar{A}x - b\|_2^2 + \delta \|x\|_2^2$

## Worst-case robust least-squares

- ▶  $\mathcal{A} = \{\bar{A} + u_1 A_1 + \cdots + u_p A_p \mid \|u\|_2 \leq 1\}$  (an ellipsoid in  $\mathbf{R}^{m \times n}$ )
- ▶ worst-case robust least-squares problem is

$$\text{minimize} \quad \sup_{A \in \mathcal{A}} \|Ax - b\|_2^2 = \sup_{\|u\|_2 \leq 1} \|P(x)u + q(x)\|_2^2$$

where  $P(x) = \begin{bmatrix} A_1 x & A_2 x & \cdots & A_p x \end{bmatrix}$ ,  $q(x) = \bar{A}x - b$

- ▶ from book appendix B, strong duality holds between the following problems

$$\begin{array}{ll} \text{maximize} & \|Pu + q\|_2^2 \\ \text{subject to} & \|u\|_2^2 \leq 1 \end{array} \qquad \begin{array}{ll} \text{minimize} & t + \lambda \\ \text{subject to} & \begin{bmatrix} I & P & q \\ P^T & \lambda I & 0 \\ q^T & 0 & t \end{bmatrix} \succeq 0 \end{array}$$


- ▶ hence, robust least-squares problem is equivalent to SDP

$$\begin{array}{ll} \text{minimize} & t + \lambda \\ \text{subject to} & \begin{bmatrix} I & P(x) & q(x) \\ P(x)^T & \lambda I & 0 \\ q(x)^T & 0 & t \end{bmatrix} \succeq 0 \end{array}$$



## Example

- ▶  $r(u) = \|(A_0 + u_1 A_1 + u_2 A_2)x - b\|_2$ ,  $u$  uniform on unit disk
- ▶ three choices of  $x$ :
  - $x_{ls}$  minimizes  $\|A_0 x - b\|_2$
  - $x_{tik}$  minimizes  $\|A_0 x - b\|_2^2 + \delta \|x\|_2^2$  (Tikhonov solution)
  - $x_{rls}$  minimizes  $\sup_{A \in \mathcal{A}} \|Ax - b\|_2^2 + \|x\|_2^2$



## 7. Statistical estimation

# Outline

Maximum likelihood estimation

Hypothesis testing

Experiment design

## Maximum likelihood estimation

- ▶ **parametric distribution estimation:** choose from a family of densities  $p_x(y)$ , indexed by a parameter  $x$  (often denoted  $\theta$ )
- ▶ we take  $p_x(y) = 0$  for invalid values of  $x$
- ▶  $p_x(y)$ , as a function of  $x$ , is called **likelihood function**
- ▶  $l(x) = \log p_x(y)$ , as a function of  $x$ , is called **log-likelihood function**
  
- ▶ **maximum likelihood estimation (MLE):** choose  $x$  to maximize  $p_x(y)$  (or  $l(x)$ )
- ▶ a convex optimization problem if  $\log p_x(y)$  is concave in  $x$  for fixed  $y$
- ▶ not the same as  $\log p_x(y)$  concave in  $y$  for fixed  $x$ , *i.e.*,  $p_x(y)$  is a family of log-concave densities

## Linear measurements with IID noise

### linear measurement model

$$y_i = a_i^T x + v_i, \quad i = 1, \dots, m$$

- ▶  $x \in \mathbf{R}^n$  is vector of unknown parameters
- ▶  $v_i$  is IID measurement noise, with density  $p(z)$
- ▶  $y_i$  is measurement:  $y \in \mathbf{R}^m$  has density  $p_x(y) = \prod_{i=1}^m p(y_i - a_i^T x)$

**maximum likelihood estimate:** any solution  $x$  of

$$\text{maximize} \quad l(x) = \sum_{i=1}^m \log p(y_i - a_i^T x)$$

( $y$  is observed value)

## Examples

- ▶ Gaussian noise  $\mathcal{N}(0, \sigma^2)$ :  $p(z) = (2\pi\sigma^2)^{-1/2} e^{-z^2/(2\sigma^2)}$ ,

$$l(x) = -\frac{m}{2} \log(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^m (a_i^T x - y_i)^2$$

ML estimate is least-squares solution

- ▶ Laplacian noise:  $p(z) = (1/(2a)) e^{-|z|/a}$ ,

$$l(x) = -m \log(2a) - \frac{1}{a} \sum_{i=1}^m |a_i^T x - y_i|$$

ML estimate is  $\ell_1$ -norm solution

- ▶ uniform noise on  $[-a, a]$ :

$$l(x) = \begin{cases} -m \log(2a) & |a_i^T x - y_i| \leq a, \quad i = 1, \dots, m \\ -\infty & \text{otherwise} \end{cases}$$

ML estimate is any  $x$  with  $|a_i^T x - y_i| \leq a$

## Logistic regression

- ▶ random variable  $y \in \{0, 1\}$  with distribution


$$p = \mathbf{prob}(y = 1) = \frac{\exp(a^T u + b)}{1 + \exp(a^T u + b)}$$

- ▶  $a, b$  are parameters;  $u \in \mathbf{R}^n$  are (observable) explanatory variables
- ▶ estimation problem: estimate  $a, b$  from  $m$  observations  $(u_i, y_i)$
- ▶ log-likelihood function (for  $y_1 = \dots = y_k = 1, y_{k+1} = \dots = y_m = 0$ ):

$$\begin{aligned} l(a, b) &= \log \left( \prod_{i=1}^k \frac{\exp(a^T u_i + b)}{1 + \exp(a^T u_i + b)} \prod_{i=k+1}^m \frac{1}{1 + \exp(a^T u_i + b)} \right) \\ &= \sum_{i=1}^k (a^T u_i + b) - \sum_{i=1}^m \log(1 + \exp(a^T u_i + b)) \end{aligned}$$

concave in  $a, b$

## Example



- ▶  $n = 1, m = 50$  measurements; circles show points  $(u_i, y_i)$
- ▶ solid curve is ML estimate of  $p = \exp(au + b)/(1 + \exp(au + b))$



## Gaussian covariance estimation

- ▶ fit Gaussian distribution  $\mathcal{N}(0, \Sigma)$  to observed data  $y_1, \dots, y_N$
- ▶ log-likelihood is

$$\begin{aligned}l(\Sigma) &= \frac{1}{2} \sum_{k=1}^N \left( -2\pi n - \log \det \Sigma - y^T \Sigma^{-1} y \right) \\ &= \frac{N}{2} \left( -2\pi n - \log \det \Sigma - \mathbf{tr} \Sigma^{-1} Y \right)\end{aligned}$$

with  $Y = (1/N) \sum_{k=1}^N y_k y_k^T$ , the empirical covariance

- ▶  $l$  is **not** concave in  $\Sigma$  (the  $\log \det \Sigma$  term has the wrong sign)
- ▶ with no constraints or regularization, MLE is empirical covariance  $\Sigma^{\text{ml}} = Y$

## Change of variables

- ▶ change variables to  $S = \Sigma^{-1}$
- ▶ recover original parameter via  $\Sigma = S^{-1}$
- ▶  $S$  is the **natural parameter** in an **exponential family** description of a Gaussian
- ▶ in terms of  $S$ , log-likelihood is

$$l(S) = \frac{N}{2} (-2\pi n + \log \det S - \mathbf{tr} SY)$$

which is **concave**

- ▶ (a similar trick can be used to handle nonzero mean)

## Fitting a sparse inverse covariance

- ▶  $S$  is the **precision matrix** of the Gaussian
- ▶  $S_{ij} = 0$  means that  $y_i$  and  $y_j$  are independent, conditioned on  $y_k, k \neq i, j$
- ▶ sparse  $S$  means
  - many pairs of components are conditionally independent, given the others
  - $y$  is described by a sparse (Gaussian) Bayes network
- ▶ to fit data with  $S$  sparse, minimize convex function

$$-\log \det S + \mathbf{tr} SY + \lambda \sum_{i \neq j} |S_{ij}|$$

over  $S \in \mathbf{S}^n$ , with hyper-parameter  $\lambda \geq 0$

## Example

- ▶ example with  $n = 4$ ,  $N = 10$  samples generated from a sparse  $S^{\text{true}}$

$$S^{\text{true}} = \begin{bmatrix} 1 & 0 & 0.5 & 0 \\ 0 & 1 & 0 & 0.1 \\ 0.5 & 0 & 1 & 0.3 \\ 0 & 0.1 & 0.3 & 1 \end{bmatrix}$$

- ▶ empirical and sparse estimate values of  $\Sigma^{-1}$  (with  $\lambda = 0.2$ )

$$Y^{-1} = \begin{bmatrix} 3 & 0.8 & 3.3 & 1.2 \\ 0.8 & 1.2 & 1.2 & 0.9 \\ 3.2 & 1.2 & 4.6 & 2.1 \\ 1.2 & 0.9 & 2.1 & 2.7 \end{bmatrix}, \quad \hat{S} = \begin{bmatrix} 0.9 & 0 & 0.6 & 0 \\ 0 & 0.7 & 0 & 0.1 \\ 0.6 & 0 & 1.1 & 0.2 \\ 0 & 0.1 & 0.2 & 1.2 \end{bmatrix}.$$

- ▶ estimation errors:  $\|S^{\text{true}} - Y^{-1}\|_F^2 = 49.8, \quad \|S^{\text{true}} - \hat{S}\|_F^2 = 0.2$

# Outline

Maximum likelihood estimation

Hypothesis testing

Experiment design

## (Binary) hypothesis testing

### detection (hypothesis testing) problem

given observation of a random variable  $X \in \{1, \dots, n\}$ , choose between:

- ▶ hypothesis 1:  $X$  was generated by distribution  $p = (p_1, \dots, p_n)$
- ▶ hypothesis 2:  $X$  was generated by distribution  $q = (q_1, \dots, q_n)$

### randomized detector

- ▶ a nonnegative matrix  $T \in \mathbf{R}^{2 \times n}$ , with  $\mathbf{1}^T T = \mathbf{1}^T$
- ▶ if we observe  $X = k$ , we choose hypothesis 1 with probability  $t_{1k}$ , hypothesis 2 with probability  $t_{2k}$
- ▶ if all elements of  $T$  are 0 or 1, it is called a **deterministic detector**

## Detection probability matrix

$$D = \begin{bmatrix} Tp & Tq \end{bmatrix} = \begin{bmatrix} 1 - P_{\text{fp}} & P_{\text{fn}} \\ P_{\text{fp}} & 1 - P_{\text{fn}} \end{bmatrix}$$

- ▶  $P_{\text{fp}}$  is probability of selecting hypothesis 2 if  $X$  is generated by distribution 1 (false positive)
- ▶  $P_{\text{fn}}$  is probability of selecting hypothesis 1 if  $X$  is generated by distribution 2 (false negative)
- ▶ **multi-objective formulation of detector design**

$$\begin{array}{ll} \text{minimize (w.r.t. } \mathbf{R}_+^2) & (P_{\text{fp}}, P_{\text{fn}}) = ((Tp)_2, (Tq)_1) \\ \text{subject to} & t_{1k} + t_{2k} = 1, \quad k = 1, \dots, n \\ & t_{ik} \geq 0, \quad i = 1, 2, \quad k = 1, \dots, n \end{array}$$

variable  $T \in \mathbf{R}^{2 \times n}$

## Scalarization

- ▶ scalarize with weight  $\lambda > 0$  to obtain

$$\begin{array}{ll}\text{minimize} & (Tp)_2 + \lambda(Tq)_1 \\ \text{subject to} & t_{1k} + t_{2k} = 1, \quad t_{ik} \geq 0, \quad i = 1, 2, \quad k = 1, \dots, n\end{array}$$

- ▶ an LP with a simple analytical solution

$$(t_{1k}, t_{2k}) = \begin{cases} (1, 0) & p_k \geq \lambda q_k \\ (0, 1) & p_k < \lambda q_k \end{cases}$$

- ▶ a deterministic detector, given by a **likelihood ratio test**
- ▶ if  $p_k = \lambda q_k$  for some  $k$ , any value  $0 \leq t_{1k} \leq 1$ ,  $t_{1k} = 1 - t_{2k}$  is optimal (i.e., Pareto-optimal detectors include non-deterministic detectors)



## Minimax detector


- ▶ minimize maximum of false positive and false negative probabilities

$$\begin{array}{ll}\text{minimize} & \max\{P_{\text{fp}}, P_{\text{fn}}\} = \max\{(Tp)_2, (Tq)_1\} \\ \text{subject to} & t_{1k} + t_{2k} = 1, \quad t_{ik} \geq 0, \quad i = 1, 2, \quad k = 1, \dots, n\end{array}$$

- ▶ an LP; solution is usually not deterministic

## Example

$$\begin{bmatrix} p & q \end{bmatrix} = \begin{bmatrix} 0.70 & 0.10 \\ 0.20 & 0.10 \\ 0.05 & 0.70 \\ 0.05 & 0.10 \end{bmatrix}$$



solutions 1, 2, 3 (and endpoints) are deterministic; 4 is minimax detector

# Outline

Maximum likelihood estimation

Hypothesis testing

Experiment design

## Experiment design

- ▶  $m$  linear measurements  $y_i = a_i^T x + w_i$ ,  $i = 1, \dots, m$  of unknown  $x \in \mathbf{R}^n$
- ▶ measurement errors  $w_i$  are IID  $\mathcal{N}(0, 1)$
- ▶ ML (least-squares) estimate is

$$\hat{x} = \left( \sum_{i=1}^m a_i a_i^T \right)^{-1} \sum_{i=1}^m y_i a_i$$

- ▶ error  $e = \hat{x} - x$  has zero mean and covariance

$$E = \mathbf{E} e e^T = \left( \sum_{i=1}^m a_i a_i^T \right)^{-1}$$

- ▶ confidence ellipsoids are given by  $\{x \mid (x - \hat{x})^T E^{-1} (x - \hat{x}) \leq \beta\}$
- ▶ **experiment design**: choose  $a_i \in \{v_1, \dots, v_p\}$  (set of possible test vectors) to make  $E$  ‘small’

## Vector optimization formulation

- formulate as vector optimization problem

$$\begin{array}{ll} \text{minimize (w.r.t. } \mathbf{S}_+^n) & E = \left( \sum_{k=1}^p m_k v_k v_k^T \right)^{-1} \\ \text{subject to} & m_k \geq 0, \quad m_1 + \cdots + m_p = m \\ & m_k \in \mathbf{Z} \end{array}$$

- variables are  $m_k$ , the number of vectors  $a_i$  equal to  $v_k$
- difficult in general, due to integer constraint
- common scalarizations: minimize  $\log \det E$ ,  $\mathbf{tr} E$ ,  $\lambda_{\max}(E)$ ,  $\dots$

## Relaxed experiment design

- ▶ assume  $m \gg p$ , use  $\lambda_k = m_k/m$  as (continuous) real variable

$$\begin{array}{ll} \text{minimize (w.r.t. } \mathbf{S}_+^n) & E = (1/m) \left( \sum_{k=1}^p \lambda_k v_k v_k^T \right)^{-1} \\ \text{subject to} & \lambda \geq 0, \quad \mathbf{1}^T \lambda = 1 \end{array}$$

- ▶ a convex relaxation, since we ignore constraint that  $m\lambda_k \in \mathbf{Z}$
- ▶ optimal value is lower bound on optimal value of (integer) experiment design problem
- ▶ simple rounding of  $\lambda_k m$  gives heuristic for experiment design problem

## *D*-optimal design

- scalarize via log determinant

$$\begin{array}{ll}\text{minimize} & \log \det \left( \sum_{k=1}^p \lambda_k v_k v_k^T \right)^{-1} \\ \text{subject to} & \lambda \geq 0, \quad \mathbf{1}^T \lambda = 1\end{array}$$

- interpretation: minimizes volume of confidence ellipsoids

## Dual of $D$ -optimal experiment design problem

### dual problem

$$\begin{array}{ll}\text{maximize} & \log \det W + n \log n \\ \text{subject to} & v_k^T W v_k \leq 1, \quad k = 1, \dots, p\end{array}$$

interpretation:  $\{x \mid x^T W x \leq 1\}$  is minimum volume ellipsoid centered at origin, that includes all test vectors  $v_k$

**complementary slackness:** for  $\lambda$ ,  $W$  primal and dual optimal


$$\lambda_k (1 - v_k^T W v_k) = 0, \quad k = 1, \dots, p$$

optimal experiment uses vectors  $v_k$  on boundary of ellipsoid defined by  $W$



## Example

( $p = 20$ )



design uses two vectors, on boundary of ellipse defined by optimal  $W$

## Derivation of dual

first reformulate primal problem with new variable  $X$ :

$$\begin{array}{ll}\text{minimize} & \log \det X^{-1} \\ \text{subject to} & X = \sum_{k=1}^p \lambda_k v_k v_k^T, \quad \lambda \geq 0, \quad \mathbf{1}^T \lambda = 1\end{array}$$

$$L(X, \lambda, Z, z, \nu) = \log \det X^{-1} + \text{tr} \left( Z \left( X - \sum_{k=1}^p \lambda_k v_k v_k^T \right) \right) - z^T \lambda + \nu (\mathbf{1}^T \lambda - 1)$$

- ▶ minimize over  $X$  by setting gradient to zero:  $-X^{-1} + Z = 0$
- ▶ minimum over  $\lambda_k$  is  $-\infty$  unless  $-v_k^T Z v_k - z_k + \nu = 0$

dual problem

$$\begin{array}{ll}\text{maximize} & n + \log \det Z - \nu \\ \text{subject to} & v_k^T Z v_k \leq \nu, \quad k = 1, \dots, p\end{array}$$

change variable  $W = Z/\nu$ , and optimize over  $\nu$  to get dual of slide 7.21

## 8. Geometric problems

# Outline

Extremal volume ellipsoids

Centering

Classification

Placement and facility location

## Minimum volume ellipsoid around a set

- ▶ **Löwner-John ellipsoid** of a set  $C$ : minimum volume ellipsoid  $\mathcal{E}$  with  $C \subseteq \mathcal{E}$
- ▶ parametrize  $\mathcal{E}$  as  $\mathcal{E} = \{v \mid \|Av + b\|_2 \leq 1\}$ ; can assume  $A \in \mathbf{S}_{++}^n$
- ▶ **vol**  $\mathcal{E}$  is proportional to  $\det A^{-1}$ ; to find Löwner-John ellipsoid, solve problem

$$\begin{array}{ll} \text{minimize (over } A, b) & \log \det A^{-1} \\ \text{subject to} & \sup_{v \in C} \|Av + b\|_2 \leq 1 \end{array}$$

convex, but evaluating the constraint can be hard (for general  $C$ )

- ▶ **finite set**  $C = \{x_1, \dots, x_m\}$ :

$$\begin{array}{ll} \text{minimize (over } A, b) & \log \det A^{-1} \\ \text{subject to} & \|Ax_i + b\|_2 \leq 1, \quad i = 1, \dots, m \end{array}$$

also gives Löwner-John ellipsoid for polyhedron **conv** $\{x_1, \dots, x_m\}$

## Maximum volume inscribed ellipsoid

- ▶ maximum volume ellipsoid  $\mathcal{E}$  with  $\mathcal{E} \subseteq C$ ,  $C \subseteq \mathbf{R}^n$  convex
- ▶ parametrize  $\mathcal{E}$  as  $\mathcal{E} = \{Bu + d \mid \|u\|_2 \leq 1\}$ ; can assume  $B \in \mathbf{S}_{++}^n$
- ▶ **vol**  $\mathcal{E}$  is proportional to  $\det B$ ; can find  $\mathcal{E}$  by solving

$$\begin{array}{ll}\text{maximize} & \log \det B \\ \text{subject to} & \sup_{\|u\|_2 \leq 1} I_C(Bu + d) \leq 0\end{array}$$

(where  $I_C(x) = 0$  for  $x \in C$  and  $I_C(x) = \infty$  for  $x \notin C$ )  
convex, but evaluating the constraint can be hard (for general  $C$ )



- ▶ **polyhedron**  $\{x \mid a_i^T x \leq b_i, i = 1, \dots, m\}$ :

$$\begin{array}{ll}\text{maximize} & \log \det B \\ \text{subject to} & \|Ba_i\|_2 + a_i^T d \leq b_i, \quad i = 1, \dots, m\end{array}$$

(constraint follows from  $\sup_{\|u\|_2 \leq 1} a_i^T (Bu + d) = \|Ba_i\|_2 + a_i^T d$ )

## Efficiency of ellipsoidal approximations

- ▶  $C \subseteq \mathbf{R}^n$  convex, bounded, with nonempty interior
- ▶ Löwner-John ellipsoid, shrunk by a factor  $n$  (around its center), lies inside  $C$
- ▶ maximum volume inscribed ellipsoid, expanded by a factor  $n$  (around its center) covers  $C$
- ▶ **example** (for polyhedra in  $\mathbf{R}^2$ )



- ▶ factor  $n$  can be improved to  $\sqrt{n}$  if  $C$  is symmetric

# Outline

Extremal volume ellipsoids

Centering



Classification

Placement and facility location



## Centering

- ▶ many possible definitions of ‘center’ of a convex set  $C$
- ▶ Chebyshev center: center of largest inscribed ball
  - for polyhedron, can be found via linear programming
- ▶ center of maximum volume inscribed ellipsoid
  - invariant under affine coordinate transformations



## Analytic center of a set of inequalities

- ▶ the **analytic center** of set of convex inequalities and linear equations

$$f_i(x) \leq 0, \quad i = 1, \dots, m, \quad Fx = g$$


is defined as solution of

$$\begin{array}{ll} \text{minimize} & -\sum_{i=1}^m \log(-f_i(x)) \\ \text{subject to} & Fx = g \end{array}$$

- ▶ objective is called the **log-barrier** for the inequalities
- ▶ (we'll see later) analytic center more easily computed than MVE or Chebyshev center
- ▶ two sets of inequalities can describe the same set, but have different analytic centers

## Analytic center of linear inequalities

- ▶  $a_i^T x \leq b_i, i = 1, \dots, m$
- ▶  $x_{ac}$  minimizes  $\phi(x) = -\sum_{i=1}^m \log(b_i - a_i^T x)$
- ▶ dashed lines are level curves of  $\phi$



## Inner and outer ellipsoids from analytic center

- ▶ we have

$$\mathcal{E}_{\text{inner}} \subseteq \{x \mid a_i^T x \leq b_i, \ i = 1, \dots, m\} \subseteq \mathcal{E}_{\text{outer}}$$

where

$$\begin{aligned}\mathcal{E}_{\text{inner}} &= \{x \mid (x - x_{\text{ac}})^T \nabla^2 \phi(x_{\text{ac}}) (x - x_{\text{ac}}) \leq 1\} \\ \mathcal{E}_{\text{outer}} &= \{x \mid (x - x_{\text{ac}})^T \nabla^2 \phi(x_{\text{ac}}) (x - x_{\text{ac}}) \leq m(m-1)\}\end{aligned}$$

- ▶ ellipsoid expansion/shrinkage factor is  $\sqrt{m(m-1)}$   
(cf.  $n$  for Löwner-John or max volume inscribed ellipsoids)

# Outline

Extremal volume ellipsoids

Centering

Classification

Placement and facility location

## Linear discrimination


- ▶ separate two sets of points  $\{x_1, \dots, x_N\}$ ,  $\{y_1, \dots, y_M\}$  by a hyperplane
- ▶ *i.e.*, find  $a \in \mathbf{R}^n$ ,  $b \in \mathbf{R}$  with

$$a^T x_i + b > 0, \quad i = 1, \dots, N, \quad a^T y_i + b < 0, \quad i = 1, \dots, M$$

- ▶ homogeneous in  $a$ ,  $b$ , hence equivalent to

$$a^T x_i + b \geq 1, \quad i = 1, \dots, N, \quad a^T y_i + b \leq -1, \quad i = 1, \dots, M$$

a set of linear inequalities in  $a$ ,  $b$ , *i.e.*, an LP feasibility problem



## Robust linear discrimination

(Euclidean) distance between hyperplanes

$$\mathcal{H}_1 = \{z \mid a^T z + b = 1\}$$


$$\mathcal{H}_2 = \{z \mid a^T z + b = -1\}$$

is  $\text{dist}(\mathcal{H}_1, \mathcal{H}_2) = 2/\|a\|_2$

to separate two sets of points by maximum margin,

$$\begin{aligned} & \text{minimize} && (1/2)\|a\|_2^2 \\ & \text{subject to} && a^T x_i + b \geq 1, \quad i = 1, \dots, N \\ & && a^T y_i + b \leq -1, \quad i = 1, \dots, M \end{aligned} \tag{2}$$


a QP in  $a, b$



## Approximate linear separation of non-separable sets

$$\begin{array}{ll}\text{minimize} & \mathbf{1}^T u + \mathbf{1}^T v \\ \text{subject to} & a^T x_i + b \geq 1 - u_i, \quad i = 1, \dots, N, \quad a^T y_i + b \leq -1 + v_i, \quad i = 1, \dots, M \\ & u \geq 0, \quad v \geq 0\end{array}$$

- ▶ an LP in  $a, b, u, v$
- ▶ at optimum,  $u_i = \max\{0, 1 - a^T x_i - b\}$ ,  $v_i = \max\{0, 1 + a^T y_i + b\}$
- ▶ equivalent to minimizing the sum of violations of the original inequalities






## Support vector classifier

$$\begin{array}{ll}\text{minimize} & \|a\|_2 + \gamma(\mathbf{1}^T u + \mathbf{1}^T v) \\ \text{subject to} & a^T x_i + b \geq 1 - u_i, \quad i = 1, \dots, N \\ & a^T y_i + b \leq -1 + v_i, \quad i = 1, \dots, M \\ & u \geq 0, \quad v \geq 0\end{array}$$

produces point on trade-off curve between inverse of margin  $2/\|a\|_2$  and classification error, measured by total slack  $\mathbf{1}^T u + \mathbf{1}^T v$

example on previous slide, with  $\gamma = 0.1$ :



## Nonlinear discrimination

- ▶ separate two sets of points by a nonlinear function  $f$ : find  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  with

$$f(x_i) > 0, \quad i = 1, \dots, N, \quad f(y_i) < 0, \quad i = 1, \dots, M$$

- ▶ choose a linearly parametrized family of functions  $f(z) = \theta^T F(z)$ 
  - $\theta \in \mathbf{R}^k$  is parameter
  - $F = (F_1, \dots, F_k) : \mathbf{R}^n \rightarrow \mathbf{R}^k$  are basis functions
- ▶ solve a set of linear inequalities in  $\theta$ :

$$\theta^T F(x_i) \geq 1, \quad i = 1, \dots, N, \quad \theta^T F(y_i) \leq -1, \quad i = 1, \dots, M$$

## Examples


- ▶ **quadratic discrimination:**  $f(z) = z^T P z + q^T z + r$ ,  $\theta = (P, q, r)$
- ▶ solve LP feasibility problem with variables  $P \in \mathbf{S}^n$ ,  $q \in \mathbf{R}^n$ ,  $r \in \mathbf{R}$

$$x_i^T P x_i + q^T x_i + r \geq 1, \quad y_i^T P y_i + q^T y_i + r \leq -1$$


- ▶ can add additional constraints (e.g.,  $P \leq -I$  to separate by an ellipsoid)
- ▶ **polynomial discrimination:**  $F(z)$  are all monomials up to a given degree  $d$
- ▶ e.g., for  $n = 2$ ,  $d = 3$

$$F(z) = (1, z_1, z_2, z_1^2, z_1 z_2, z_2^2, z_1^3, z_1^2 z_2, z_1 z_2^2, z_2^3)$$

## Example



separation by ellipsoid



separation by 4th degree polynomial

# Outline

Extremal volume ellipsoids

Centering

Classification




Placement and facility location

## Placement and facility location




- ▶  $N$  points with coordinates  $x_i \in \mathbf{R}^2$  (or  $\mathbf{R}^3$ )
- ▶ some positions  $x_i$  are given; the other  $x_i$ 's are variables
- ▶ for each pair of points, a cost function  $f_{ij}(x_i, x_j)$
- ▶ **placement problem:** minimize  $\sum_{i \neq j} f_{ij}(x_i, x_j)$
- ▶ **interpretations**
  - points are locations of plants or warehouses;  $f_{ij}$  is transportation cost between facilities  $i$  and  $j$
  - points are locations of cells in an integrated circuit;  $f_{ij}$  represents wirelength

## Example

- ▶ minimize  $\sum_{(i,j) \in \mathcal{E}} h(\|x_i - x_j\|_2)$ , with 6 free points, 27 edges
- ▶ optimal placements for  $h(z) = z$ ,  $h(z) = z^2$ ,  $h(z) = z^4$



- ▶ histograms of edge lengths  $\|x_i - x_j\|_2$ ,  $(i, j) \in \mathcal{E}$



## B. Numerical linear algebra background



# Outline

Flop counts and BLAS

Solving systems of linear equations

Block elimination

## Flop count

- ▶ **flop** (floating-point operation): one addition, subtraction, multiplication, or division of two floating-point numbers
- ▶ to estimate complexity of an algorithm
  - express number of flops as a (polynomial) function of the problem dimensions
  - simplify by keeping only the leading terms
- ▶ not an accurate predictor of computation time on modern computers, but useful as a rough estimate of complexity

## Basic linear algebra subroutines (BLAS)

**vector-vector operations** ( $x, y \in \mathbf{R}^n$ ) (BLAS level 1)

- ▶ inner product  $x^T y$ :  $2n - 1$  flops ( $\approx 2n, O(n)$ )
- ▶ sum  $x + y$ , scalar multiplication  $\alpha x$ :  $n$  flops

**matrix-vector product**  $y = Ax$  with  $A \in \mathbf{R}^{m \times n}$  (BLAS level 2)

- ▶  $m(2n - 1)$  flops ( $\approx 2mn$ )
- ▶  $2N$  if  $A$  is sparse with  $N$  nonzero elements
- ▶  $2p(n + m)$  if  $A$  is given as  $A = UV^T$ ,  $U \in \mathbf{R}^{m \times p}$ ,  $V \in \mathbf{R}^{n \times p}$

**matrix-matrix product**  $C = AB$  with  $A \in \mathbf{R}^{m \times n}$ ,  $B \in \mathbf{R}^{n \times p}$  (BLAS level 3)

- ▶  $mp(2n - 1)$  flops ( $\approx 2mnp$ )
- ▶ less if  $A$  and/or  $B$  are sparse
- ▶  $(1/2)m(m + 1)(2n - 1) \approx m^2 n$  if  $m = p$  and  $C$  symmetric

## BLAS on modern computers

- ▶ there are good implementations of BLAS and variants (*e.g.*, for sparse matrices)
- ▶ CPU single thread speeds typically 1–10 Gflops/s ( $10^9$  flops/sec)
- ▶ CPU multi threaded speeds typically 10–100 Gflops/s
- ▶ GPU speeds typically 100 Gflops/s–1 Tflops/s ( $10^{12}$  flops/sec)

# Outline

Flop counts and BLAS

Solving systems of linear equations

Block elimination

## Complexity of solving linear equations

- ▶  $A \in \mathbf{R}^{n \times n}$  is invertible,  $b \in \mathbf{R}^n$
- ▶ solution of  $Ax = b$  is  $x = A^{-1}b$
- ▶ solving  $Ax = b$ , i.e., computing  $x = A^{-1}b$ 
  - almost never done by computing  $A^{-1}$ , then multiplying by  $b$
  - for general methods,  $O(n^3)$
  - (much) less if  $A$  is structured (banded, sparse, Toeplitz, ...)
  - e.g., for  $A$  with half-bandwidth  $k$  ( $A_{ij} = 0$  for  $|i - j| > k$ ,  $O(k^2n)$ )
- ▶ it's super useful to recognize matrix structure that can be exploited in solving  $Ax = b$

## Linear equations that are easy to solve

- ▶ diagonal matrices:  $n$  flops;  $x = A^{-1}b = (b_1/a_{11}, \dots, b_n/a_{nn})$
- ▶ lower triangular:  $n^2$  flops via **forward substitution**

$$\begin{aligned}x_1 &:= b_1/a_{11} \\x_2 &:= (b_2 - a_{21}x_1)/a_{22} \\x_3 &:= (b_3 - a_{31}x_1 - a_{32}x_2)/a_{33} \\&\vdots \\x_n &:= (b_n - a_{n1}x_1 - a_{n2}x_2 - \dots - a_{n,n-1}x_{n-1})/a_{nn}\end{aligned}$$

- ▶ upper triangular:  $n^2$  flops via **backward substitution**

## Linear equations that are easy to solve

- ▶ orthogonal matrices ( $A^{-1} = A^T$ ):
  - $2n^2$  flops to compute  $x = A^T b$  for general  $A$
  - less with structure, e.g., if  $A = I - 2uu^T$  with  $\|u\|_2 = 1$ , we can compute  $x = A^T b = b - 2(u^T b)u$  in  $4n$  flops
- ▶ permutation matrices: for  $\pi = (\pi_1, \pi_2, \dots, \pi_n)$  a permutation of  $(1, 2, \dots, n)$

$$a_{ij} = \begin{cases} 1 & j = \pi_i \\ 0 & \text{otherwise} \end{cases}$$

- interpretation:  $Ax = (x_{\pi_1}, \dots, x_{\pi_n})$
- satisfies  $A^{-1} = A^T$ , hence cost of solving  $Ax = b$  is 0 flops
- example:

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad A^{-1} = A^T = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$



## Factor-solve method for solving $Ax = b$

- ▶ factor  $A$  as a product of simple matrices (usually 2–5):

$$A = A_1 A_2 \cdots A_k$$

- ▶ e.g.,  $A_i$  diagonal, upper or lower triangular, orthogonal, permutation, ...
- ▶ compute  $x = A^{-1}b = A_k^{-1} \cdots A_2^{-1} A_1^{-1} b$  by solving  $k$  ‘easy’ systems of equations

$$A_1 x_1 = b, \quad A_2 x_2 = x_1, \quad \dots \quad A_k x_k = x_{k-1}$$

- ▶ cost of factorization step usually dominates cost of solve step

## Solving equations with multiple righthand sides

- ▶ we wish to solve

$$Ax_1 = b_1, \quad Ax_2 = b_2, \quad \dots \quad Ax_m = b_m$$

- ▶ cost: one factorization plus  $m$  solves
- ▶ called **factorization caching**
- ▶ when factorization cost dominates solve cost, we can solve a modest number of equations at the same cost as one (!!)

## LU factorization

- ▶ every nonsingular matrix  $A$  can be factored as  $A = PLU$  with  $P$  a permutation,  $L$  lower triangular,  $U$  upper triangular
- ▶ factorization cost:  $(2/3)n^3$  flops

---

*Solving linear equations by LU factorization.*

**given** a set of linear equations  $Ax = b$ , with  $A$  nonsingular.

1. *LU factorization.* Factor  $A$  as  $A = PLU$  ( $(2/3)n^3$  flops).
2. *Permutation.* Solve  $Pz_1 = b$  (0 flops).
3. *Forward substitution.* Solve  $Lz_2 = z_1$  ( $n^2$  flops).
4. *Backward substitution.* Solve  $Ux = z_2$  ( $n^2$  flops).

- 
- ▶ total cost:  $(2/3)n^3 + 2n^2 \approx (2/3)n^3$  for large  $n$

## Sparse LU factorization

- ▶ for  $A$  sparse and invertible, factor as  $A = P_1 L U P_2$
- ▶ adding permutation matrix  $P_2$  offers possibility of sparser  $L, U$
- ▶ hence, less storage and cheaper factor and solve steps
- ▶  $P_1$  and  $P_2$  chosen (heuristically) to yield sparse  $L, U$
- ▶ choice of  $P_1$  and  $P_2$  depends on sparsity pattern and values of  $A$
- ▶ cost is usually much less than  $(2/3)n^3$ ; exact value depends in a complicated way on  $n$ , number of zeros in  $A$ , sparsity pattern
- ▶ often practical to solve very large sparse systems of equations

## Cholesky factorization

- ▶ every positive definite  $A$  can be factored as  $A = LL^T$
- ▶  $L$  is lower triangular with positive diagonal entries
- ▶ Cholesky factorization cost:  $(1/3)n^3$  flops

---

*Solving linear equations by Cholesky factorization.*

**given** a set of linear equations  $Ax = b$ , with  $A \in \mathbf{S}_{++}^n$ .

1. *Cholesky factorization.* Factor  $A$  as  $A = LL^T$  ( $(1/3)n^3$  flops).
2. *Forward substitution.* Solve  $Lz_1 = b$  ( $n^2$  flops).
3. *Backward substitution.* Solve  $L^T x = z_1$  ( $n^2$  flops).

- 
- ▶ total cost:  $(1/3)n^3 + 2n^2 \approx (1/3)n^3$  for large  $n$

## Sparse Cholesky factorization

- ▶ for sparse positive definite  $A$ , factor as  $A = PLL^T P^T$
- ▶ adding permutation matrix  $P$  offers possibility of sparser  $L$
- ▶ same as
  - permuting rows and columns of  $A$  to get  $\tilde{A} = P^T A P$
  - then finding Cholesky factorization of  $\tilde{A}$
- ▶  $P$  chosen (heuristically) to yield sparse  $L$
- ▶ choice of  $P$  only depends on sparsity pattern of  $A$  (unlike sparse LU)
- ▶ cost is usually much less than  $(1/3)n^3$ ; exact value depends in a complicated way on  $n$ , number of zeros in  $A$ , sparsity pattern

## Example

- ▶ sparse  $A$  with upper arrow sparsity pattern

$$A = \begin{bmatrix} * & * & * & * & * \\ * & * & & & \\ * & & * & & \\ * & & & * & \\ * & & & & * \end{bmatrix} \quad L = \begin{bmatrix} * & & & & \\ * & * & & & \\ * & * & * & & \\ * & * & * & * & \\ * & * & * & * & * \end{bmatrix}$$

$L$  is full, with  $O(n^2)$  nonzeros; solve cost is  $O(n^2)$

- ▶ reverse order of entries (*i.e.*, permute) to get lower arrow sparsity pattern

$$\tilde{A} = \begin{bmatrix} * & & & & * \\ & * & & & * \\ & & * & & * \\ & & & * & * \\ * & * & * & * & * \end{bmatrix} \quad L = \begin{bmatrix} * & & & & \\ & * & & & \\ & & * & & \\ & & & * & \\ * & * & * & * & * \end{bmatrix}$$

$L$  is sparse with  $O(n)$  nonzeros; cost of solve is  $O(n)$

## LDL<sup>T</sup> factorization

- ▶ every nonsingular symmetric matrix  $A$  can be factored as

$$A = PLDL^T P^T$$

with  $P$  a permutation matrix,  $L$  lower triangular,  $D$  block diagonal with  $1 \times 1$  or  $2 \times 2$  diagonal blocks

- ▶ factorization cost:  $(1/3)n^3$
- ▶ cost of solving linear equations with symmetric  $A$  by LDL<sup>T</sup> factorization:  
 $(1/3)n^3 + 2n^2 \approx (1/3)n^3$  for large  $n$
- ▶ for sparse  $A$ , can choose  $P$  to yield sparse  $L$ ; cost  $\ll (1/3)n^3$



# Outline

Flop counts and BLAS

Solving systems of linear equations

Block elimination

## Equations with structured sub-blocks

- express  $Ax = b$  in blocks as

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

with  $x_1 \in \mathbf{R}^{n_1}$ ,  $x_2 \in \mathbf{R}^{n_2}$ ; blocks  $A_{ij} \in \mathbf{R}^{n_i \times n_j}$

- assuming  $A_{11}$  is nonsingular, can eliminate  $x_1$  as

$$x_1 = A_{11}^{-1}(b_1 - A_{12}x_2)$$

- to compute  $x_2$ , solve

$$(A_{22} - A_{21}A_{11}^{-1}A_{12})x_2 = b_2 - A_{21}A_{11}^{-1}b_1$$

- $S = A_{22} - A_{21}A_{11}^{-1}A_{12}$  is the **Schur complement**

## Bock elimination method

---

*Solving linear equations by block elimination.*

**given** a nonsingular set of linear equations with  $A_{11}$  nonsingular.

1. Form  $A_{11}^{-1}A_{12}$  and  $A_{11}^{-1}b_1$ .
  2. Form  $S = A_{22} - A_{21}A_{11}^{-1}A_{12}$  and  $\tilde{b} = b_2 - A_{21}A_{11}^{-1}b_1$ .
  3. Determine  $x_2$  by solving  $Sx_2 = \tilde{b}$ .
  4. Determine  $x_1$  by solving  $A_{11}x_1 = b_1 - A_{12}x_2$ .
- 

### dominant terms in flop count

- ▶ step 1:  $f + n_2s$  ( $f$  is cost of factoring  $A_{11}$ ;  $s$  is cost of solve step)
- ▶ step 2:  $2n_2^2n_1$  (cost dominated by product of  $A_{21}$  and  $A_{11}^{-1}A_{12}$ )
- ▶ step 3:  $(2/3)n_2^3$

total:  $f + n_2s + 2n_2^2n_1 + (2/3)n_2^3$

## Examples

- ▶ for general  $A_{11}$ ,  $f = (2/3)n_1^3$ ,  $s = 2n_1^2$

$$\text{\#flops} = (2/3)n_1^3 + 2n_1^2n_2 + 2n_2^2n_1 + (2/3)n_2^3 = (2/3)(n_1 + n_2)^3$$

so, no gain over standard method

- ▶ block elimination is useful for structured  $A_{11}$  ( $f \ll n_1^3$ )
- ▶ for example,  $A_{11}$  diagonal ( $f = 0$ ,  $s = n_1$ ):  $\text{\#flops} \approx 2n_2^2n_1 + (2/3)n_2^3$

## Structured plus low rank matrices

- ▶ we wish to solve  $(A + BC)x = b$ ,  $A \in \mathbf{R}^{n \times n}$ ,  $B \in \mathbf{R}^{n \times p}$ ,  $C \in \mathbf{R}^{p \times n}$
- ▶ assume  $A$  has structure (*i.e.*,  $Ax = b$  easy to solve)
- ▶ first **uneliminate** to write as block equations with new variable  $y$

$$\begin{bmatrix} A & B \\ C & -I \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} b \\ 0 \end{bmatrix}$$

- ▶ now apply block elimination: solve

$$(I + CA^{-1}B)y = CA^{-1}b,$$

then solve  $Ax = b - By$

- ▶ this proves the **matrix inversion lemma**: if  $A$  and  $A + BC$  are nonsingular,

$$(A + BC)^{-1} = A^{-1} - A^{-1}B(I + CA^{-1}B)^{-1}CA^{-1}$$

## Example: Solving diagonal plus low rank equations

- ▶ with  $A$  diagonal,  $p \ll n$ ,  $A + BC$  is called **diagonal plus low rank**
- ▶ for covariance matrices, called a **factor model**
- ▶ method 1: form  $D = A + BC$ , then solve  $Dx = b$ 
  - storage  $n^2$
  - solve cost  $(2/3)n^3 + 2pn^2$  (**cubic** in  $n$ )
- ▶ method 2: solve  $(I + CA^{-1}B)y = CA^{-1}b$ , then compute  $x = A^{-1}b - A^{-1}By$ 
  - storage  $O(np)$
  - solve cost  $2p^2n + (2/3)p^3$  (**linear** in  $n$ )

## 9. Unconstrained minimization

# Outline

Terminology and assumptions

Gradient descent method

Steepest descent method

Newton's method

Self-concordant functions

Implementation



## Unconstrained minimization

- ▶ unconstrained minimization problem

$$\text{minimize } f(x)$$

- ▶ we assume
  - $f$  convex, twice continuously differentiable (hence **dom**  $f$  open)
  - optimal value  $p^\star = \inf_x f(x)$  is attained at  $x^\star$  (not necessarily unique)
- ▶ optimality condition is  $\nabla f(x) = 0$
- ▶ minimizing  $f$  is the same as solving  $\nabla f(x) = 0$
- ▶ a set of  $n$  equations with  $n$  unknowns

## Quadratic functions

- ▶ convex quadratic:  $f(x) = (1/2)x^T Px + q^T x + r, P \succeq 0$
- ▶ we can solve exactly via linear equations

$$\nabla f(x) = Px + q = 0$$

- ▶ much more on this special case later

## Iterative methods

- ▶ for most non-quadratic functions, we use **iterative methods**
- ▶ these produce a sequence of points  $x^{(k)} \in \text{dom} f$ ,  $k = 0, 1, \dots$
- ▶  $x^{(0)}$  is the **initial point** or **starting point**
- ▶  $x^{(k)}$  is the  $k$ th **iterate**
- ▶ we hope that the method **converges**, *i.e.*,

$$f(x^{(k)}) \rightarrow p^\star, \quad \nabla f(x^{(k)}) \rightarrow 0$$

## Initial point and sublevel set

- ▶ algorithms in this chapter require a starting point  $x^{(0)}$  such that
  - $x^{(0)} \in \mathbf{dom} f$
  - sublevel set  $S = \{x \mid f(x) \leq f(x^{(0)})\}$  is closed
- ▶ 2nd condition is hard to verify, except when **all** sublevel sets are closed
  - equivalent to condition that **epi**  $f$  is closed
  - true if  $\mathbf{dom} f = \mathbf{R}^n$
  - true if  $f(x) \rightarrow \infty$  as  $x \rightarrow \mathbf{bd} \mathbf{dom} f$
- ▶ examples of differentiable functions with closed sublevel sets:

$$f(x) = \log \left( \sum_{i=1}^m \exp(a_i^T x + b_i) \right), \quad f(x) = - \sum_{i=1}^m \log(b_i - a_i^T x)$$

## Strong convexity and implications

- ▶  $f$  is **strongly convex** on  $S$  if there exists an  $m > 0$  such that

$$\nabla^2 f(x) \succeq mI \text{ for all } x \in S$$

- ▶ same as  $f(x) - (m/2)\|x\|_2^2$  is convex
- ▶ if  $f$  is strongly convex, for  $x, y \in S$ ,

$$f(y) \geq f(x) + \nabla f(x)^T (y - x) + \frac{m}{2} \|x - y\|_2^2$$

- ▶ hence,  $S$  is bounded
- ▶ we conclude  $p^\star > -\infty$ , and for  $x \in S$ ,

$$f(x) - p^\star \leq \frac{1}{2m} \|\nabla f(x)\|_2^2$$

- ▶ useful as stopping criterion (if you know  $m$ , which usually you do not)

# Outline

Terminology and assumptions

Gradient descent method

Steepest descent method

Newton's method

Self-concordant functions

Implementation

## Descent methods

- ▶ **descent methods** generate iterates as

$$x^{(k+1)} = x^{(k)} + t^{(k)} \Delta x^{(k)}$$

with  $f(x^{(k+1)}) < f(x^{(k)})$  (hence the name)

- ▶ other notations:  $x^+ = x + t\Delta x$ ,  $x := x + t\Delta x$
- ▶  $\Delta x^{(k)}$  is the **step**, or **search direction**
- ▶  $t^{(k)} > 0$  is the **step size**, or **step length**
- ▶ from convexity,  $f(x^+) < f(x)$  implies  $\nabla f(x)^T \Delta x < 0$
- ▶ this means  $\Delta x$  is a **descent direction**

## Generic descent method

---

**General descent method.**

**given** a starting point  $x \in \text{dom } f$ .

**repeat**

1. Determine a descent direction  $\Delta x$ .
2. **Line search.** Choose a step size  $t > 0$ .
3. **Update.**  $x := x + t\Delta x$ .


**until** stopping criterion is satisfied.

---



## Line search types

- ▶ **exact line search:**  $t = \operatorname{argmin}_{t \geq 0} f(x + t\Delta x)$
- ▶ **backtracking line search** (with parameters  $\alpha \in (0, 1/2)$ ,  $\beta \in (0, 1)$ )
  - starting at  $t = 1$ , repeat  $t := \beta t$  until  $f(x + t\Delta x) < f(x) + \alpha t \nabla f(x)^T \Delta x$
- ▶ graphical interpretation: reduce  $t$  (i.e., backtrack) until  $t \leq t_0$



## Gradient descent method

- ▶ general descent method with  $\Delta x = -\nabla f(x)$

---

**given** a starting point  $x \in \text{dom } f$ .

**repeat**

1.  $\Delta x := -\nabla f(x)$ .
2. **Line search.** Choose step size  $t$  via exact or backtracking line search.
3. **Update.**  $x := x + t\Delta x$ .

**until** stopping criterion is satisfied.

---

- ▶ stopping criterion usually of the form  $\|\nabla f(x)\|_2 \leq \epsilon$
- ▶ convergence result: for strongly convex  $f$ ,

$$f(x^{(k)}) - p^\star \leq c^k (f(x^{(0)}) - p^\star)$$

$c \in (0, 1)$  depends on  $m$ ,  $x^{(0)}$ , line search type


- ▶ very simple, but can be very slow

## Example: Quadratic function on $\mathbf{R}^2$

- ▶ take  $f(x) = (1/2)(x_1^2 + \gamma x_2^2)$ , with  $\gamma > 0$
- ▶ with exact line search, starting at  $x^{(0)} = (\gamma, 1)$ :


$$x_1^{(k)} = \gamma \left( \frac{\gamma - 1}{\gamma + 1} \right)^k, \quad x_2^{(k)} = \left( -\frac{\gamma - 1}{\gamma + 1} \right)^k$$

- very slow if  $\gamma \gg 1$  or  $\gamma \ll 1$
- example for  $\gamma = 10$  at right
- called **zig-zagging**




## Example: Nonquadratic function on $\mathbb{R}^2$

►  $f(x_1, x_2) = e^{x_1+3x_2-0.1} + e^{x_1-3x_2-0.1} + e^{-x_1-0.1}$




backtracking line search



exact line search

## Example: A problem in $\mathbf{R}^{100}$

- ▶  $f(x) = c^T x - \sum_{i=1}^{500} \log(b_i - a_i^T x)$
- ▶ **linear convergence**, *i.e.*, a straight line on a semilog plot



# Outline

Terminology and assumptions

Gradient descent method

**Steepest descent method**

Newton's method

Self-concordant functions

Implementation

## Steepest descent method



- ▶ **normalized steepest descent direction** (at  $x$ , for norm  $\|\cdot\|$ ):

$$\Delta x_{\text{nsd}} = \operatorname{argmin}\{\nabla f(x)^T v \mid \|v\| = 1\}$$

- ▶ interpretation: for small  $v$ ,  $f(x+v) \approx f(x) + \nabla f(x)^T v$ ;
- ▶ direction  $\Delta x_{\text{nsd}}$  is unit-norm step with most negative directional derivative
- ▶ **(unnormalized) steepest descent direction:**  $\Delta x_{\text{sd}} = \|\nabla f(x)\|_* \Delta x_{\text{nsd}}$
- ▶ satisfies  $\nabla f(x)^T \Delta x_{\text{sd}} = -\|\nabla f(x)\|_*^2$
- ▶ **steepest descent method**
  - general descent method with  $\Delta x = \Delta x_{\text{sd}}$
  - convergence properties similar to gradient descent



## Examples

- ▶ Euclidean norm:  $\Delta x_{\text{sd}} = -\nabla f(x)$
- ▶ quadratic norm  $\|x\|_P = (x^T P x)^{1/2}$  ( $P \in \mathbf{S}_{++}^n$ ):  $\Delta x_{\text{sd}} = -P^{-1} \nabla f(x)$
- ▶  $\ell_1$ -norm:  $\Delta x_{\text{sd}} = -(\partial f(x) / \partial x_i) e_i$ , where  $|\partial f(x) / \partial x_i| = \|\nabla f(x)\|_\infty$
- ▶ unit balls, normalized steepest descent directions for quadratic norm and  $\ell_1$ -norm:





## Choice of norm for steepest descent



- ▶ steepest descent with backtracking line search for two quadratic norms
- ▶ ellipses show  $\{x \mid \|x - x^{(k)}\|_P = 1\}$
- ▶ interpretation of steepest descent with quadratic norm  $\|\cdot\|_P$ : gradient descent after change of variables  $\bar{x} = P^{1/2}x$
- ▶ shows choice of  $P$  has strong effect on speed of convergence

# Outline

Terminology and assumptions

Gradient descent method

Steepest descent method

**Newton's method**


Self-concordant functions

Implementation

## Newton step

- ▶ **Newton step** is  $\Delta x_{\text{nt}} = -\nabla^2 f(x)^{-1} \nabla f(x)$
- ▶ **interpretation:**  $x + \Delta x_{\text{nt}}$  minimizes second order approximation


$$\widehat{f}(x+v) = f(x) + \nabla f(x)^T v + \frac{1}{2} v^T \nabla^2 f(x) v$$



## Another interpretation


- ▶  $x + \Delta x_{\text{nt}}$  solves linearized optimality condition

$$\nabla f(x + v) \approx \widehat{\nabla f}(x + v) = \nabla f(x) + \nabla^2 f(x)v = 0$$



## And one more interpretation

- ▶  $\Delta x_{\text{nt}}$  is steepest descent direction at  $x$  in local Hessian norm  $\|u\|_{\nabla^2 f(x)} = (u^T \nabla^2 f(x) u)^{1/2}$



- ▶ dashed lines are contour lines of  $f$ ; ellipse is  $\{x + v \mid v^T \nabla^2 f(x) v = 1\}$
- ▶ arrow shows  $-\nabla f(x)$

## Newton decrement

- ▶ **Newton decrement** is  $\lambda(x) = (\nabla f(x)^T \nabla^2 f(x)^{-1} \nabla f(x))^{1/2}$
- ▶ a measure of the proximity of  $x$  to  $x^\star$
- ▶ gives an estimate of  $f(x) - p^\star$ , using quadratic approximation  $\widehat{f}$ :

$$f(x) - \inf_y \widehat{f}(y) = \frac{1}{2} \lambda(x)^2$$

- ▶ equal to the norm of the Newton step in the quadratic Hessian norm

$$\lambda(x) = \left( \Delta x_{\text{nt}}^T \nabla^2 f(x) \Delta x_{\text{nt}} \right)^{1/2}$$

- ▶ directional derivative in the Newton direction:  $\nabla f(x)^T \Delta x_{\text{nt}} = -\lambda(x)^2$
- ▶ affine invariant (unlike  $\|\nabla f(x)\|_2$ )

## Newton's method

---

**given** a starting point  $x \in \text{dom } f$ , tolerance  $\epsilon > 0$ .

**repeat**

1. **Compute the Newton step and decrement.**

$$\Delta x_{\text{nt}} := -\nabla^2 f(x)^{-1} \nabla f(x); \quad \lambda^2 := \nabla f(x)^T \nabla^2 f(x)^{-1} \nabla f(x).$$

2. **Stopping criterion.** **quit** if  $\lambda^2/2 \leq \epsilon$ .

3. **Line search.** Choose step size  $t$  by backtracking line search.

4. **Update.**  $x := x + t\Delta x_{\text{nt}}$ .

---

- ▶ **affine invariant**, *i.e.*, independent of linear changes of coordinates
- ▶ Newton iterates for  $\tilde{f}(y) = f(Ty)$  with starting point  $y^{(0)} = T^{-1}x^{(0)}$  are  $y^{(k)} = T^{-1}x^{(k)}$

## Classical convergence analysis

### assumptions

- ▶  $f$  strongly convex on  $S$  with constant  $m$
- ▶  $\nabla^2 f$  is Lipschitz continuous on  $S$ , with constant  $L > 0$ :

$$\|\nabla^2 f(x) - \nabla^2 f(y)\|_2 \leq L\|x - y\|_2$$

( $L$  measures how well  $f$  can be approximated by a quadratic function)

**outline:** there exist constants  $\eta \in (0, m^2/L)$ ,  $\gamma > 0$  such that

- ▶ if  $\|\nabla f(x)\|_2 \geq \eta$ , then  $f(x^{(k+1)}) - f(x^{(k)}) \leq -\gamma$
- ▶ if  $\|\nabla f(x)\|_2 < \eta$ , then

$$\frac{L}{2m^2} \|\nabla f(x^{(k+1)})\|_2 \leq \left( \frac{L}{2m^2} \|\nabla f(x^{(k)})\|_2 \right)^2$$



## Classical convergence analysis

**damped Newton phase** ( $\|\nabla f(x)\|_2 \geq \eta$ )

- ▶ most iterations require backtracking steps
- ▶ function value decreases by at least  $\gamma$
- ▶ if  $p^\star > -\infty$ , this phase ends after at most  $(f(x^{(0)}) - p^\star)/\gamma$  iterations

**quadratically convergent phase** ( $\|\nabla f(x)\|_2 < \eta$ )

- ▶ all iterations use step size  $t = 1$
- ▶  $\|\nabla f(x)\|_2$  converges to zero quadratically: if  $\|\nabla f(x^{(k)})\|_2 < \eta$ , then

$$\frac{L}{2m^2} \|\nabla f(x^l)\|_2 \leq \left( \frac{L}{2m^2} \|\nabla f(x^k)\|_2 \right)^{2^{l-k}} \leq \left( \frac{1}{2} \right)^{2^{l-k}}, \quad l \geq k$$

## Classical convergence analysis



**conclusion:** number of iterations until  $f(x) - p^\star \leq \epsilon$  is bounded above by

$$\frac{f(x^{(0)}) - p^\star}{\gamma} + \log_2 \log_2(\epsilon_0/\epsilon)$$

- ▶  $\gamma, \epsilon_0$  are constants that depend on  $m, L, x^{(0)}$
- ▶ second term is small (of the order of 6) and almost constant for practical purposes
- ▶ in practice, constants  $m, L$  (hence  $\gamma, \epsilon_0$ ) are usually unknown
- ▶ provides qualitative insight in convergence properties (*i.e.*, explains two algorithm phases)

## Example: $\mathbf{R}^2$



(same problem as slide 9.13)



- ▶ backtracking parameters  $\alpha = 0.1, \beta = 0.7$
- ▶ converges in only 5 steps
- ▶ quadratic local convergence

## Example in $\mathbf{R}^{100}$

(same problem as slide 9.14)




- ▶ backtracking parameters  $\alpha = 0.01$ ,  $\beta = 0.5$
- ▶ backtracking line search almost as fast as exact l.s. (and much simpler)
- ▶ clearly shows two phases in algorithm

## Example in $\mathbf{R}^{10000}$

(with sparse  $a_i$ )

$$f(x) = - \sum_{i=1}^{10000} \log(1 - x_i^2) - \sum_{i=1}^{100000} \log(b_i - a_i^T x)$$



- ▶ backtracking parameters  $\alpha = 0.01$ ,  $\beta = 0.5$ .
- ▶ performance similar as for small examples

# Outline

Terminology and assumptions

Gradient descent method

Steepest descent method

Newton's method

**Self-concordant functions**

Implementation

## Self-concordance

### shortcomings of classical convergence analysis

- ▶ depends on unknown constants ( $m, L, \dots$ )
- ▶ bound is not affinely invariant, although Newton's method is

### convergence analysis via self-concordance (Nesterov and Nemirovski)

- ▶ does not depend on any unknown constants
- ▶ gives affine-invariant bound
- ▶ applies to special class of convex **self-concordant** functions
- ▶ developed to analyze polynomial-time interior-point methods for convex optimization

## Convergence analysis for self-concordant functions

### definition

- ▶ convex  $f : \mathbf{R} \rightarrow \mathbf{R}$  is self-concordant if  $|f'''(x)| \leq 2f''(x)^{3/2}$  for all  $x \in \text{dom } f$
- ▶  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is self-concordant if  $g(t) = f(x + tv)$  is self-concordant for all  $x \in \text{dom } f$ ,  $v \in \mathbf{R}^n$

### examples on $\mathbf{R}$

- ▶ linear and quadratic functions
- ▶ negative logarithm  $f(x) = -\log x$
- ▶ negative entropy plus negative logarithm:  $f(x) = x \log x - \log x$

**affine invariance:** if  $f : \mathbf{R} \rightarrow \mathbf{R}$  is s.c., then  $\tilde{f}(y) = f(ay + b)$  is s.c.:

$$\tilde{f}'''(y) = a^3 f'''(ay + b), \quad \tilde{f}''(y) = a^2 f''(ay + b)$$



## Self-concordant calculus

### properties

- ▶ preserved under positive scaling  $\alpha \geq 1$ , and sum
- ▶ preserved under composition with affine function
- ▶ if  $g$  is convex with **dom**  $g = \mathbf{R}_{++}$  and  $|g'''(x)| \leq 3g''(x)/x$  then

$$f(x) = \log(-g(x)) - \log x$$

is self-concordant

**examples:** properties can be used to show that the following are s.c.

- ▶  $f(x) = -\sum_{i=1}^m \log(b_i - a_i^T x)$  on  $\{x \mid a_i^T x < b_i, i = 1, \dots, m\}$
- ▶  $f(X) = -\log \det X$  on  $\mathbf{S}_{++}^n$
- ▶  $f(x) = -\log(y^2 - x^T x)$  on  $\{(x, y) \mid \|x\|_2 < y\}$

## Convergence analysis for self-concordant functions

**summary:** there exist constants  $\eta \in (0, 1/4]$ ,  $\gamma > 0$  such that

- ▶ if  $\lambda(x) > \eta$ , then  $f(x^{(k+1)}) - f(x^{(k)}) \leq -\gamma$
- ▶ if  $\lambda(x) \leq \eta$ , then  $2\lambda(x^{(k+1)}) \leq (2\lambda(x^{(k)}))^2$

( $\eta$  and  $\gamma$  only depend on backtracking parameters  $\alpha, \beta$ )


**complexity bound:** number of Newton iterations bounded by

$$\frac{f(x^{(0)}) - p^\star}{\gamma} + \log_2 \log_2(1/\epsilon)$$

for  $\alpha = 0.1$ ,  $\beta = 0.8$ ,  $\epsilon = 10^{-10}$ , bound evaluates to  $375(f(x^{(0)}) - p^\star) + 6$

## Numerical example

- ▶ 150 randomly generated instances of  $f(x) = -\sum_{i=1}^m \log(b_i - a_i^T x)$ ,  $x \in \mathbf{R}^n$
- ▶  $\circ$ :  $m = 100, n = 50$ ;  $\square$ :  $m = 1000, n = 500$ ;  $\diamond$ :  $m = 1000, n = 50$



- ▶ number of iterations much smaller than  $375(f(x^{(0)}) - p^*) + 6$
- ▶ bound of the form  $c(f(x^{(0)}) - p^*) + 6$  with smaller  $c$  (empirically) valid

# Outline

Terminology and assumptions

Gradient descent method

Steepest descent method

Newton's method

Self-concordant functions

Implementation

## Implementation

main effort in each iteration: evaluate derivatives and solve Newton system

$$H\Delta x = -g$$

where  $H = \nabla^2 f(x)$ ,  $g = \nabla f(x)$

**via Cholesky factorization**

$$H = LL^T, \quad \Delta x_{\text{nt}} = -L^{-T}L^{-1}g, \quad \lambda(x) = \|L^{-1}g\|_2$$

- ▶ cost  $(1/3)n^3$  flops for unstructured system
- ▶ cost  $\ll (1/3)n^3$  if  $H$  is sparse, banded, or has other structure

## Example

- ▶  $f(x) = \sum_{i=1}^n \psi_i(x_i) + \psi_0(Ax + b)$ , with  $A \in \mathbf{R}^{p \times n}$  dense,  $p \ll n$
- ▶ Hessian has low rank plus diagonal structure  $H = D + A^T H_0 A$
- ▶  $D$  diagonal with diagonal elements  $\psi_i''(x_i)$ ;  $H_0 = \nabla^2 \psi_0(Ax + b)$

**method 1:** form  $H$ , solve via dense Cholesky factorization: (cost  $(1/3)n^3$ )

**method 2** (block elimination): factor  $H_0 = L_0 L_0^T$ ; write Newton system as

$$D\Delta x + A^T L_0 w = -g, \quad L_0^T A \Delta x - w = 0$$

eliminate  $\Delta x$  from first equation; compute  $w$  and  $\Delta x$  from

$$(I + L_0^T A D^{-1} A^T L_0) w = -L_0^T A D^{-1} g, \quad D\Delta x = -g - A^T L_0 w$$

cost:  $2p^2 n$  (dominated by computation of  $L_0^T A D^{-1} A^T L_0$ )

## 10. Equality constrained minimization

# Outline

Equality constrained minimization

Newton's method with equality constraints

Infeasible start Newton method

Implementation



## Equality constrained minimization

- ▶ equality constrained smooth minimization problem:

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & Ax = b\end{array}$$

- ▶ we assume

- $f$  convex, twice continuously differentiable
- $A \in \mathbf{R}^{p \times n}$  with  $\mathbf{rank} A = p$
- $p^\star$  is finite and attained

- ▶ **optimality conditions:**  $x^\star$  is optimal if and only if there exists a  $\nu^\star$  such that

$$\nabla f(x^\star) + A^T \nu^\star = 0, \quad Ax^\star = b$$

## Equality constrained quadratic minimization

- ▶  $f(x) = (1/2)x^T Px + q^T x + r, P \in \mathbf{S}_+^n$
- ▶  $\nabla f(x) = Px + q$
- ▶ optimality conditions are a **system of linear equations**

$$\begin{bmatrix} P & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} x^\star \\ v^\star \end{bmatrix} = \begin{bmatrix} -q \\ b \end{bmatrix}$$

- ▶ coefficient matrix is called KKT matrix
- ▶ KKT matrix is nonsingular if and only if

$$Ax = 0, \quad x \neq 0 \quad \implies \quad x^T Px > 0$$

- ▶ equivalent condition for nonsingularity:  $P + A^T A \succ 0$

## Eliminating equality constraints

- ▶ represent feasible set  $\{x \mid Ax = b\}$  as  $\{Fz + \hat{x} \mid z \in \mathbf{R}^{n-p}\}$ 
  - $\hat{x}$  is (any) **particular solution** of  $Ax = b$
  - range of  $F \in \mathbf{R}^{n \times (n-p)}$  is nullspace of  $A$  (**rank**  $F = n - p$  and  $AF = 0$ )
- ▶ **reduced or eliminated problem**: minimize  $f(Fz + \hat{x})$
- ▶ an unconstrained problem with variable  $z \in \mathbf{R}^{n-p}$
- ▶ from solution  $z^\star$ , obtain  $x^\star$  and  $v^\star$  as

$$x^\star = Fz^\star + \hat{x}, \quad v^\star = -(AA^T)^{-1}A\nabla f(x^\star)$$

## Example: Optimal resource allocation

- ▶ allocate resource amount  $x_i \in \mathbf{R}$  to agent  $i$
- ▶ agent  $i$  cost if  $f_i(x_i)$
- ▶ resource budget is  $b$ , so  $x_1 + \cdots + x_n = b$
- ▶ resource allocation problem is

$$\begin{array}{ll}\text{minimize} & f_1(x_1) + f_2(x_2) + \cdots + f_n(x_n) \\ \text{subject to} & x_1 + x_2 + \cdots + x_n = b\end{array}$$

- ▶ eliminate  $x_n = b - x_1 - \cdots - x_{n-1}$ , *i.e.*, choose

$$\hat{x} = be_n, \quad F = \begin{bmatrix} I \\ -\mathbf{1}^T \end{bmatrix} \in \mathbf{R}^{n \times (n-1)}$$

- ▶ reduced problem: minimize  $f_1(x_1) + \cdots + f_{n-1}(x_{n-1}) + f_n(b - x_1 - \cdots - x_{n-1})$

# Outline

Equality constrained minimization

Newton's method with equality constraints

Infeasible start Newton method

Implementation

## Newton step

- ▶ Newton step  $\Delta x_{\text{nt}}$  of  $f$  at feasible  $x$  is given by solution  $v$  of

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = \begin{bmatrix} -\nabla f(x) \\ 0 \end{bmatrix}$$

- ▶  $\Delta x_{\text{nt}}$  solves second order approximation (with variable  $v$ )

$$\begin{array}{ll} \text{minimize} & \widehat{f}(x+v) = f(x) + \nabla f(x)^T v + (1/2)v^T \nabla^2 f(x)v \\ \text{subject to} & A(x+v) = b \end{array}$$

- ▶  $\Delta x_{\text{nt}}$  equations follow from linearizing optimality conditions

$$\nabla f(x+v) + A^T w \approx \nabla f(x) + \nabla^2 f(x)v + A^T w = 0, \quad A(x+v) = b$$

## Newton decrement

- ▶ Newton decrement for equality constrained minimization is

$$\lambda(x) = \left( \Delta x_{\text{nt}}^T \nabla^2 f(x) \Delta x_{\text{nt}} \right)^{1/2} = \left( -\nabla f(x)^T \Delta x_{\text{nt}} \right)^{1/2}$$

- ▶ gives an estimate of  $f(x) - p^\star$  using quadratic approximation  $\widehat{f}$ :

$$f(x) - \inf_{Ay=b} \widehat{f}(y) = \lambda(x)^2 / 2$$

- ▶ directional derivative in Newton direction:

$$\left. \frac{d}{dt} f(x + t \Delta x_{\text{nt}}) \right|_{t=0} = -\lambda(x)^2$$

- ▶ in general,  $\lambda(x) \neq \left( \nabla f(x)^T \nabla^2 f(x)^{-1} \nabla f(x) \right)^{1/2}$

## Newton's method with equality constraints

---

**given** starting point  $x \in \text{dom } f$  with  $Ax = b$ , tolerance  $\epsilon > 0$ .

**repeat**

1. Compute the Newton step and decrement  $\Delta x_{\text{nt}}, \lambda(x)$ .
  2. *Stopping criterion.* **quit** if  $\lambda^2/2 \leq \epsilon$ .
  3. *Line search.* Choose step size  $t$  by backtracking line search.
  4. *Update.*  $x := x + t\Delta x_{\text{nt}}$ .
- 

- ▶ a feasible descent method:  $x^{(k)}$  feasible and  $f(x^{(k+1)}) < f(x^{(k)})$
- ▶ affine invariant



## Newton's method and elimination

- ▶ reduced problem: minimize  $\tilde{f}(z) = f(Fz + \hat{x})$ 
  - variables  $z \in \mathbf{R}^{n-p}$
  - $\hat{x}$  satisfies  $A\hat{x} = b$ ; **rank**  $F = n - p$  and  $AF = 0$
- ▶ (unconstrained) Newton's method for  $\tilde{f}$ , started at  $z^{(0)}$ , generates iterates  $z^{(k)}$
- ▶ iterates of Newton's method with equality constraints, started at  $x^{(0)} = Fz^{(0)} + \hat{x}$ , are

$$x^{(k)} = Fz^{(k)} + \hat{x}$$

- ▶ hence, don't need separate convergence analysis

# Outline

Equality constrained minimization

Newton's method with equality constraints

Infeasible start Newton method

Implementation

## Newton step at infeasible points

- ▶ with  $y = (x, v)$ , write optimality condition as  $r(y) = 0$ , where

$$r(y) = (\nabla f(x) + A^T v, Ax - b)$$

is **primal-dual residual**

- ▶ consider  $x \in \text{dom} f$ ,  $Ax \neq b$ , i.e.,  $x$  is infeasible
- ▶ linearizing  $r(y) = 0$  gives  $r(y + \Delta y) \approx r(y) + Dr(y)\Delta y = 0$ :

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x_{\text{nt}} \\ \Delta v_{\text{nt}} \end{bmatrix} = - \begin{bmatrix} \nabla f(x) + A^T v \\ Ax - b \end{bmatrix}$$

- ▶  $(\Delta x_{\text{nt}}, \Delta v_{\text{nt}})$  is called **infeasible** or **primal-dual** Newton step at  $x$

## Infeasible start Newton method

---

**given** starting point  $x \in \text{dom} f$ ,  $v$ , tolerance  $\epsilon > 0$ ,  $\alpha \in (0, 1/2)$ ,  $\beta \in (0, 1)$ .

**repeat**

1. Compute primal and dual Newton steps  $\Delta x_{\text{nt}}$ ,  $\Delta v_{\text{nt}}$ .

2. *Backtracking line search* on  $\|r\|_2$ .

$t := 1$ .

**while**  $\|r(x + t\Delta x_{\text{nt}}, v + t\Delta v_{\text{nt}})\|_2 > (1 - \alpha t)\|r(x, v)\|_2$ ,  $t := \beta t$ .

3. *Update*.  $x := x + t\Delta x_{\text{nt}}$ ,  $v := v + t\Delta v_{\text{nt}}$ .

**until**  $Ax = b$  and  $\|r(x, v)\|_2 \leq \epsilon$ .

---

- ▶ not a descent method:  $f(x^{(k+1)}) > f(x^{(k)})$  is possible
- ▶ directional derivative of  $\|r(y)\|_2$  in direction  $\Delta y = (\Delta x_{\text{nt}}, \Delta v_{\text{nt}})$  is

$$\left. \frac{d}{dt} \|r(y + t\Delta y)\|_2 \right|_{t=0} = -\|r(y)\|_2$$

# Outline

Equality constrained minimization

Newton's method with equality constraints

Infeasible start Newton method

Implementation

## Solving KKT systems

- ▶ feasible and infeasible Newton methods require solving KKT system

$$\begin{bmatrix} H & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = - \begin{bmatrix} g \\ h \end{bmatrix}$$

- ▶ in general, can use  $LDL^T$  factorization
- ▶ or elimination (if  $H$  nonsingular and easily inverted):
  - solve  $AH^{-1}A^Tw = h - AH^{-1}g$  for  $w$
  - $v = -H^{-1}(g + A^Tw)$

## Example: Equality constrained analytic centering


- ▶ **primal problem:** minimize  $-\sum_{i=1}^n \log x_i$  subject to  $Ax = b$
- ▶ **dual problem:** maximize  $-b^T \nu + \sum_{i=1}^n \log(A^T \nu)_i + n$ 
  - recover  $x^\star$  as  $x_i^\star = 1/(A^T \nu)_i$
- ▶ three methods to solve:
  - Newton method with equality constraints
  - Newton method applied to dual problem
  - infeasible start Newton method

these have **different requirements for initialization**

- ▶ we'll look at an example with  $A \in \mathbf{R}^{100 \times 500}$ , different starting points

## Newton's method with equality constraints


- requires  $x^{(0)} \succ 0$ ,  $Ax^{(0)} = b$






## Newton method applied to dual problem

- requires  $A^T \nu^{(0)} \succ 0$



## Infeasible start Newton method

- requires  $x^{(0)} \succ 0$



## Complexity per iteration of three methods is identical

- ▶ for feasible Newton method, use block elimination to solve KKT system

$$\begin{bmatrix} \mathbf{diag}(x)^{-2} & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ w \end{bmatrix} = \begin{bmatrix} \mathbf{diag}(x)^{-1} \mathbf{1} \\ 0 \end{bmatrix}$$

reduces to solving  $A \mathbf{diag}(x)^2 A^T w = b$

- ▶ for Newton system applied to dual, solve  $A \mathbf{diag}(A^T v)^{-2} A^T \Delta v = -b + A \mathbf{diag}(A^T v)^{-1} \mathbf{1}$
- ▶ for infeasible start Newton method, use block elimination to solve KKT system

$$\begin{bmatrix} \mathbf{diag}(x)^{-2} & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta v \end{bmatrix} = \begin{bmatrix} \mathbf{diag}(x)^{-1} \mathbf{1} - A^T v \\ b - Ax \end{bmatrix}$$

reduces to solving  $A \mathbf{diag}(x)^2 A^T w = 2Ax - b$

- ▶ conclusion: in each case, solve  $ADA^T w = h$  with  $D$  positive diagonal

## Example: Network flow optimization

- ▶ directed graph with  $n$  arcs,  $p + 1$  nodes
- ▶  $x_i$ : flow through arc  $i$ ;  $\phi_i$ : strictly convex flow cost function for arc  $i$
- ▶ **incidence matrix**  $\tilde{A} \in \mathbf{R}^{(p+1) \times n}$  defined as

$$\tilde{A}_{ij} = \begin{cases} 1 & \text{arc } j \text{ leaves node } i \\ -1 & \text{arc } j \text{ enters node } i \\ 0 & \text{otherwise} \end{cases}$$

- ▶ **reduced incidence matrix**  $A \in \mathbf{R}^{p \times n}$  is  $\tilde{A}$  with last row removed
- ▶ **rank**  $A = p$  if graph is connected
- ▶ flow conservation is  $Ax = b$ ,  $b \in \mathbf{R}^p$  is (reduced) source vector
- ▶ **network flow optimization problem**: minimize  $\sum_{i=1}^n \phi_i(x_i)$  subject to  $Ax = b$

## KKT system

- ▶ KKT system is

$$\begin{bmatrix} H & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = - \begin{bmatrix} g \\ h \end{bmatrix}$$

- ▶  $H = \mathbf{diag}(\phi_1''(x_1), \dots, \phi_n''(x_n))$ , positive diagonal
- ▶ solve via elimination:

$$AH^{-1}A^T w = h - AH^{-1}g, \quad v = -H^{-1}(g + A^T w)$$

- ▶ sparsity pattern of  $AH^{-1}A^T$  is given by graph connectivity

$$\begin{aligned} (AH^{-1}A^T)_{ij} \neq 0 &\iff (AA^T)_{ij} \neq 0 \\ &\iff \text{nodes } i \text{ and } j \text{ are connected by an arc} \end{aligned}$$

## Analytic center of linear matrix inequality

- ▶ minimize  $-\log \det X$  subject to  $\mathbf{tr}(A_i X) = b_i, \quad i = 1, \dots, p$
- ▶ optimality conditions

$$X^\star \succ 0, \quad -(X^\star)^{-1} + \sum_{j=1}^p v_j^\star A_j = 0, \quad \mathbf{tr}(A_i X^\star) = b_i, \quad i = 1, \dots, p$$

- ▶ Newton step  $\Delta X$  at feasible  $X$  is defined by

$$X^{-1}(\Delta X)X^{-1} + \sum_{j=1}^p w_j A_j = X^{-1}, \quad \mathbf{tr}(A_i \Delta X) = 0, \quad i = 1, \dots, p$$

- ▶ follows from linear approximation  $(X + \Delta X)^{-1} \approx X^{-1} - X^{-1}(\Delta X)X^{-1}$
- ▶  $n(n+1)/2 + p$  variables  $\Delta X, w$

## Solution by block elimination

- ▶ eliminate  $\Delta X$  from first equation to get  $\Delta X = X - \sum_{j=1}^p w_j X A_j X$
- ▶ substitute  $\Delta X$  in second equation to get

$$\sum_{j=1}^p \text{tr}(A_i X A_j X) w_j = b_i, \quad i = 1, \dots, p$$

- ▶ a dense positive definite set of linear equations with variable  $w \in \mathbf{R}^p$
- ▶ form and solve this set of equations to get  $w$ , then get  $\Delta X$  from equation above

## Flop count

- ▶ find Cholesky factor  $L$  of  $X$   $(1/3)n^3$
- ▶ form  $p$  products  $L^T A_j L$   $(3/2)pn^3$
- ▶ form  $p(p+1)/2$  inner products  $\text{tr}((L^T A_i L)(L^T A_j L))$  to get coefficient matrix  $(1/2)p^2 n^2$
- ▶ solve  $p \times p$  system of equations via Cholesky factorization  $(1/3)p^3$
- ▶ flop count dominated by  $pn^3 + p^2 n^2$
- ▶ cf. naïve method,  $(n^2 + p)^3$



## 11. Interior-point methods

# Outline

Inequality constrained minimization

Logarithmic barrier and central path

Barrier method

Phase I methods

Complexity analysis

Generalized inequalities

## Inequality constrained minimization

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

we assume

- ▶  $f_i$  convex, twice continuously differentiable
- ▶  $A \in \mathbf{R}^{p \times n}$  with **rank**  $A = p$
- ▶  $p^\star$  is finite and attained
- ▶ problem is strictly feasible: there exists  $\tilde{x}$  with

$$\tilde{x} \in \mathbf{dom} f_0, \quad f_i(\tilde{x}) < 0, \quad i = 1, \dots, m, \quad A\tilde{x} = b$$

hence, strong duality holds and dual optimum is attained

## Examples

- ▶ LP, QP, QCQP, GP
- ▶ entropy maximization with linear inequality constraints

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^n x_i \log x_i \\ \text{subject to} & Fx \leq g, \quad Ax = b\end{array}$$

with  $\text{dom } f_0 = \mathbf{R}_{++}^n$

- ▶ differentiability may require reformulating the problem, e.g., piecewise-linear minimization or  $\ell_\infty$ -norm approximation via LP
- ▶ SDPs and SOCPs are better handled as problems with generalized inequalities (see later)

# Outline

Inequality constrained minimization

Logarithmic barrier and central path

Barrier method

Phase I methods

Complexity analysis

Generalized inequalities

## Logarithmic barrier

- reformulation via **indicator function**:

$$\begin{array}{ll}\text{minimize} & f_0(x) + \sum_{i=1}^m I_-(f_i(x)) \\ \text{subject to} & Ax = b\end{array}$$


where  $I_-(u) = 0$  if  $u \leq 0$ ,  $I_-(u) = \infty$  otherwise

- **approximation via logarithmic barrier**:

$$\begin{array}{ll}\text{minimize} & f_0(x) - (1/t) \sum_{i=1}^m \log(-f_i(x)) \\ \text{subject to} & Ax = b\end{array}$$

- an equality constrained problem
- for  $t > 0$ ,  $-(1/t) \log(-u)$  is a smooth approximation of  $I_-$
- approximation improves as  $t \rightarrow \infty$

- $-(1/t) \log u$  for three values of  $t$ , and  $I_-(u)$



## Logarithmic barrier function

- ▶ log barrier function for constraints  $f_1(x) \leq 0, \dots, f_m(x) \leq 0$

$$\phi(x) = - \sum_{i=1}^m \log(-f_i(x)), \quad \text{dom } \phi = \{x \mid f_1(x) < 0, \dots, f_m(x) < 0\}$$

- ▶ convex (from composition rules)
- ▶ twice continuously differentiable, with derivatives

$$\begin{aligned}\nabla \phi(x) &= \sum_{i=1}^m \frac{1}{-f_i(x)} \nabla f_i(x) \\ \nabla^2 \phi(x) &= \sum_{i=1}^m \frac{1}{f_i(x)^2} \nabla f_i(x) \nabla f_i(x)^T + \sum_{i=1}^m \frac{1}{-f_i(x)} \nabla^2 f_i(x)\end{aligned}$$



## Central path

- ▶ for  $t > 0$ , define  $x^\star(t)$  as the solution of

$$\begin{array}{ll}\text{minimize} & tf_0(x) + \phi(x) \\ \text{subject to} & Ax = b\end{array}$$


(for now, assume  $x^\star(t)$  exists and is unique for each  $t > 0$ )

- ▶ central path is  $\{x^\star(t) \mid t > 0\}$

**example:** central path for an LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & a_i^T x \leq b_i, \quad i = 1, \dots, 6\end{array}$$

hyperplane  $c^T x = c^T x^\star(t)$  is tangent to level curve of  $\phi$  through  $x^\star(t)$



## Dual points on central path

- ▶  $x = x^\star(t)$  if there exists a  $w$  such that

$$t\nabla f_0(x) + \sum_{i=1}^m \frac{1}{-f_i(x)} \nabla f_i(x) + A^T w = 0, \quad Ax = b$$

- ▶ therefore,  $x^\star(t)$  minimizes the Lagrangian

$$L(x, \lambda^\star(t), \nu^\star(t)) = f_0(x) + \sum_{i=1}^m \lambda_i^\star(t) f_i(x) + \nu^\star(t)^T (Ax - b)$$

where we define  $\lambda_i^\star(t) = 1/(-tf_i(x^\star(t)))$  and  $\nu^\star(t) = w/t$

- ▶ this confirms the intuitive idea that  $f_0(x^\star(t)) \rightarrow p^\star$  if  $t \rightarrow \infty$ :

$$p^\star \geq g(\lambda^\star(t), \nu^\star(t)) = L(x^\star(t), \lambda^\star(t), \nu^\star(t)) = f_0(x^\star(t)) - m/t$$

## Interpretation via KKT conditions

$x = x^\star(t)$ ,  $\lambda = \lambda^\star(t)$ ,  $\nu = \nu^\star(t)$  satisfy

1. primal constraints:  $f_i(x) \leq 0$ ,  $i = 1, \dots, m$ ,  $Ax = b$
2. dual constraints:  $\lambda \geq 0$
3. approximate complementary slackness:  $-\lambda_i f_i(x) = 1/t$ ,  $i = 1, \dots, m$
4. gradient of Lagrangian with respect to  $x$  vanishes:

$$\nabla f_0(x) + \sum_{i=1}^m \lambda_i \nabla f_i(x) + A^T \nu = 0$$

difference with KKT is that condition 3 replaces  $\lambda_i f_i(x) = 0$

## Force field interpretation

- ▶ **centering problem** (for problem with no equality constraints)

$$\text{minimize} \quad tf_0(x) - \sum_{i=1}^m \log(-f_i(x))$$

- ▶ **force field interpretation**

- $tf_0(x)$  is potential of force field  $F_0(x) = -t\nabla f_0(x)$
- $-\log(-f_i(x))$  is potential of force field  $F_i(x) = (1/f_i(x))\nabla f_i(x)$

- ▶ forces balance at  $x^\star(t)$ :



$$F_0(x^\star(t)) + \sum_{i=1}^m F_i(x^\star(t)) = 0$$

## Example: LP

- ▶ minimize  $c^T x$  subject to  $a_i^T x \leq b_i, i = 1, \dots, m$ , with  $x \in \mathbf{R}^n$
- ▶ objective force field is constant:  $F_0(x) = -tc$
- ▶ constraint force field decays as inverse distance to constraint hyperplane:

$$F_i(x) = \frac{-a_i}{b_i - a_i^T x}, \quad \|F_i(x)\|_2 = \frac{1}{\text{dist}(x, \mathcal{H}_i)}$$

where  $\mathcal{H}_i = \{x \mid a_i^T x = b_i\}$



# Outline

Inequality constrained minimization

Logarithmic barrier and central path

**Barrier method**

Phase I methods

Complexity analysis

Generalized inequalities

## Barrier method

---

**given** strictly feasible  $x$ ,  $t := t^{(0)} > 0$ ,  $\mu > 1$ , tolerance  $\epsilon > 0$ .



**repeat**

1. *Centering step.* Compute  $x^\star(t)$  by minimizing  $tf_0 + \phi$ , subject to  $Ax = b$ .
  2. *Update.*  $x := x^\star(t)$ .
  3. *Stopping criterion.* **quit** if  $m/t < \epsilon$ .
  4. *Increase  $t$ .*  $t := \mu t$ .
- 

- ▶ terminates with  $f_0(x) - p^\star \leq \epsilon$  (stopping criterion follows from  $f_0(x^\star(t)) - p^\star \leq m/t$ )
- ▶ centering usually done using Newton's method, starting at current  $x$
- ▶ choice of  $\mu$  involves a trade-off: large  $\mu$  means fewer outer iterations, more inner (Newton) iterations; typical values:  $\mu = 10$  or  $20$
- ▶ several heuristics for choice of  $t^{(0)}$

## Example: Inequality form LP

( $m = 100$  inequalities,  $n = 50$  variables)




- ▶ starts with  $x$  on central path ( $t^{(0)} = 1$ , duality gap 100)
- ▶ terminates when  $t = 10^8$  (gap  $10^{-6}$ )
- ▶ total number of Newton iterations not very sensitive for  $\mu \geq 10$



## Example: Geometric program in convex form

( $m = 100$  inequalities and  $n = 50$  variables)

$$\begin{array}{ll}\text{minimize} & \log \left( \sum_{k=1}^5 \exp(a_{0k}^T x + b_{0k}) \right) \\ \text{subject to} & \log \left( \sum_{k=1}^5 \exp(a_{ik}^T x + b_{ik}) \right) \leq 0, \quad i = 1, \dots, m\end{array}$$




## Family of standard LPs

$$(A \in \mathbf{R}^{m \times 2m})$$

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b, \quad x \geq 0\end{array}$$

$m = 10, \dots, 1000$ ; for each  $m$ , solve 100 randomly generated instances



number of iterations grows very slowly as  $m$  ranges over a 100 : 1 ratio

# Outline

Inequality constrained minimization

Logarithmic barrier and central path

Barrier method

**Phase I methods**

Complexity analysis

Generalized inequalities

## Phase I methods

- ▶ barrier method needs strictly feasible starting point, *i.e.*,  $x$  with

$$f_i(x) < 0, \quad i = 1, \dots, m, \quad Ax = b$$

- ▶ (like the infeasible start Newton method, more sophisticated interior-point methods do not require a feasible starting point)
- ▶ **phase I** method forms an optimization problem that
  - is itself strictly feasible
  - finds a strictly feasible point for original problem, if one exists
  - certifies original problem as infeasible otherwise
- ▶ **phase II** uses barrier method starting from strictly feasible point found in phase I

## Basic phase I method

- ▶ introduce slack variable  $s$  in **phase I problem**

$$\begin{array}{ll}\text{minimize (over } x, s) & s \\ \text{subject to} & f_i(x) \leq s, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

with optimal value  $\bar{p}^\star$

- if  $\bar{p}^\star < 0$ , original inequalities are strictly feasible
  - if  $\bar{p}^\star > 0$ , original inequalities are infeasible
  - $\bar{p}^\star = 0$  is an ambiguous case
- ▶ start phase I problem with
    - any  $\tilde{x}$  in problem domain with  $A\tilde{x} = b$
    - $s = 1 + \max_i f_i(\tilde{x})$

## Sum of infeasibilities phase I method



- ▶ minimize **sum** of slacks, not max:

$$\begin{array}{ll}\text{minimize} & \mathbf{1}^T s \\ \text{subject to} & s \geq 0, \quad f_i(x) \leq s_i, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

- ▶ will find a strictly feasible point if one exists
- ▶ for infeasible problems, produces a solution that satisfies many (but not all) inequalities
- ▶ can weight slacks to set **priorities** (in satisfying constraints)




## Example

- ▶ infeasible set of 100 linear inequalities in 50 variables
- ▶ left: basic phase I solution; satisfies 39 inequalities
- ▶ right: sum of infeasibilities phase I solution; satisfies 79 inequalities



## Example: Family of linear inequalities

- ▶  $Ax \leq b + \gamma \Delta b$ ; strictly feasible for  $\gamma > 0$ , infeasible for  $\gamma < 0$
- ▶ use basic phase I, terminate when  $s < 0$  or dual objective is positive
- ▶ number of iterations roughly proportional to  $\log(1/|\gamma|)$





# Outline

Inequality constrained minimization

Logarithmic barrier and central path

Barrier method

Phase I methods

Complexity analysis

Generalized inequalities

## Number of outer iterations

- ▶ in each iteration duality gap is reduced by exactly the factor  $\mu$
- ▶ **number of outer (centering) iterations** is exactly

$$\left\lceil \frac{\log(m/(\epsilon t^{(0)}))}{\log \mu} \right\rceil$$

plus the initial centering step (to compute  $x^\star(t^{(0)})$ )

- ▶ we will bound **number of Newton steps per centering iteration** using self-concordance analysis

## Complexity analysis via self-concordance

same assumptions as on slide 11.2, plus:

- ▶ sublevel sets (of  $f_0$ , on the feasible set) are bounded
- ▶  $tf_0 + \phi$  is self-concordant with closed sublevel sets

second condition

- ▶ holds for LP, QP, QCQP
- ▶ may require reformulating the problem, *e.g.*,

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n x_i \log x_i \\ \text{subject to} & Fx \leq g \end{array} \quad \longrightarrow \quad \begin{array}{ll} \text{minimize} & \sum_{i=1}^n x_i \log x_i \\ \text{subject to} & Fx \leq g, \quad x \geq 0 \end{array}$$

- ▶ needed for complexity analysis; barrier method works even when self-concordance assumption does not apply

## Newton iterations per centering step

- ▶ we compute  $x^+ = x^\star(\mu t)$ , by minimizing  $\mu t f_0(x) + \phi(x)$  starting from  $x = x^\star(t)$
- ▶ from self-concordance theory,

$$\text{\#Newton iterations} \leq \frac{\mu t f_0(x) + \phi(x) - \mu t f_0(x^+) - \phi(x^+)}{\gamma} + c$$

- ▶  $\gamma, c$  are constants (that depend only on Newton algorithm parameters)
- ▶ we will bound numerator  $\mu t f_0(x) + \phi(x) - \mu t f_0(x^+) - \phi(x^+)$
- ▶ with  $\lambda_i = \lambda_i^\star(t) = -1/(t f_i(x))$ , we have  $-f_i(x) = 1/(t \lambda_i)$ , so

$$\phi(x) = \sum_{i=1}^m -\log(-f_i(x)) = \sum_{i=1}^m \log(t \lambda_i)$$

so

$$\phi(x) - \phi(x^+) = \sum_{i=1}^m (\log(t \lambda_i) + \log(-f_i(x^+))) = \sum_{i=1}^m \log(-\mu t \lambda_i f_i(x^+)) - m \log \mu$$


using  $\log u \leq u - 1$  we have  $\phi(x) - \phi(x^+) \leq -\mu t \sum_{i=1}^m \lambda_i f_i(x^+) - m - m \log \mu$ , so

$$\begin{aligned}
 & \mu t f_0(x) + \phi(x) - \mu t f_0(x^+) - \phi(x^+) \\
 & \leq \mu t f_0(x) - \mu t f_0(x^+) - \mu t \sum_{i=1}^m \lambda_i f_i(x^+) - m - m \log \mu \\
 & = \mu t f_0(x) - \mu t \left( f_0(x^+) + \sum_{i=1}^m \lambda_i f_i(x^+) + v^T (Ax^+ - b) \right) - m - m \log \mu \\
 & = \mu t f_0(x) - \mu t L(x^+, \lambda, v) - m - m \log \mu \\
 & \leq \mu t f_0(x) - \mu t g(\lambda, v) - m - m \log \mu \\
 & = m(\mu - 1 - \log \mu)
 \end{aligned}$$

using  $L(x^+, \lambda, \nu) \geq g(\lambda, \nu)$  in second last line and  $f_0(x) - g(\lambda, \nu) = m/t$  in last line

## Total number of Newton iterations

$$\text{\#Newton iterations} \leq N = \left\lceil \frac{\log(m/(t^{(0)}\epsilon))}{\log \mu} \right\rceil \left( \frac{m(\mu - 1 - \log \mu)}{\gamma} + c \right)$$



$N$  versus  $\mu$  for typical values of  $\gamma, c$ ;  
 $m = 100$ , initial duality gap  $\frac{m}{t^{(0)}\epsilon} = 10^5$

- confirms trade-off in choice of  $\mu$
- in practice, #iterations is in the tens; not very sensitive for  $\mu \geq 10$

## Polynomial-time complexity of barrier method

- ▶ for  $\mu = 1 + 1/\sqrt{m}$ :

$$N = O\left(\sqrt{m} \log\left(\frac{m/t^{(0)}}{\epsilon}\right)\right)$$

- ▶ number of Newton iterations for fixed gap reduction is  $O(\sqrt{m})$
- ▶ multiply with cost of one Newton iteration (a polynomial function of problem dimensions), to get bound on number of flops
- ▶ this choice of  $\mu$  optimizes worst-case complexity; in practice we choose  $\mu$  fixed and larger

# Outline

Inequality constrained minimization

Logarithmic barrier and central path

Barrier method

Phase I methods

Complexity analysis

Generalized inequalities



## Generalized inequalities

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \preceq_{K_i} 0, \quad i = 1, \dots, m \\ & Ax = b\end{array}$$

- ▶  $f_0$  convex,  $f_i : \mathbf{R}^n \rightarrow \mathbf{R}^{k_i}$ ,  $i = 1, \dots, m$ , convex with respect to proper cones  $K_i \in \mathbf{R}^{k_i}$
- ▶ we assume
  - $f_i$  twice continuously differentiable
  - $A \in \mathbf{R}^{p \times n}$  with **rank**  $A = p$
  - $p^\star$  is finite and attained
  - problem is strictly feasible; hence strong duality holds and dual optimum is attained
- ▶ examples of greatest interest: SOCP, SDP

## Generalized logarithm for proper cone

$\psi : \mathbf{R}^q \rightarrow \mathbf{R}$  is **generalized logarithm** for proper cone  $K \subseteq \mathbf{R}^q$  if:

- ▶ **dom**  $\psi = \mathbf{int} K$  and  $\nabla^2 \psi(y) \prec 0$  for  $y \succ_K 0$
- ▶  $\psi(sy) = \psi(y) + \theta \log s$  for  $y \succ_K 0, s > 0$  ( $\theta$  is the degree of  $\psi$ )

### examples

- ▶ nonnegative orthant  $K = \mathbf{R}_+^n$ :  $\psi(y) = \sum_{i=1}^n \log y_i$ , with degree  $\theta = n$
- ▶ positive semidefinite cone  $K = \mathbf{S}_+^n$ :  $\psi(Y) = \log \det Y$ , with degree  $\theta = n$
- ▶ second-order cone  $K = \{y \in \mathbf{R}^{n+1} \mid (y_1^2 + \cdots + y_n^2)^{1/2} \leq y_{n+1}\}$ :

$$\psi(y) = \log(y_{n+1}^2 - y_1^2 - \cdots - y_n^2) \quad \text{with degree } (\theta = 2)$$

## Properties

- ▶ (without proof): for  $y \succ_K 0$ ,

$$\nabla\psi(y) \succeq_{K^*} 0, \quad y^T \nabla\psi(y) = \theta$$

- ▶ nonnegative orthant  $\mathbf{R}_+^n$ :  $\psi(y) = \sum_{i=1}^n \log y_i$

$$\nabla\psi(y) = (1/y_1, \dots, 1/y_n), \quad y^T \nabla\psi(y) = n$$

- ▶ positive semidefinite cone  $\mathbf{S}_+^n$ :  $\psi(Y) = \log \det Y$

$$\nabla\psi(Y) = Y^{-1}, \quad \text{tr}(Y \nabla\psi(Y)) = n$$

- ▶ second-order cone  $K = \{y \in \mathbf{R}^{n+1} \mid (y_1^2 + \dots + y_n^2)^{1/2} \leq y_{n+1}\}$ :

$$\nabla\psi(y) = \frac{2}{y_{n+1}^2 - y_1^2 - \dots - y_n^2} \begin{bmatrix} -y_1 \\ \vdots \\ -y_n \\ y_{n+1} \end{bmatrix}, \quad y^T \nabla\psi(y) = 2$$

## Logarithmic barrier and central path

**logarithmic barrier** for  $f_1(x) \leq_{K_1} 0, \dots, f_m(x) \leq_{K_m} 0$ :

$$\phi(x) = - \sum_{i=1}^m \psi_i(-f_i(x)), \quad \text{dom } \phi = \{x \mid f_i(x) \prec_{K_i} 0, i = 1, \dots, m\}$$

- ▶  $\psi_i$  is generalized logarithm for  $K_i$ , with degree  $\theta_i$
- ▶  $\phi$  is convex, twice continuously differentiable

**central path:**  $\{x^\star(t) \mid t > 0\}$  where  $x^\star(t)$  is solution of

$$\begin{array}{ll} \text{minimize} & tf_0(x) + \phi(x) \\ \text{subject to} & Ax = b \end{array}$$

## Dual points on central path

$x = x^\star(t)$  if there exists  $w \in \mathbf{R}^p$ ,

$$t \nabla f_0(x) + \sum_{i=1}^m Df_i(x)^T \nabla \psi_i(-f_i(x)) + A^T w = 0$$

$(Df_i(x) \in \mathbf{R}^{k_i \times n}$  is derivative matrix of  $f_i$ )

► therefore,  $x^\star(t)$  minimizes Lagrangian  $L(x, \lambda^\star(t), \nu^\star(t))$ , where

$$\lambda_i^\star(t) = \frac{1}{t} \nabla \psi_i(-f_i(x^\star(t))), \quad \nu^\star(t) = \frac{w}{t}$$

► from properties of  $\psi_i$ :  $\lambda_i^\star(t) \succ_{K_i^\star} 0$ , with duality gap

$$f_0(x^\star(t)) - g(\lambda^\star(t), \nu^\star(t)) = (1/t) \sum_{i=1}^m \theta_i$$

## Example: Semidefinite programming

(with  $F_i \in \mathbf{S}^p$ )

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & F(x) = \sum_{i=1}^n x_i F_i + G \leq 0\end{array}$$

- ▶ logarithmic barrier:  $\phi(x) = \log \det(-F(x))^{-1}$
- ▶ central path:  $x^\star(t)$  minimizes  $tc^T x - \log \det(-F(x))$ ; hence

$$tc_i - \text{tr}(F_i F(x^\star(t))^{-1}) = 0, \quad i = 1, \dots, n$$

- ▶ dual point on central path:  $Z^\star(t) = -(1/t)F(x^\star(t))^{-1}$  is feasible for

$$\begin{array}{ll}\text{maximize} & \text{tr}(GZ) \\ \text{subject to} & \text{tr}(F_i Z) + c_i = 0, \quad i = 1, \dots, n \\ & Z \geq 0\end{array}$$

- ▶ duality gap on central path:  $c^T x^\star(t) - \text{tr}(GZ^\star(t)) = p/t$

## Barrier method

---

**given** strictly feasible  $x$ ,  $t := t^{(0)} > 0$ ,  $\mu > 1$ , tolerance  $\epsilon > 0$ .

**repeat**

1. *Centering step.* Compute  $x^\star(t)$  by minimizing  $tf_0 + \phi$ , subject to  $Ax = b$ .
  2. *Update.*  $x := x^\star(t)$ .
  3. *Stopping criterion.* **quit** if  $(\sum_i \theta_i)/t < \epsilon$ .
  4. *Increase  $t$ .*  $t := \mu t$ .
- 



- ▶ only difference is duality gap  $m/t$  on central path is replaced by  $\sum_i \theta_i/t$
- ▶ number of outer iterations:

$$\left\lceil \frac{\log((\sum_i \theta_i)/(\epsilon t^{(0)}))}{\log \mu} \right\rceil$$

- ▶ complexity analysis via self-concordance applies to SDP, SOCP

## Example: SOCP



(50 variables, 50 SOC constraints in  $\mathbf{R}^6$ )





## Example: SDP

(100 variables, LMI constraint in  $\mathbf{S}^{100}$ )




## Example: Family of SDPs

$$(A \in \mathbf{S}^n, x \in \mathbf{R}^n)$$

$$\begin{array}{ll}\text{minimize} & \mathbf{1}^T x \\ \text{subject to} & A + \mathbf{diag}(x) \geq 0\end{array}$$

$n = 10, \dots, 1000$ ; for each  $n$  solve 100 randomly generated instances



## Primal-dual interior-point methods

- ▶ more efficient than barrier method when high accuracy is needed
- ▶ update primal and dual variables, and  $\kappa$ , at each iteration; no distinction between inner and outer iterations
- ▶ often exhibit superlinear asymptotic convergence
- ▶ search directions can be interpreted as Newton directions for modified KKT conditions
- ▶ can start at infeasible points
- ▶ cost per iteration same as barrier method

## 12. Conclusions

# Modeling

## mathematical optimization

- ▶ problems in engineering design, data analysis and statistics, economics, management, . . . , can often be expressed as mathematical optimization problems
- ▶ techniques exist to take into account multiple objectives or uncertainty in the data

## tractability

- ▶ roughly speaking, tractability in optimization requires convexity
- ▶ algorithms for nonconvex optimization find local (suboptimal) solutions, or are very expensive
- ▶ surprisingly many applications can be formulated as convex problems

# Theoretical consequences of convexity

- ▶ local optima are global
- ▶ extensive duality theory
  - systematic way of deriving lower bounds on optimal value
  - necessary and sufficient optimality conditions
  - certificates of infeasibility
  - sensitivity analysis
- ▶ solution methods with polynomial worst-case complexity theory (with self-concordance)

## Practical consequences of convexity

(most) **convex problems can be solved globally and efficiently**

- ▶ interior-point methods require 20 – 80 steps in practice
- ▶ basic algorithms (*e.g.*, Newton, barrier method, ...) are easy to implement and work well for small and medium size problems (larger problems if structure is exploited)
- ▶ high-quality solvers (some open-source) are available
- ▶ high level modeling tools like CVXPY ease modeling and problem specification

## How to use convex optimization

to use convex optimization in some applied context

- ▶ use rapid prototyping, approximate modeling
  - start with simple models, small problem instances, inefficient solution methods
  - if you don't like the results, no need to expend further effort on more accurate models or efficient algorithms
- ▶ work out, simplify, and interpret optimality conditions and dual
- ▶ even if the problem is quite nonconvex, you can use convex optimization
  - in subproblems, *e.g.*, to find search direction
  - by repeatedly forming and solving a convex approximation at the current point



## Further topics

some topics we didn't cover:

- ▶ methods for very large scale problems
- ▶ subgradient calculus, convex analysis
- ▶ localization, subgradient, proximal and related methods
- ▶ distributed convex optimization
- ▶ applications that build on or use convex optimization

these are all covered in EE364b.

## Related classes

- ▶ EE364b — convex optimization II (Pilanci)
- ▶ EE364m — mathematics of convexity (Duchi)
- ▶ CS261, CME334, MSE213 — theory and algorithm analysis (Sidford)
- ▶ AA222 — algorithms for nonconvex optimization (Kochenderfer)
- ▶ CME307 — linear and conic optimization (Ye)