

# The Boundary of the Mandelbrot set for a pair of linear maps

Stefano Silvestri

IUPUI

## Setting: Iterated Function Systems

For each  $\lambda \in \mathbb{D} \setminus \{0\}$  associate to it the compact sets

$$A_\lambda := \left\{ \sum_{j=0}^{\infty} a_j \lambda^j \mid a_j \in \{-1, +1\} \right\}$$

$$\tilde{A}_\lambda := \left\{ \sum_{j=0}^{\infty} a_j \lambda^j \mid a_j \in \{-1, 0, +1\} \right\}$$

These are, respectively, the *attractors* of the IFS  $\{\mathfrak{s}_-, \mathfrak{s}_+\}$  and  $\{\mathfrak{s}_-, \mathfrak{s}_0, \mathfrak{s}_+\}$  where

$$\mathfrak{s}_-(z) = \lambda z - 1 \quad \mathfrak{s}_0(z) = \lambda z \quad \mathfrak{s}_+(z) = \lambda z + 1$$

# Setting: Iterated Function Systems

For each  $\lambda \in \mathbb{D} \setminus \{0\}$  associate to it the compact sets

$$A_\lambda := \left\{ \sum_{j=0}^{\infty} a_j \lambda^j \mid a_j \in \{-1, +1\} \right\}$$

$$\tilde{A}_\lambda := \left\{ \sum_{j=0}^{\infty} a_j \lambda^j \mid a_j \in \{-1, 0, +1\} \right\}$$


These are, respectively, the *attractors* of the IFS  $\{\mathfrak{s}_-, \mathfrak{s}_+\}$  and  $\{\mathfrak{s}_-, \mathfrak{s}_0, \mathfrak{s}_+\}$  where

$$\mathfrak{s}_-(z) = \lambda z - 1 \quad \mathfrak{s}_0(z) = \lambda z \quad \mathfrak{s}_+(z) = \lambda z + 1$$


In other words, they are the unique non-empty compact sets satisfying

$$A_\lambda = \mathfrak{s}_-(A_\lambda) \cup \mathfrak{s}_+(A_\lambda) \quad \tilde{A}_\lambda = \mathfrak{s}_-(\tilde{A}_\lambda) \cup \mathfrak{s}_0(\tilde{A}_\lambda) \cup \mathfrak{s}_+(\tilde{A}_\lambda)$$


# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda \approx -0.366 + 0.52i$




# Constructing $A_\lambda$ for $\lambda = 0.25 + 0.5i$




# Constructing $A_\lambda$ for $\lambda = 0.25 + 0.5i$




# Constructing $A_\lambda$ for $\lambda = 0.25 + 0.5i$



# Constructing $A_\lambda$ for $\lambda = 0.25 + 0.5i$




# Constructing $A_\lambda$ for $\lambda = 0.25 + 0.5i$




# The Mandelbrot set for $\{\mathfrak{s}_-, \mathfrak{s}_+\}$

$$A_\lambda = \mathfrak{s}_-(A_\lambda) \cup \mathfrak{s}_+(A_\lambda)$$



# The Mandelbrot set for $\{\mathfrak{s}_-, \mathfrak{s}_+\}$


$$A_\lambda = \mathfrak{s}_-(A_\lambda) \cup \mathfrak{s}_+(A_\lambda)$$



$$\mathcal{M} = \{\lambda \in \mathbb{D} \mid A_\lambda \text{ is connected}\}$$

# The Mandelbrot set for $\{\mathfrak{s}_-, \mathfrak{s}_+\}$

$$A_\lambda = \mathfrak{s}_-(A_\lambda) \cup \mathfrak{s}_+(A_\lambda)$$




$$\mathcal{M} = \{\lambda \in \mathbb{D} \mid A_\lambda \text{ is connected}\}$$

$$\mathcal{M}_0 = \{\lambda \in \mathbb{D} \mid 0 \in A_\lambda\}$$


# The Mandelbrot set for $\{\mathfrak{s}_-, \mathfrak{s}_+\}$

$$A_\lambda = \mathfrak{s}_-(A_\lambda) \cup \mathfrak{s}_+(A_\lambda) \quad \tilde{A}_\lambda = \mathfrak{s}_-(\tilde{A}_\lambda) \cup \mathfrak{s}_0(\tilde{A}_\lambda) \cup \mathfrak{s}_+(\tilde{A}_\lambda)$$




$$\begin{aligned} \mathcal{M} &= \{\lambda \in \mathbb{D} \mid A_\lambda \text{ is connected}\} = \left\{ \lambda \in \mathbb{D} \mid 0 \in \tilde{A}_\lambda \right\} \\ \mathcal{M}_0 &= \{\lambda \in \mathbb{D} \mid 0 \in A_\lambda\} \end{aligned}$$





$\mathcal{M}$  (in blue)



$$\mathcal{M}_0 \subset \mathcal{M}$$



# Interesting features of $\mathcal{M}$ and $\mathcal{M}_0$



# History

1985 Barnsley & Harrington introduce  $\mathcal{M}$ .

# History

1985 Barnsley & Harrington introduce  $\mathcal{M}$ .

1988 – 1992 Bousch proved both  $\mathcal{M}$  and  $\mathcal{M}_0$  are connected and locally connected.

# History

1985 Barnsley & Harrington introduce  $\mathcal{M}$ .

1988 – 1992 Bousch proved both  $\mathcal{M}$  and  $\mathcal{M}_0$  are connected and locally connected.

2002 Bandt proved  $\mathbb{D} \setminus \mathcal{M}$  is disconnected.

# History

1985 Barnsley & Harrington introduce  $\mathcal{M}$ .

1988 – 1992 Bousch proved both  $\mathcal{M}$  and  $\mathcal{M}_0$  are connected and locally connected.

2002 Bandt proved  $\mathbb{D} \setminus \mathcal{M}$  is disconnected.

2014 Calegari-Koch-Walker proved that both  $\mathbb{D} \setminus \mathcal{M}$  and  $\mathbb{D} \setminus \mathcal{M}_0$  have infinitely many components.

# History

1985 Barnsley & Harrington introduce  $\mathcal{M}$ .

1988 – 1992 Bousch proved both  $\mathcal{M}$  and  $\mathcal{M}_0$  are connected and locally connected.

2002 Bandt proved  $\mathbb{D} \setminus \mathcal{M}$  is disconnected.

2013 Tiozzo showed  $\mathcal{M}_0$  arises as the closure of the set (restricted to  $\mathbb{D}$ ) of Galois conjugates of growth rates of postcritically finite real quadratic polynomials.

2014 Calegari-Koch-Walker proved that both  $\mathbb{D} \setminus \mathcal{M}$  and  $\mathbb{D} \setminus \mathcal{M}_0$  have infinitely many components.

# History

1985 Barnsley & Harrington introduce  $\mathcal{M}$ .

1988 – 1992 Bousch proved both  $\mathcal{M}$  and  $\mathcal{M}_0$  are connected and locally connected.

2002 Bandt proved  $\mathbb{D} \setminus \mathcal{M}$  is disconnected.

2009 Eroğlu-Rohde-Solomyak showed quasisymmetric conjugacy between quadratic dynamics and some IFS.

2013 Tiozzo showed  $\mathcal{M}_0$  arises as the closure of the set (restricted to  $\mathbb{D}$ ) of Galois conjugates of growth rates of postcritically finite real quadratic polynomials.

2014 Calegari-Koch-Walker proved that both  $\mathbb{D} \setminus \mathcal{M}$  and  $\mathbb{D} \setminus \mathcal{M}_0$  have infinitely many components.

# Motivation

Is there a way to discern holes in  $\mathcal{M}$ ? Given a  $\lambda \in \partial\mathcal{M}$  is it possible to construct a path in  $\mathbb{D} \setminus \mathcal{M}$  that converges to it?



# Motivation

Is there a way to discern holes in  $\mathcal{M}_0$ ? Given a  $\lambda \in \partial\mathcal{M}_0$  is it possible to construct a path in  $\mathbb{D} \setminus \mathcal{M}_0$  that converges to it?

# Motivation

Is there a way to discern holes in  $\mathcal{M}_0$ ? Given a  $\lambda \in \partial\mathcal{M}_0$  is it possible to construct a path in  $\mathbb{D} \setminus \mathcal{M}_0$  that converges to it?

**Conjecture:** The parameters  $\lambda \in (\partial\mathcal{M} \cap \partial\mathcal{M}_0) \setminus \mathbb{R}$  are those for which the dynamics on  $A_\lambda$  is quasisymmetric conjugate to the dynamics of  $z^2 + c$  for Misiurewicz  $c$  on its Julia set.



## Other definition of $\mathcal{M}$

$A_\lambda$  is also the closure of the fixed points of finite composition of the maps  $s_-$  and  $s_+$ . For example, let  $s_{ab} = s_a \circ s_b$  then

$$\begin{aligned} z = s_{-+}(z) = -1 + \lambda + \lambda^2 z &\iff z = \frac{-1}{1 + \lambda} = -\sum_{j=0}^{\infty} (-1)^j \lambda^j \\ &= \lim_{n \rightarrow \infty} s_{(-+)^n}(0) \end{aligned}$$

## Other definition of $\mathcal{M}$

$A_\lambda$  is also the closure of the fixed points of finite composition of the maps  $s_-$  and  $s_+$ . For example, let  $s_{ab} = s_a \circ s_b$  then

$$\begin{aligned} z = s_{-+}(z) = -1 + \lambda + \lambda^2 z &\iff z = \frac{-1}{1 + \lambda} = -\sum_{j=0}^{\infty} (-1)^j \lambda^j \\ &= \lim_{n \rightarrow \infty} s_{(-+)^n}(0) \end{aligned}$$

Define the surjection  $\pi_\lambda : \Sigma = \{-, +\}^{\mathbb{N}} \rightarrow A_\lambda$

$$a = a_0 a_1 \cdots \mapsto \sum_{j=0}^{\infty} a_j \lambda^j$$

## Other definition of $\mathcal{M}$

$A_\lambda$  is also the closure of the fixed points of finite composition of the maps  $s_-$  and  $s_+$ . For example, let  $s_{ab} = s_a \circ s_b$  then

$$\begin{aligned} z = s_{-+}(z) = -1 + \lambda + \lambda^2 z &\iff z = \frac{-1}{1 + \lambda} = -\sum_{j=0}^{\infty} (-1)^j \lambda^j \\ &= \lim_{n \rightarrow \infty} s_{(-+)^n}(0) \end{aligned}$$

Define the surjection  $\pi_\lambda : \Sigma = \{-, +\}^{\mathbb{N}} \rightarrow A_\lambda$

$$a = a_0 a_1 \cdots \mapsto \sum_{j=0}^{\infty} a_j \lambda^j$$

$A_\lambda$  is connected  $\iff s_-(A_\lambda) \cap s_+(A_\lambda) \neq \emptyset \iff \exists a, b \in \Sigma$   
with  $a_0 = +, b_0 = -$  such that  $\pi_\lambda(b) = \pi_\lambda(a)$

$$\iff \sum_{j=0}^{\infty} (a_j - b_j) \lambda^j = 2 \sum_{j=0}^{\infty} c_j \lambda^j = 0, \quad c_j \in \{-1, 0, +1\}.$$

## Other definition of $\mathcal{M}$

Let  $\mathcal{F}_\lambda = \left\{ f(z) = 1 + \sum_{j=1}^{\infty} c_j z^j \mid c_j \in \{-1, 0, +1\}; f(\lambda) = 0 \right\}$   
then

$$\mathcal{M} = \{\lambda \in \mathbb{D} \mid A_\lambda \text{ is connected}\} = \{\lambda \in \mathbb{D} \mid |\mathcal{F}_\lambda| \neq 0\}$$

## Other definition of $\mathcal{M}$

Let  $\mathcal{F}_\lambda = \left\{ f(z) = 1 + \sum_{j=1}^{\infty} c_j z^j \mid c_j \in \{-1, 0, +1\}; f(\lambda) = 0 \right\}$   
then

$$\mathcal{M} = \{\lambda \in \mathbb{D} \mid A_\lambda \text{ is connected}\} = \{\lambda \in \mathbb{D} \mid |\mathcal{F}_\lambda| \neq 0\}$$

Intuitively  $\lambda \in \partial\mathcal{M} \iff O_\lambda := \mathfrak{s}_-(A_\lambda) \cap \mathfrak{s}_+(A_\lambda)$  is “thin”  
Elements in  $\mathcal{F}_\lambda$  give a complete description of  $O_\lambda$

## Other definition of $\mathcal{M}$

Let  $\mathcal{F}_\lambda = \left\{ f(z) = 1 + \sum_{j=1}^{\infty} c_j z^j \mid c_j \in \{-1, 0, +1\}; f(\lambda) = 0 \right\}$   
then

$$\mathcal{M} = \{\lambda \in \mathbb{D} \mid A_\lambda \text{ is connected}\} = \{\lambda \in \mathbb{D} \mid |\mathcal{F}_\lambda| \neq 0\}$$

Intuitively  $\lambda \in \partial\mathcal{M} \iff O_\lambda := \mathfrak{s}_-(A_\lambda) \cap \mathfrak{s}_+(A_\lambda)$  is “thin”  
Elements in  $\mathcal{F}_\lambda$  give a complete description of  $O_\lambda$

Lemma (Solomyak, 2005)

If  $|O_\lambda| \leq 2$  then  $|\mathcal{F}_\lambda| = 1$ :

- (i.)  $|O_\lambda| = 1 \iff f \in \mathcal{F}_\lambda$  has no zero coefficients.
- (ii.)  $|O_\lambda| = 2 \iff f \in \mathcal{F}_\lambda$  has exactly one zero coefficient.

# Important Theorems

Remember  $\tilde{A}_\lambda$  is the attractor of the IFS  $\{\mathfrak{s}_-, \mathfrak{s}_0, \mathfrak{s}_+\}$ .

Theorem (Solomyak, 2005)

Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with

$$f(z) = q(z) + \frac{c_{\ell+1}z^{\ell+1} + \dots + c_{\ell+p}z^p}{1 - z^p}, \quad q(z) = \sum_{j=0}^{\ell} c_j z^j$$

then  $f'(\lambda) \neq 0$  and

# Important Theorems

Remember  $\tilde{A}_\lambda$  is the attractor of the IFS  $\{\mathfrak{s}_-, \mathfrak{s}_0, \mathfrak{s}_+\}$ .

Theorem (Solomyak, 2005)

Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with

$$f(z) = q(z) + \frac{c_{\ell+1}z^{\ell+1} + \dots + c_{\ell+p}z^p}{1 - z^p}, \quad q(z) = \sum_{j=0}^{\ell} c_j z^j$$

then  $f'(\lambda) \neq 0$  and

(i.)  $\tilde{A}_\lambda$  is  $|\lambda^{-p}|$ -self similar about  $-q(\lambda)/\lambda^{\ell+1} =: \zeta$ ;

# Important Theorems

Remember  $\tilde{A}_\lambda$  is the attractor of the IFS  $\{\mathfrak{s}_-, \mathfrak{s}_0, \mathfrak{s}_+\}$ .

Theorem (Solomyak, 2005)

Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with

$$f(z) = q(z) + \frac{c_{\ell+1}z^{\ell+1} + \dots + c_{\ell+p}z^p}{1 - z^p}, \quad q(z) = \sum_{j=0}^{\ell} c_j z^j$$

then  $f'(\lambda) \neq 0$  and

- (i.)  $\tilde{A}_\lambda$  is  $|\lambda^{-p}|$ -self similar about  $-q(\lambda)/\lambda^{\ell+1} =: \zeta$ ;
- (ii.)  $\mathcal{M}$  about  $\lambda$  and  $\frac{\lambda^{\ell+1}}{f'(\lambda)}\tilde{A}_\lambda$  about  $\frac{\lambda^{\ell+1}}{f'(\lambda)}\zeta$  are asymptotically similar.

# Important Theorems

Remember  $\tilde{A}_\lambda$  is the attractor of the IFS  $\{\mathfrak{s}_-, \mathfrak{s}_0, \mathfrak{s}_+\}$ .

Theorem (Solomyak, 2005)

Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with

$$f(z) = q(z) + \frac{c_{\ell+1}z^{\ell+1} + \dots + c_{\ell+p}z^p}{1 - z^p}, \quad q(z) = \sum_{j=0}^{\ell} c_j z^j$$

then  $f'(\lambda) \neq 0$  and

- (i.)  $\tilde{A}_\lambda$  is  $|\lambda^{-p}|$ -self similar about  $-q(\lambda)/\lambda^{\ell+1} =: \zeta$ ;
- (ii.)  $\mathcal{M}$  about  $\lambda$  and  $\frac{\lambda^{\ell+1}}{f'(\lambda)}\tilde{A}_\lambda$  about  $\frac{\lambda^{\ell+1}}{f'(\lambda)}\zeta$  are asymptotically similar.
- (iii.)  $\mathcal{M}$  is asymptotically  $|\lambda^{-p}|$ -self similar about  $\lambda$ ;

# Important Theorems

Remember  $\tilde{A}_\lambda$  is the attractor of the IFS  $\{\mathfrak{s}_-, \mathfrak{s}_0, \mathfrak{s}_+\}$ .

Theorem (Solomyak, 2005)

Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with



$$f(z) = q(z) + \frac{c_{\ell+1}z^{\ell+1} + \dots + c_{\ell+p}z^p}{1 - z^p}, \quad q(z) = \sum_{j=0}^{\ell} c_j z^j$$

then  $f'(\lambda) \neq 0$  and

- (i.)  $\tilde{A}_\lambda$  is  $|\lambda^{-p}|$ -self similar about  $-q(\lambda)/\lambda^{\ell+1} =: \zeta$ ;
- (ii.)  $\mathcal{M}$  about  $\lambda$  and  $\frac{\lambda^{\ell+1}}{f'(\lambda)}\tilde{A}_\lambda$  about  $\frac{\lambda^{\ell+1}}{f'(\lambda)}\zeta$  are asymptotically similar.
- (iii.)  $\mathcal{M}$  is asymptotically  $|\lambda^{-p}|$ -self similar about  $\lambda$ ;

**Remark.** If  $f$  has no zero coefficients then we can also replace  $\mathcal{M}$  with  $\mathcal{M}_0$  and  $\tilde{A}_\lambda$  with  $A_\lambda$ .

# Important Theorems




# Important Theorems

Theorem (Calegari-Koch-Walker, 2014)

Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with  $f$  rational as before.

Then  $\exists \delta > 0$  such that  $\forall C \notin \frac{\lambda^{\ell+1}}{f'(\lambda)} (\tilde{A}_\lambda - \zeta)$  with  $|C| < \delta$ , the parameter  $C\lambda^{pn} + \lambda$  is not in  $\mathcal{M}$  for all sufficiently large  $n$ .



# Main Result

Theorem (Pérez-S., 2018)

*Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with  $f$  rational as before.*

*If  $f$  has finitely many zero coefficients and its Taylor polynomials satisfy certain conditions then  $\lambda$  is on the boundary of a hole of  $\mathcal{M}$ .*

# Main Result

Theorem (Pérez-S., 2018)

*Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with  $f$  rational as before.*

*If  $f$  has finitely many zero coefficients and its Taylor polynomials satisfy certain conditions then  $\lambda$  is on the boundary of a hole of  $\mathcal{M}$ .*

Minor adjustments to the conditions on the Taylor polynomials gives



Theorem (Pérez-S., 2018)

*Suppose  $\lambda \in \mathcal{M} \setminus \mathbb{R}$  and  $\mathcal{F}_\lambda = \{f\}$  with  $f$  rational as before.*

*If  $f$  has no zero coefficient and its Taylor polynomials satisfy certain conditions then  $\lambda$  is on the boundary of a hole of  $\mathcal{M}_0$ .*



# Idea of proof

Use the asymptotic similarity:



# Idea of proof

Use the asymptotic similarity:




Show  $\zeta \in \tilde{A}_\lambda$  is accessible (locally) from  $\mathbb{C} \setminus \tilde{A}_\lambda$ .

# Idea of proof

Show  $\zeta \in \tilde{A}_\lambda$  is accessible (locally) from  $\mathbb{C} \setminus \tilde{A}_\lambda$ .


Use the iterative construction of  $\tilde{A}_\lambda = \bigcap_{n \geq 0} \tilde{l}_\lambda^n$  where  $\tilde{l}_\lambda^n$  is the finite union of closed discs covering  $\tilde{A}_\lambda$  and the self-similarity of  $\tilde{A}_\lambda$  at  $\zeta$ .



# Idea of proof

Show  $\zeta \in \tilde{A}_\lambda$  is accessible (locally) from  $\mathbb{C} \setminus \tilde{A}_\lambda$ .


Use the iterative construction of  $\tilde{A}_\lambda = \bigcap_{n \geq 0} \tilde{l}_\lambda^n$  where  $\tilde{l}_\lambda^n$  is the finite union of closed discs covering  $\tilde{A}_\lambda$  and the self-similarity of  $\tilde{A}_\lambda$  at  $\zeta$ .



# Idea of proof

Show  $\zeta \in \tilde{A}_\lambda$  is accessible (locally) from  $\mathbb{C} \setminus \tilde{A}_\lambda$ .


Use the iterative construction of  $\tilde{A}_\lambda = \bigcap_{n \geq 0} \tilde{l}_\lambda^n$  where  $\tilde{l}_\lambda^n$  is the finite union of closed discs covering  $\tilde{A}_\lambda$  and the self-similarity of  $\tilde{A}_\lambda$  at  $\zeta$ .



# Idea of proof

Show  $\zeta \in \tilde{A}_\lambda$  is accessible (locally) from  $C \setminus A_\lambda C \setminus \tilde{A}_\lambda$ .


Use the iterative construction of  $\tilde{A}_\lambda = \bigcap_{n \geq 0} \tilde{l}_\lambda^n$  where  $\tilde{l}_\lambda^n$  is the finite union of closed discs covering  $\tilde{A}_\lambda$  and the self-similarity of  $\tilde{A}_\lambda$  at  $\zeta$ .



# Idea of proof

Show  $\zeta \in A_\lambda$  is accessible (locally) from  $\mathbb{C} \setminus A_\lambda$ .


Use the iterative construction of  $A_\lambda = \cap_{n \geq 0} I_\lambda^n$  where  $I_\lambda^n$  is the finite union of closed discs covering  $A_\lambda$  and the self-similarity of  $A_\lambda$  at  $\zeta$ .



# Idea of proof

Show  $\zeta \in A_\lambda$  is accessible (locally) from  $\mathbb{C} \setminus A_\lambda$ .


Use the iterative construction of  $A_\lambda = \cap_{n \geq 0} I_\lambda^n$  where  $I_\lambda^n$  is the finite union of closed discs covering  $A_\lambda$  and the self-similarity of  $A_\lambda$  at  $\zeta$ .



# Idea of proof

Show  $\zeta \in A_\lambda$  is accessible (locally) from  $\mathbb{C} \setminus A_\lambda$ .

Use the iterative construction of  $A_\lambda = \bigcap_{n \geq 0} I_\lambda^n$  where  $I_\lambda^n$  is the finite union of closed discs covering  $A_\lambda$  and the self-similarity of  $A_\lambda$  at  $\zeta$ .



# Notation

$\mathcal{F}_\lambda = \{f\}$  with  $f(z) = \sum_{j=0}^{\ell} c_j z^j + \frac{c_{\ell+1} z^{\ell+1} + \dots + c_{\ell+p} z^p}{1 - z^p}$  with  
 $\mathbf{c} = c_0 \cdots c_\ell (c_{\ell+1} \cdots c_{\ell+p}) \in \Sigma$  then  $O_\lambda = \{0\}$ .

# Notation

$\mathcal{F}_\lambda = \{f\}$  with  $f(z) = \sum_{j=0}^{\ell} c_j z^j + \frac{c_{\ell+1} z^{\ell+1} + \dots + c_{\ell+p} z^p}{1 - z^p}$  with

$\mathbf{c} = c_0 \cdots c_\ell (c_{\ell+1} \cdots c_{\ell+p}) \in \Sigma$  then  $O_\lambda = \{0\}$ .

Let  $\mathbf{c}|n = c_0 \cdots c_n$  and set  $f_n(\lambda) := \sum_{j=0}^n c_j \lambda^j$ .

# Notation

$\mathcal{F}_\lambda = \{f\}$  with  $f(z) = \sum_{j=0}^{\ell} c_j z^j + \frac{c_{\ell+1} z^{\ell+1} + \dots + c_{\ell+p} z^p}{1 - z^p}$  with

$\mathbf{c} = c_0 \cdots c_\ell (c_{\ell+1} \cdots c_{\ell+p}) \in \Sigma$  then  $O_\lambda = \{0\}$ .

Let  $\mathbf{c}|n = c_0 \cdots c_n$  and set  $f_n(\lambda) := \sum_{j=0}^n c_j \lambda^j$ .

Then

$$\zeta = -\frac{f_\ell(\lambda)}{\lambda^{\ell+1}}, \quad \zeta_n = \frac{f_{\ell+1+n}(\lambda) - f_\ell(\lambda)}{\lambda^{\ell+1}}.$$

# Notation

$\mathcal{F}_\lambda = \{f\}$  with  $f(z) = \sum_{j=0}^{\ell} c_j z^j + \frac{c_{\ell+1} z^{\ell+1} + \dots + c_{\ell+p} z^p}{1 - z^p}$  with

$\mathbf{c} = c_0 \cdots c_\ell (c_{\ell+1} \cdots c_{\ell+p}) \in \Sigma$  then  $O_\lambda = \{0\}$ .

Let  $\mathbf{c}|n = c_0 \cdots c_n$  and set  $f_n(\lambda) := \sum_{j=0}^n c_j \lambda^j$ .

Then

$$\zeta = -\frac{f_\ell(\lambda)}{\lambda^{\ell+1}}, \quad \zeta_n = \frac{f_{\ell+1+n}(\lambda) - f_\ell(\lambda)}{\lambda^{\ell+1}}.$$

So  $B_n$  is centered at

$$\omega_n = -(\zeta_n - \zeta) + \zeta = -\zeta_n + 2\zeta = -\frac{1}{\lambda^{\ell+1}} f_{\ell+1+n}(\lambda) + \zeta$$

with a radius

$$r_n = |\zeta_n - \omega_n| - |\lambda^{n+1}| R = \frac{2}{|\lambda^{\ell+1}|} |f_{\ell+1+n}(\lambda)| - |\lambda^{n+1}| R$$

where  $R = (1 - |\lambda|)^{-1}$ .

## Conditions on Taylor polynomials of $f$

For all  $n \geq 0$

$B_n$  exists if and only if  $r_n > 0$

$$|f_{\ell+1+n}(\lambda)| > \frac{1}{2} \frac{|\lambda^{\ell+1+n+1}|}{1 - |\lambda|}.$$

# Conditions on Taylor polynomials of $f$

For all  $n \geq 0$

$B_n$  exists if and only if  $r_n > 0$

$$|f_{\ell+1+n}(\lambda)| > \frac{1}{2} \frac{|\lambda^{\ell+1+n+1}|}{1 - |\lambda|}.$$

$B_n \cap B_{n+1} \neq \emptyset$  if and only if  $r_n + r_{n+1} > |\omega_n - \omega_{n+1}|$

$$|f_{\ell+1+n}(\lambda)| + |f_{\ell+1+n+1}(\lambda)| > \frac{|\lambda^{\ell+1+n+1}|}{1 - |\lambda|}$$

# Conditions on Taylor polynomials of $f$

For all  $n \geq 0$

$B_n$  exists if and only if  $r_n > 0$

$$|f_{\ell+1+n}(\lambda)| > \frac{1}{2} \frac{|\lambda^{\ell+1+n+1}|}{1 - |\lambda|}.$$

$B_n \cap B_{n+1} \neq \emptyset$  if and only if  $r_n + r_{n+1} > |\omega_n - \omega_{n+1}|$

$$|f_{\ell+1+n}(\lambda)| + |f_{\ell+1+n+1}(\lambda)| > \frac{|\lambda^{\ell+1+n+1}|}{1 - |\lambda|}$$

$B_n \cap \mathbb{I}_\lambda^n = \emptyset$  if and only if  $|\omega_n - \nu_{\mathbf{a}|n}| > r_n + |\lambda^{n+1}| R$  for any  $\mathbf{a} \in \Sigma$  and  $\nu_{\mathbf{a}|n} = \sum_{j \leq n} a_j \lambda^j$

$$2 |f_{\ell+1+n}(\lambda)| < \left| f_\ell(\lambda) + f_{\ell+1+n}(\lambda) + \lambda^{\ell+1} \nu_{\mathbf{a}|n} \right|$$

## Self-similarity of $|_\lambda^n$ at 0

Since

$$f(\lambda) = 0 \implies \sum_{j=\ell+1}^{\ell+p} c_j \lambda^j = (\lambda^p - 1) \sum_{j=0}^{\ell} c_j \lambda^j$$

then for any  $0 \leq n \leq p$

$$\frac{1}{\lambda^p} \left( |_\lambda^{\bar{c}|\ell+p+n} \cup |_\lambda^{c|\ell+p+n} \right) = |_\lambda^{\bar{c}|\ell+n} \cup |_\lambda^{c|\ell+n}$$



where  $\bar{c}$  is the opposite of  $c$ .

We need to check finitely many inequalities to certify  $\lambda \in \partial \mathcal{M}_0$ !

$+ (+ + -)$

## Proposition



Let  $\lambda \approx -0.366 + 0.520i$  be the root of  $f(z) = 1 + \frac{z+z^2-z^3}{1-z^3}$  then  $\lambda$  is on the boundary of a hole of both  $\mathcal{M}_0$  and  $\mathcal{M}$ .



$+ (+ + -)$

## Proposition

Let  $\lambda \approx -0.366 + 0.520i$  be the root of  $f(z) = 1 + \frac{z+z^2-z^3}{1-z^3}$  then  $\lambda$  is on the boundary of a hole of both  $\mathcal{M}_0$  and  $\mathcal{M}$ .



To show the inequalities are satisfied use estimates on  $|\lambda|$  and  $\arg(\lambda)$ , and the fact that  $1 + \lambda + \lambda^2 = 2\lambda^3$ .

$+ (+ + -)^\infty$

Proposition (Pérez-S., 2019)

*Let  $\lambda \approx -0.366 + 0.520i$  be the root of  $f(z) = 1 + \frac{z+z^2-z^3}{1-z^3}$  then  $\lambda$  is on the boundary of the main connected component of  $\mathbb{D} \setminus \mathcal{M}$ .*

$+ (+ + -)^\infty$

Proposition (Pérez-S., 2019)

*Let  $\lambda \approx -0.366 + 0.520i$  be the root of  $f(z) = 1 + \frac{z+z^2-z^3}{1-z^3}$  then  $\lambda$  is on the boundary of the main connected component of  $\mathbb{D} \setminus \mathcal{M}$ .*

$\lambda$  is a root of the polynomial  $1 + z + z^2 - 2z^3$ .

$+ (+ + -)^\infty$

Proposition (Pérez-S., 2019)

Let  $\lambda \approx -0.366 + 0.520i$  be the root of  $f(z) = 1 + \frac{z+z^2-z^3}{1-z^3}$  then  $\lambda$  is on the boundary of the main connected component of  $\mathbb{D} \setminus \mathcal{M}$ .

$\lambda$  is a root of the polynomial  $1 + z + z^2 - 2z^3$ .

$\implies (1 - \lambda^{3k})f(\lambda) = f_{3k-1}(\lambda) - 2\lambda^{3k} = 0$  for every  $k \in \mathbb{N}$ .

$+ (+ + -)^\infty$

Proposition (Pérez-S., 2019)

Let  $\lambda \approx -0.366 + 0.520i$  be the root of  $f(z) = 1 + \frac{z+z^2-z^3}{1-z^3}$  then  $\lambda$  is on the boundary of the main connected component of  $\mathbb{D} \setminus \mathcal{M}$ .

$\lambda$  is a root of the polynomial  $1 + z + z^2 - 2z^3$ .


$$\implies (1 - \lambda^{3k})f(\lambda) = f_{3k-1}(\lambda) - 2\lambda^{3k} = 0 \text{ for every } k \in \mathbb{N}.$$

Geometrically, it means that the discs with centers

$$\pm f_{3k-1}(\lambda) \pm \lambda^{3k}$$

are aligned.


# Geometry of $A_\lambda$ for $\lambda$ associated to $+ (+ + -)^\infty$



$k = 1$


# Geometry of $A_\lambda$ for $\lambda$ associated to $+ (+ + -)^\infty$

$k = 2$




# Level Curves


$\lambda$  is on the level curve  $\text{Im}\left(\frac{f_{3k-1}(z)}{z^{3k}}\right) = 0$  for every  $k \in \mathbb{N}$ .




# Block Tower



# Block Tower




# Shape Conditions




Level 2

# Shape Conditions



# Shape Conditions



**Thank You!**