

Lotka-Volterra systems with stochastic resetting

Silvio Martinico

University of Pisa
Computational Models for Complex Systems



Lotka-Volterra model

Purpose

Description of prey-predator populations interaction

The (basic) model

$$\begin{cases} \frac{\partial a(t)}{\partial t} = -\mu a(t) + \lambda a(t)b(t) \\ \frac{\partial b(t)}{\partial t} = \sigma b(t) - \lambda' a(t)b(t) \end{cases}$$

- $a(t)$ density of the population of predators
- μ mortality rate of predators
- λ reproductive rate of predators
- $b(t)$ density of the population of prey
- σ reproductive rate of prey
- λ' predation rate



Lotka-Volterra model

Purpose

Description of prey-predator populations interaction

The (basic) model

$$\begin{cases} \frac{\partial a(t)}{\partial t} = -\mu a(t) + \lambda a(t)b(t) \\ \frac{\partial b(t)}{\partial t} = \sigma b(t) - \lambda' a(t)b(t) \end{cases}$$

- $a(t)$ density of the population of predators
- μ mortality rate of predators
- λ reproductive rate of predators
- $b(t)$ density of the population of prey
- σ reproductive rate of prey
- λ' predation rate



Reinterpretation of the model

Rapid population growth + limited resources \Rightarrow scarcity and risk of extinction

How to survive?

Two things have an important role in populations surviving/persistence under unfavorable conditions:

- Spatial structure of the environment
- Movements

Our model

Fragile ecosystem, prey are scarce, i.e. confined to the origin cell (patch).

- Prey confined to a single patch
- Predators only reproduce in the prey patch
- Predators movements modeled by *Lévy flights*
- Predators subject to resetting to the prey patch



Reinterpretation of the model

Rapid population growth + limited resources \Rightarrow scarcity and risk of extinction

How to survive?

Two things have an important role in populations surviving/persistence under unfavorable conditions:

- Spatial structure of the environment
- Movements

Our model

Fragile ecosystem, prey are scarce, i.e. confined to the origin cell (patch).

- Prey confined to a single patch
- Predators only reproduce in the prey patch
- Predators movements modeled by *Lévy flights*
- Predators subject to resetting to the prey patch



Reinterpretation of the model

Rapid population growth + limited resources \Rightarrow scarcity and risk of extinction

How to survive?

Two things have an important role in populations surviving/persistence under unfavorable conditions:

- Spatial structure of the environment
- Movements

Our model

Fragile ecosystem, prey are scarce, i.e. confined to the origin cell (patch).

- Prey confined to a single patch
- Predators only reproduce in the prey patch
- Predators movements modeled by *Lévy flights*
- Predators subject to resetting to the prey patch



Motivations behind the model

Single patch

It is a limiting case of fragile system and it allows us to study the effect of predator mobility on species survival, within and outside the patch

Predators movements and resetting

- Standard models uses *Brownian diffusion* (Markovian random walks)
- Low prey density \Rightarrow foraging uncertainty \Rightarrow larger displacements and site fidelity

Random walk ignores multiscale movements and site fidelity and causes vanishing densities

- Lévy flights \Rightarrow more sites visited without preventing returns to previous sites
- Stochastic resetting to prey patch models spatial memory, traplining and cropping
- Resetting \Rightarrow stationary probability density around resetting point (home range behavior)



Motivations behind the model

Single patch

It is a limiting case of fragile system and it allows us to study the effect of predator mobility on species survival, within and outside the patch

Predators movements and resetting

- Standard models uses *Brownian diffusion* (Markovian random walks)
- Low prey density \Rightarrow foraging uncertainty \Rightarrow larger displacements and site fidelity

Random walk ignores multiscale movements and site fidelity and causes vanishing densities

- Lévy flights \Rightarrow more sites visited without preventing returns to previous sites
- Stochastic resetting to prey patch models spatial memory, traplining and cropping
- Resetting \Rightarrow stationary probability density around resetting point (home range behavior)



Focus and expectations

We want to analyze the strategies which maximize the predator populations

Efficient reproduction and foraging $\not\Rightarrow$ population maximization [42]

Predictions

- Low predator's reproductive rate + NO resetting $\xrightarrow{(?)}$ low abundance or extinction
- Low reproductive rate + large resetting rate $\xrightarrow{(?)}$ increasing density around prey patch

But...

- High density around the patch $\xrightarrow{(?)}$ high competition for prey $\xrightarrow{(?)}$ decrease density

So, we expect that predators population reaches a maximum at some finite resetting rate

Similarly, fixing resetting rate, predators density could probably be maximized by adjusting Lévy flights parameters



Focus and expectations

We want to analyze the strategies which maximize the predator populations

Efficient reproduction and foraging $\not\Rightarrow$ population maximization [42]

Predictions

- Low predator's reproductive rate + NO resetting $\xrightarrow{(?)}$ low abundance or extinction
- Low reproductive rate + large resetting rate $\xrightarrow{(?)}$ increasing density around prey patch

But...

- High density around the patch $\xrightarrow{(?)}$ high competition for prey $\xrightarrow{(?)}$ decrease density

So, we expect that predators population reaches a maximum at some finite resetting rate

Similarly, fixing resetting rate, predators density could probably be maximized by adjusting Lévy flights parameters



Setup

Environment

- D -dimensional lattice of square cells with length $R > 1$
- The center of each cell is represented by $\mathbf{n} \in \mathbb{Z}^D$

Movements

- Predators perform independent random steps of length > 1
- Probability distribution of dispersal of a predator between cells:

$$p(\mathbf{l}) = p_0 \delta_{\mathbf{l}, \mathbf{0}} + (1 - p_0) f(\mathbf{l}) , \quad \mathbf{l} \in \mathbb{Z}^D$$

- $p_0 := \mathbb{P}(\text{predator remains in the same cell after moving})$
- f dispersal distribution (\approx inverse power-law with exponent β), such that $f(0) = 0$ and:

$$\sum_{\mathbf{l}, |\mathbf{l}| \neq 0} f(\mathbf{l}) = 1$$



Setup

Environment

- D -dimensional lattice of square cells with length $R > 1$
- The center of each cell is represented by $\mathbf{n} \in \mathbb{Z}^D$

Movements

- Predators perform independent random steps of length > 1
- Probability distribution of dispersal of a predator between cells:

$$p(\mathbf{l}) = p_0 \delta_{\mathbf{l}, \mathbf{0}} + (1 - p_0) f(\mathbf{l}) , \quad \mathbf{l} \in \mathbb{Z}^D$$

- $p_0 := \mathbb{P}(\text{predator remains in the same cell after moving})$
- f dispersal distribution (\approx inverse power-law with exponent β), such that $f(0) = 0$ and:

$$\sum_{\mathbf{l}, |\mathbf{l}| \neq 0} f(\mathbf{l}) = 1$$



New model

Predators equation

$$\frac{\partial a(\mathbf{n}, t)}{\partial t} = - \underbrace{\alpha(1 - p_0)a(\mathbf{n}, t)}_{\text{1st term}} + \underbrace{\alpha \sum_{\mathbf{l}, |\mathbf{l}| \neq 0} p(\mathbf{l})a(\mathbf{n} - \mathbf{l}, t)}_{\text{2nd term}}$$
$$+ \underbrace{\lambda a_0(t)b(t)\delta_{\mathbf{n}, \mathbf{0}}}_{\text{3rd term}} - \underbrace{\mu a(\mathbf{n}, t)}_{\text{4th term}} - \underbrace{r a(\mathbf{n}, t)}_{\text{5th term}} + \underbrace{\left[r \sum_{\mathbf{m}} a(\mathbf{m}, t) \right] \delta_{\mathbf{n}, \mathbf{0}}}_{\text{6th term}}$$

Prey equation

$$\frac{\partial b(t)}{\partial t} = \sigma b(t) \left(1 - \frac{b(t) + a_0(t)}{K} \right) - \lambda' a_0(t)b(t)$$



New model

Predators equation

$$\frac{\partial a(\mathbf{n}, t)}{\partial t} = - \underbrace{\alpha(1 - p_0)a(\mathbf{n}, t)}_{\text{1st term}} + \underbrace{\alpha \sum_{\mathbf{l}, |\mathbf{l}| \neq 0} p(\mathbf{l})a(\mathbf{n} - \mathbf{l}, t)}_{\text{2nd term}}$$
$$+ \underbrace{\lambda a_0(t)b(t)\delta_{\mathbf{n}, \mathbf{0}}}_{\text{3rd term}} - \underbrace{\mu a(\mathbf{n}, t)}_{\text{4th term}} - \underbrace{r a(\mathbf{n}, t)}_{\text{5th term}} + \underbrace{\left[r \sum_{\mathbf{m}} a(\mathbf{m}, t) \right] \delta_{\mathbf{n}, \mathbf{0}}}_{\text{6th term}}$$

Prey equation

$$\frac{\partial b(t)}{\partial t} = \sigma b(t) \left(1 - \frac{b(t) + a_0(t)}{\kappa} \right) - \lambda' a_0(t)b(t)$$



Steady states

Trivial case: no movements

Absence of movements: $\alpha = r = 0$

$$\frac{\partial a(\mathbf{n}, t)}{\partial t} = \lambda a_0(t) b(t) \delta_{\mathbf{n}, \mathbf{0}} - \mu a(\mathbf{n}, t)$$

- Populations vanishing outside prey patch at large t
- Inside the prey patch:

$$\frac{\partial a(\mathbf{n}, t)}{\partial t} = \lambda a_0(t) b(t) - \mu a_0(t) = 0 \iff a_0(t) (\lambda b(t) - \mu) = 0$$

$$\text{i } a_0(t) = 0 \implies \left[\frac{\partial b(t)}{\partial t} = \sigma b(t) \left(1 - \frac{b(t)}{\kappa} \right) = 0 \iff b(t) = 0 \vee b(t) = \kappa \right]$$

$$\text{ii } b(t) = \frac{\mu}{\lambda} \implies \left[\frac{\partial b(t)}{\partial t} = \sigma \frac{\mu}{\lambda} \left(1 - \frac{\mu/\lambda + a_0(t)}{\kappa} \right) - \lambda' \frac{\mu}{\lambda} a_0(t) = 0 \iff a_0(t) = \frac{\kappa - \mu/\lambda}{1 + \lambda' \kappa/\sigma} \right]$$



General case

Prey steady states

- $b(t) = 0$
- $b(t) \neq 0 \Rightarrow \left[\frac{\partial b(t)}{\partial t} = 0 \iff \sigma \left(1 - \frac{b+a_0}{\kappa} \right) - \lambda' a_0 = 0 \iff b = K - a_0 \left(1 - \frac{\lambda' \kappa}{\sigma} \right) \right]$

Predators steady states

1. Define the Fourier series whose coefficients are the a_n s: $\hat{a}(\mathbf{k}) := \sum_n a_n e^{-ik \cdot n}$
2. Obtain an expression for a_n from $\frac{\partial a_n(t)}{\partial t} = 0$ and $b = K - a_0 \left(1 - \frac{\lambda' \kappa}{\sigma} \right)$
3. Replace it in $\hat{a}(\mathbf{k})$
4. Get a new equation for $\hat{a}(\mathbf{k})$: $\hat{a}(\mathbf{k}) = \frac{\lambda a_0 [K - a_0 (1 + \kappa \lambda' / \sigma)] + r A}{\alpha (1 - p_0) [1 - \hat{f}(\mathbf{k})] + \mu + r}$, where $A := \hat{a}(\mathbf{0})$
5. $A = \hat{a}(\mathbf{0}) = \frac{\lambda a_0 [K - a_0 (1 + \kappa \lambda' / \sigma)] + r A}{\mu + r} \implies A = \frac{\lambda a_0}{\mu} \left[K - a_0 \left(1 + \kappa \lambda' / \sigma \right) \right]$



General case

Prey steady states

- $b(t) = 0$
- $b(t) \neq 0 \Rightarrow \left[\frac{\partial b(t)}{\partial t} = 0 \iff \dot{\frac{b}{b}} = 0 \iff \sigma \left(1 - \frac{b+a_0}{\kappa} \right) - \lambda' a_0 = 0 \iff b = K - a_0 \left(1 - \frac{\lambda' \kappa}{\sigma} \right) \right]$

Predators steady states

1. Define the Fourier series whose coefficients are the a_n s: $\hat{a}(\mathbf{k}) := \sum_n a_n e^{-i\mathbf{k} \cdot \mathbf{n}}$
2. Obtain an expression for a_n from $\frac{\partial a_n(t)}{\partial t} = 0$ and $b = K - a_0 \left(1 - \frac{\lambda' \kappa}{\sigma} \right)$
3. Replace it in $\hat{a}(\mathbf{k})$
4. Get a new equation for $\hat{a}(\mathbf{k})$: $\hat{a}(\mathbf{k}) = \frac{\lambda a_0 \left[K - a_0 (1 + \kappa \lambda' / \sigma) \right] + r A}{\alpha (1 - p_0) [1 - \hat{f}(\mathbf{k})] + \mu + r}$, where $A := \hat{a}(\mathbf{0})$
5. $A = \hat{a}(\mathbf{0}) = \frac{\lambda a_0 \left[K - a_0 (1 + \kappa \lambda' / \sigma) \right] + r A}{\mu + r} \implies A = \frac{\lambda a_0}{\mu} \left[K - a_0 \left(1 + \kappa \lambda' / \sigma \right) \right]$



Stationary predators density at prey patch

- Fourier coefficients: $a_n = \frac{1}{(2\pi)^D} \int_B \hat{a}(\mathbf{k}) e^{-i\mathbf{k}\cdot\mathbf{n}} d\mathbf{k}$, where $B := (-\pi, \pi)^D$

- $a_0 = \frac{1}{(2\pi)^D} \int_B \hat{a}(\mathbf{k}) d\mathbf{k} =$

$$= \frac{1}{(2\pi)^D} \int_B \frac{\lambda a_0 \left[K - a_0 (1 + \kappa \lambda' / \sigma) \right] + r \cdot \frac{\lambda a_0}{\mu} \left[K - a_0 \left(1 + \kappa \lambda' / \sigma \right) \right]}{\alpha (1 - p_0) [1 - \hat{f}(\mathbf{k})] + \mu + r} d\mathbf{k}$$


- $a_0 = 0$
- $a_0 \neq 0$

$$\Rightarrow a_0 = \frac{1}{1 + \kappa \lambda' / \sigma} \left[K - \left(\frac{\lambda}{\mu (2\pi)^D} \int_B \frac{\mu + r}{\alpha (1 - p_0) [1 - \hat{f}(\mathbf{k})] + \mu + r} d\mathbf{k} \right)^{-1} \right]$$



Total predators population

- Population in cell $n = R^D \cdot a_n$
- Total population $N_p = \sum_n R^D a_n = R^D \sum_n a_n = R^D A = R^D \cdot \frac{\lambda a_0}{\mu} \left[K - a_0 \left(1 + \kappa \lambda' / \sigma \right) \right]$
- $\frac{\partial N_p(a_0)}{\partial a_0} \implies a_0^{max} = \frac{K}{2(1+\kappa\lambda'/\sigma)}$
- $N_p(a_0) = 0 \implies a_0 = 0 \vee a_0 = 2a_0^{max}$



Experiments

Parameters setting

- $D = 1$

- $K = 1$


- $\sigma = 1$

- $R = 10$


- $f(x) = \begin{cases} \frac{\beta-1}{2}|x|^{-\beta} & |x| > 1 \\ 0 & |x| \leq 1 \end{cases}$
- $p_0 = \frac{2}{R} \int_0^R dr \int_0^r f(x) dx = \begin{cases} 1 - \frac{1}{R} + \frac{1 - R^{2-\beta}}{(2-\beta)R} & \beta \neq 2 \\ 1 - \frac{1}{R} - \frac{\log R}{R} & \beta = 2 \end{cases}$




Total predators population in the (μ, β) -plane



Total predators population in the (β, r) -plane



Total predators population as function of r/λ




Total predators population as function of β

- $\mu = 0.01$


$r = 0$

$r = 0.01$


$r = 0.15$



(a)



(b)




(c)

- $\mu = 0.0001$


$r = 0$

$r = 0.01$


$r = 0.15$



(a)




(b)




(c)

Optimal resetting rate



(a)




(a) $\lambda = 0.1$


(b) $\lambda = 0.5$

(c) $\lambda = 0.0001$

(d) $\lambda = 0.0005$




(c)




(d)

Optimal Lévy exponent



(a)




(a) $\lambda = 0.1$


(b) $\lambda = 0.5$

(c) $\lambda = 0.0001$

(d) $\lambda = 0.0005$




(c)




(d)




Monte Carlo simulations of 2D stochastic population model




$\lambda = 0.1 \quad \mu = 0.05 \quad \sigma = 1$







$\lambda = 0.1 \quad \mu = 0.08 \quad \sigma = 1$




$\lambda = 0.1 \quad \mu = 0.01 \quad \sigma = 1$




2D stochastic model VS 1D deterministic model



(a)



(b)



(c)

Grazie per l'attenzione!



$$\begin{aligned} \sigma b(t) \left(1 - \frac{b(t) + a_0(t)}{K} \right) - \lambda' a_0(t) b(t) &= 0 \\ \sigma K - \sigma b(t) - \sigma a_0(t) - \lambda' K a_0(t) &= 0 \\ \Rightarrow a_0(t) \left(\sigma + \lambda' K \right) &= \sigma K - \sigma b(t) = \frac{K - b(t)}{\frac{\sigma}{\sigma + \lambda' K}} \\ m = 0 \quad \frac{(\lambda b - \mu)(K - b(t))}{\sigma + \lambda' K} &= 0 \Rightarrow b = \frac{\mu}{\lambda} \\ \lambda a_0 b \delta_{m_0} - \mu a(m, t) &= 0 \quad \begin{cases} m \neq 0 & \Rightarrow a_0(t) = \frac{K - \mu}{1 + \lambda' K} \\ \mu a(m, t) = 0 & \Rightarrow a = 0 \end{cases} \end{aligned}$$

$$\begin{aligned} \frac{db}{dt} &= \sigma b \left(1 - \frac{b + a_0}{K} \right) - \lambda' a_0 b = 0 \\ \bullet b = 0 &\Rightarrow \frac{db}{dt} = 0 \rightsquigarrow \text{fixed point} \\ \bullet b \neq 0 &\Rightarrow \left[\frac{db}{dt} = 0 \Leftrightarrow \sigma \left(1 - \frac{b + a_0}{K} \right) - \lambda' a_0 = 0 \Leftrightarrow \sigma - \frac{\sigma b}{K} - \frac{\sigma a_0}{K} - \lambda' a_0 = 0 \right. \\ &\quad \left. \Leftrightarrow K - b - a_0 - \frac{\lambda' K}{\sigma} a_0 = 0 \Leftrightarrow b = K - a_0 \left(1 - \frac{\lambda' K}{\sigma} \right) \right] \end{aligned}$$

$$\begin{aligned}
 \frac{\partial a(\mathbf{n}, t)}{\partial t} = & -\underbrace{\alpha(1-p_0)a(\mathbf{n}, t)}_{\text{1st term}} + \underbrace{\alpha \sum_{\mathbf{l}, |\mathbf{l}| \neq 0} p(\mathbf{l})a(\mathbf{n}-\mathbf{l}, t)}_{\text{2nd term}} \\
 & + \underbrace{\lambda a_0(t)b(t)\delta_{\mathbf{n}, \mathbf{0}}}_{\text{3rd term}} - \underbrace{\mu a(\mathbf{n}, t)}_{\text{4th term}} - \underbrace{r a(\mathbf{n}, t)}_{\text{5th term}} + \underbrace{\left[r \sum_m a(\mathbf{m}, t) \right] \delta_{\mathbf{n}, \mathbf{0}}}_{\text{6th term}}. \\
 (\ast) \quad = & -\alpha(1-p_0)a_m + \alpha \sum_{\ell} p(\ell)a_{m-\ell} + \lambda a_0 \left(K - a_0 \left(1 + \frac{K\lambda}{\sigma} \right) \right) \delta_{\mathbf{n}, \mathbf{0}} \\
 & - \mu a_m - r a_m + \left(r \sum_m a_m \right) \delta_{m, 0} = 0 \\
 \Rightarrow & a_m \left(\alpha(1-p_0) + \mu + r \right) = \alpha \sum_{\ell \neq 0} p(\ell)a_{m-\ell}
 \end{aligned}$$

$$\begin{aligned}
 p(l) = & p_0 \delta_{l,0} + (1-p_0) f(l) \\
 \Rightarrow a_m = & \begin{cases} \frac{\alpha \sum_{\ell \neq 0} p(\ell) a_{m-\ell}}{\alpha(1-p_0) + \mu + r} = \frac{\alpha \sum_{\ell \neq 0} (1-p_0) f(\ell) a_{m-\ell}}{\alpha(1-p_0) + \mu + r} & \text{if } m \neq 0 \\ \frac{\alpha \sum_{\ell \neq 0} (1-p_0) f(\ell) a_{m-\ell} + \lambda a_0 \left(K - a_0 \left(1 + \frac{K\lambda}{\sigma} \right) \right) + r \sum_m a_m}{\alpha(1-p_0) + \mu + r} & \text{if } m = 0 \end{cases}
 \end{aligned}$$



$$\begin{aligned}
 \hat{a}(\tilde{\mathbf{k}}) &\equiv \sum_n a_n e^{-i\tilde{\mathbf{k}} \cdot \mathbf{n}} \\
 &= \sum_{m \neq 0} \frac{\alpha \sum_{\ell \neq 0} (1-p_0) f(\ell) a_{m-\ell}}{\alpha (1-p_0) + \mu + \gamma} e^{-i\tilde{\mathbf{k}} \cdot \mathbf{n}} + \gamma \sum_m a_m \\
 &\quad \underbrace{\hspace{10em}}_1 \quad \underbrace{\hspace{10em}}_2 \\
 1) & e^{i\tilde{\mathbf{k}} \cdot \mathbf{n}} = e^{i\tilde{\mathbf{k}}(m-\ell) - i\tilde{\mathbf{k}} \ell} e^{\tilde{\mathbf{k}} \cdot \mathbf{n}} \\
 2) &= \sum_{m \neq 0} \frac{\alpha (1-p_0) \sum_{\ell \neq 0} f(\ell) a_{m-\ell} e^{i\tilde{\mathbf{k}}(m-\ell)} e^{-i\tilde{\mathbf{k}} \ell}}{\alpha (1-p_0) + \mu + \gamma} \\
 &= \frac{\alpha (1-p_0)}{\alpha (1-p_0) + \mu + \gamma} \sum_{\ell \neq 0} f(\ell) e^{-i\tilde{\mathbf{k}} \ell} \sum_{m \neq 0} a_{m-\ell} e^{-i\tilde{\mathbf{k}}(m-\ell)}
 \end{aligned}$$

$$\begin{aligned}
 2) &= \frac{\gamma (1-p_0) \sum_{\ell \neq 0} f(\ell) a_{\ell} + \lambda a_0 (K - a_0 (1 + \frac{K\gamma}{\alpha})) + \gamma \sum_m a_m}{\alpha (1-p_0) + \mu + \gamma} \\
 &= \underbrace{\frac{\alpha (1-p_0)}{\alpha (1-p_0) + \mu + \gamma} \sum_{\ell \neq 0} f(\ell) a_{\ell} e^{-i\tilde{\mathbf{k}} \ell} e^{-i\tilde{\mathbf{k}}(0-\ell)}}_3 + \underbrace{\frac{\lambda a_0 (K - a_0 (1 + \frac{K\gamma}{\alpha})) + \gamma \sum_m a_m}{\alpha (1-p_0) + \mu + \gamma}}_4
 \end{aligned}$$

$1) + 3)$

$$1) + 3) = \frac{d(-p_0)}{d(-p_0) + \mu + r} \left(\sum_{l \neq 0} f(l) e^{-ikl} \sum_{m \neq 0} a_{m-l} e^{-i\tilde{k}(m-l)} + \sum_{l \neq 0} f(l) e^{i\tilde{k}l} a_{-l} e^{-i\tilde{k}(0-l)} \right)$$

$$= \frac{d(-p_0)}{d(-p_0) + \mu + r} \left[\sum_{l \neq 0} f(l) e^{-ikl} \left(\sum_{m \neq 0} a_{m-l} e^{-i\tilde{k}(m-l)} + a_{-l} e^{-i\tilde{k}(0-l)} \right) \right]$$

$$= \frac{d(-p_0)}{d(-p_0) + \mu + r} \left[\sum_{l \neq 0} f(l) e^{-ikl} \underbrace{\sum_m a_{m-l} e^{-i\tilde{k}(m-l)}}_{{\hat{Q}}(\tilde{k})} \right]$$

$\hat{Q}(\tilde{k})$ (translation is the identity time
we are summing over \mathbb{Z}).

$$f(l)=0$$

$$\downarrow \frac{d(-p_0)}{d(-p_0) + \mu + r} \hat{a}(\tilde{k}) \underbrace{\sum_l f(l) e^{-ikl}}_{\hat{f}(\tilde{k})} = \frac{d(-p_0)}{d(-p_0) + \mu + r} \hat{a}(\tilde{k}) \hat{f}(\tilde{k})$$

4 + 5

$$\Rightarrow \hat{\alpha}(\tilde{k}) = \frac{\alpha(1-p_0)}{\alpha(1-p_0)+\mu+r} \hat{f}(\tilde{k}) \hat{\alpha}(\tilde{k}) + \frac{\lambda a_0(K - a_0(1 + \frac{K\lambda'}{\sigma})) + r \sum_m a_m}{\alpha(1-p_0)+\mu+r}$$

$$\Rightarrow \hat{\alpha}(\tilde{k}) \left(1 - \frac{\alpha(1-p_0)}{\alpha(1-p_0)+\mu+r} \hat{f}(\tilde{k}) \right) = \frac{\lambda a_0(K - a_0(1 + \frac{K\lambda'}{\sigma})) + r \sum_m a_m}{\alpha(1-p_0)+\mu+r}$$

$$\Rightarrow \hat{\alpha}(\tilde{k}) \left(\frac{\alpha(1-p_0)[1 - \hat{f}(\tilde{k})] + \mu + r}{\alpha(1-p_0)+\mu+r} \right) = \frac{\lambda a_0(K - a_0(1 + \frac{K\lambda'}{\sigma})) + r \sum_m a_m}{\alpha(1-p_0)+\mu+r}$$

$$\Rightarrow \hat{\alpha}(\tilde{k}) = \frac{\lambda a_0(K - a_0(1 + \frac{K\lambda'}{\sigma})) + r \sum_m a_m}{\alpha(1-p_0)[1 - \hat{f}(\tilde{k})] + \mu + r} = \frac{\lambda a_0[K - a_0(1 + K\lambda'/\sigma)] + rA}{\alpha(1-p_0)[1 - \hat{f}(k)] + \mu + r},$$

$$A := \sum_m a_m = \hat{\alpha}(0) = \frac{\lambda a_0(K - a_0(1 + \frac{K\lambda'}{\sigma})) + rA}{\mu + r}$$

□

$$\Rightarrow \mu A + rA = \lambda a_0 \left(K - a_0 \left(1 + \frac{K\lambda'}{\sigma} \right) \right) + rA$$

$$\Rightarrow A = \frac{\lambda a_0}{\mu} \left(K - a_0 \left(1 + \frac{K\lambda'}{\sigma} \right) \right)$$



$$\sqrt{2\pi} a_n = \langle \hat{a}(\kappa), e^{ink} \rangle = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} \hat{a}(\kappa) e^{-ink} d\kappa$$

$$\implies a_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} \hat{a}(\kappa) e^{-ink} d\kappa$$



$$\alpha_0 = \frac{1}{(2\pi)^D} \int_B \hat{Q}(\vec{k}) d\vec{k} = \frac{1}{(2\pi)^D} \int_B \frac{\lambda a_0 [K - a_0(1 + K\lambda'/\sigma)] + rA}{\alpha(1 - p_0)[1 - \hat{f}(\vec{k})] + \mu + r} d\vec{k}$$

$$= \frac{1}{(2\pi)^D} \int_B \frac{\lambda a_0 [K - a_0(1 + K\lambda'/\sigma)] + r \frac{\lambda}{\mu} a_0 (K - a_0(1 + \frac{K\lambda'}{\sigma}))}{\alpha(1 - p_0)[1 - \hat{f}(\vec{k})] + \mu + r} d\vec{k}$$

- $\bullet Q_0 = 0 \Leftrightarrow \alpha > 0$

- $\bullet Q_0 \neq 0 \Leftrightarrow$

$$1 = \frac{\lambda}{(2\pi)^D} \left[\int_B \frac{K(1 + \frac{\lambda}{\mu}) d\vec{k}}{\alpha(1 - p_0)[1 - \hat{f}(\vec{k})] + \mu + r} - (1 + K\lambda'/\sigma) \int_B \frac{(1 + \frac{\lambda}{\mu}) d\vec{k}}{\alpha(1 - p_0)[1 - \hat{f}(\vec{k})] + \mu + r} a_0 \right]$$

$$\Rightarrow 1 = \frac{\lambda}{\mu(2\pi)^D} \int_B \frac{\mu + r}{\alpha(1 - p_0)[1 - \hat{f}(\vec{k})] + \mu + r} d\vec{k} \cdot \left[K - (1 + K\lambda'/\sigma)a_0 \right]$$

$$\Rightarrow a_0 = \frac{1}{1 + K\lambda'/\sigma} \left[K - \left(\frac{\lambda}{\mu(2\pi)^D} \int_B \frac{\mu + r}{\alpha(1 - p_0)[1 - \hat{f}(\vec{k})] + \mu + r} d\vec{k} \right)^{-1} \right]$$

