
# 15.5. The Simple Pendulum



A simple pendulum is one which can be considered to be a point mass suspended from a string or rod of negligible mass.

$$\theta(t) = \theta_{\text{max}} \cos(\omega t + \phi)$$

where


$$\omega = \sqrt{\frac{g}{L}}$$

$$f = \frac{\omega}{2\pi} = \frac{1}{2\pi} \sqrt{\frac{g}{L}}$$

$$T = \frac{1}{f} = 2\pi \sqrt{\frac{L}{g}}$$

T and f of a simple pendulum depend only on the length L of the string and the acceleration due to gravity. They are independent of the mass m.

# 15.6. The Physical Pendulum



When the pendulum is displaced through an angle  $\theta$  from its equilibrium position, a restoring torque  $\tau$  appears.

$$\tau = -(mg\sin\theta)(h)$$


With small angle approximation:

$$\frac{d^{2}\theta}{dt^{2}} + \frac{mgh}{I}\theta = 0$$
$$\theta(t) = \theta_{\text{max}}\cos(\omega t + \phi)$$

where 
$$\omega = \sqrt{\frac{mgh}{I}}$$

$$T = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{I}{mgh}}$$

#### • Torsional Pendulum




A torsional pendulum consists of a rigid object suspended by a wire attached to a rigid support. The object oscillates about the line OP with an amplitude  $\theta_{\text{max}}$ .

$$T=2\pi\sqrt{\frac{I}{\kappa}}$$

where  $\kappa$  is the torsion constant.

### 15.6. Damped oscillations (descriptive)





One example of a damped oscillator is an object attached to a spring and submersed in a viscous liquid.

$$x(t) = A e^{-\alpha t} \cos (\omega t + \phi)$$
where  $\alpha = b/2m$ 

$$\omega = \sqrt{\frac{k}{m} - \frac{b^2}{4m^2}}$$


$$= \sqrt{\omega_0^2 - \left(\frac{b}{2m}\right)^2}$$





Graphs of position versus time for (a) an underdamped oscillator, (b) a critically damped oscillator, and (c) an overdamped oscillator.

# 15.7. Forced Oscillations (descriptive)



Graph of amplitude versus frequency for a damped oscillator when a periodic driving force is present. When the frequency  $\omega$  of the driving force equals the natural frequency  $\omega_0$  of the oscillator, *resonance* occurs. Note that the shape of the resonance curve depends on the size of the damping coefficient b.

$$x = A\cos(\omega t + \phi)$$

$$A = \frac{F_0/m}{\sqrt{(\omega^2 - \omega_0^2)^2 + \left(\frac{b\omega}{m}\right)^2}}$$

### **Chapter 16.** Wave Motion

# Main types of waves:

- Mechanical waves : sound waves, water waves... (physical medium involved)
- Electromagnetic waves: light, radio waves, television signals ... (no medium)
- Matter waves: a beam of particles, e.g. electrons (Quantum Physics)