

DATE UTC

1/1/2021

8/7/2026

CITY

All

STATION NAME

All

QUALITY BAND

All

40.21

Avg PM2.5


Kampala

Most Polluted City


84.48%

% Exceeding WHO guideline

Avg PM2.5 by city




% Exceeding WHO guideline and % Within Guideline by city




Monthly PM2.5 by City

city	January	February	March	April	May	June	July	August	September	October	November
Addis Ababa	24.65	22.70	21.29	21.44	28.78	38.53	32.67	29.51	35.37	21.40	19.1
Johannesburg								44.70			
Kampala	66.16	53.25	48.21	35.48	40.88	48.93	56.61	53.18	51.80	62.16	61.1
Kigali	48.44	39.79	30.25	25.88	28.90	45.78	56.29	56.06	48.17	34.00	26.1
Lagos	64.27	36.15	26.46	22.58	26.23	35.34	34.08	40.75	42.44	44.13	56.1
Nairobi	24.23	24.41	18.31	18.34	45.84	26.21	23.82	22.74	22.99	19.76	17.1
Total	53.40	39.96	32.43	26.43	33.10	40.60	42.49	40.81	39.50	41.13	46.1


Station Reliability Score by Station



Monitoring stations by PM2.5 level



Station Count by pollutant



Executive summary

Kampala records the highest average PM2.5 concentration among the six cities studied (52.6 $\mu\text{g}/\text{m}^3$), followed by Kigali (38.9 $\mu\text{g}/\text{m}^3$) and Lagos (38.6 $\mu\text{g}/\text{m}^3$). All six cities exceed the WHO annual PM2.5 guideline of 5 $\mu\text{g}/\text{m}^3$, and among cities with reliable sample sizes, Kigali shows the highest WHO 24-hour guideline exceedance rate at 97.1% of readings, followed closely by Kampala (95.6%) and Addis Ababa (87.3%).

Based on pollution severity, exceedance frequency, and seasonal volatility, **Kampala, Kigali, and Lagos** are identified as the top-priority cities for UNEP intervention. Johannesburg is excluded from this ranking due to insufficient monitoring infrastructure (1 active PM2.5 station, 3 total readings across the study period) and is instead flagged for monitoring-expansion investment rather than direct pollution intervention.