

Causes of variation in vaccine efficacy in a wild rodent population

Saudamini Venkatesan¹, Ewan Smith², Amy B. Pedersen^{1,†}, and Simon A. Babayan^{2,†,*}

¹The Moredun Research Institute, Pentlands Science Park, EH26 0PZ, Scotland, UK,

²Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow, G12 8QQ, Scotland, UK,

[†]These authors contributed equally to this work

^{*}Corresponding authors: amy.pedersen@ed.ac.uk, simon.babayan@glasgow.ac.uk


Individuals vary broadly in their response to vaccination and subsequent challenge infection, with poor vaccine responders causing persistence of both infection and transmission in populations.

Keywords: Helminth; vaccine; immunoparasitology; machine learning; pathway analysis; ageing.

INTRODUCTION

16 Individuals vary widely in their responses to vaccination and little is known about the
 17 causes underlying low vaccine efficacy, nor how effective a vaccine will be “in the real
 18 world” against the pathogen for which it has been developed.[1]

RESULTS



METHODS

ACKNOWLEDGEMENTS

COMPETING INTERESTS

20 The authors declare that there are no competing interests.

AUTHOR CONTRIBUTIONS

REFERENCES

- 21 1. Belkin, M. & Niyogi, P. *Using manifold structure for partially labeled classification in Advances in neural*
 22 *information processing systems* (2002), 929–936.

FIGURE LEGENDS

SUPPLEMENTARY INFORMATION

²³ **Supplementary Data 1**

²⁴ **Supplementary Data 2**

²⁵ **Supplementary Data 3**

²⁶ **Supplementary Figures**