

Theoretical particle physics

Chenhuan Wang

October 7, 2019

Contents

0	Organisational	3
1	Introduction	4
1.1	Standard Model	4
1.2	Energy scales	5
1.3	Natural units	6

0 Organisational

Tutorials: Thursday: 8-10, 10-12 Friday: 10-12, 13-15

Exam consists of four problems

- first quickies
- 2nd-4th: similar in style to homework; two will be very close to homework

One needs 50% of points from homework. May hand in pairs.

Content of the lectures

- Standard Model of particle physics
- Electroweak sector
 - gauge principle
 - Higgs mechanism
 - Yukawa interactions
 - CP-violation

Exercises

- go through basics of computing Feynman diagrams
- not to derive the formalism
- Lagrange \rightarrow Feynman rules \rightarrow amplitudes \rightarrow cross section and decay rates (measured quantities)

Literature

- Halzen and Martin, Quarks and Leptons (a lot of basics of QCD)
- Cheng and Li (includes also quantum field theory topics CP-violation in Standard Model)
- Mark Thomson
- QFT basics
 - Peskin and Schroeder
 - M.Schwartz
 - Ryder
- Okun, Leptons and Quarks

1 Introduction

1.1 Standard Model

It is the fundamental theory of nature. There are three interactions included

- electromagnetic
- weak
- strong
- Higgs boson exchange

Electromagnetic and weak interactions can be unified into electroweak interactions.

In the Sun all these three interactions and gravity are present

- Photons reaching us clearly indicate QED's involvement.
- Neutrinos produced in weak interaction. Four protons to two protons and two neutrons (Helium). Only weak interaction can change the colour of quarks.
- Binding of Helium via strong interaction and binding energy released as kinetic energy.
- Gravity brings protons together and at high temperature to give helium.

Gauge theories (Lie groups algebras)

- EM: $U(1)_{EM}, U(1)_Y$
- weak: $SU(2)_L$
- strong: $SU(3)_c$

Forces in quantum theories involve exchange particles spin 1, vector bosons

- photon γ
- weak W^\pm, W^0 and Z^0 (mixture of W^0 and hyper charge) (discovered at CERN)
- strong $g^a, a = 1, \dots, 8$ gluons (discovered at DESY)

particles with spin $\frac{1}{2}$
Leptons

$$\begin{pmatrix} \nu_e \\ e^- \end{pmatrix}_L, e_R^-; \quad \begin{pmatrix} \nu_\mu \\ \mu^- \end{pmatrix}_L, \mu_R^-; \quad \begin{pmatrix} \nu_\tau \\ \tau^- \end{pmatrix}_L, \tau_R^-$$

Quarks

$$\begin{pmatrix} u \\ d \end{pmatrix}_L, u_R, d_R; \quad \begin{pmatrix} c \\ s \end{pmatrix}_L, c_R, s_R; \quad \begin{pmatrix} t \\ b \end{pmatrix}_L, t_R, b_R$$

One complete generation

$$\begin{pmatrix} \nu_e \\ e^- \end{pmatrix}_L, e_R^-; \quad \begin{pmatrix} u \\ d \end{pmatrix}_L, u_R, d_R;$$

To remove any one part, then gauge theory is inconsistent. It is known as "anomaly". (AQFT)

Higgs boson h^0 , spin 0

In Standard Model Higgs bosons are described by complex scalar fields $\begin{pmatrix} H^+ \\ H^0 \end{pmatrix}$. h^0 is the only fundamental scalar in nature, as far as we know.

1.2 Energy scales

- binding energy of atoms 1 – 10eV
- binding energy of nucleons $\approx 1\text{MeV}$
- no known binding energy in particle physics
- protons and neutrons $\approx \Lambda_{QCD} \approx \mathcal{O}(100\text{MeV})$

Particles have masses

- electron $m_e = 511\text{keV}$
- muon $m_\mu = 105\text{MeV}$
- tau $m_\tau = 1.7\text{GeV}$
- neutrinos $m_\nu < 1\text{eV}$
- quarks*
 - $m_u \approx 3\text{MeV}$
 - $m_d \approx 5\text{MeV}$
 - ...
- photon $m_\gamma = 0$
- gluon $m_g = 0$
- Higgs $m_{Higgs} \approx 125\text{GeV}$

*mass of proton mainly comes from dynamical effect "gluon"

Colliders

- LEP 91 GeV – 200 GeV
- Tevatron($p\bar{p}$) 800 GeV – 2 TeV
- LHC 7 TeV – 13 TeV

1.3 Natural units

$$\hbar = c = 1 \quad (1.3.1)$$

$$k_B = 1 \quad (1.3.2)$$

Everything expressed in term of powers of energy.

$$1 = 1 \times 10^{-15} \text{ m} = 5 \text{ GeV}^{-1}$$