

LINMA2171 Numerical Analysis

SIMON DESMIDT

Academic year 2024-2025 - Q1



UCLouvain

Contents

1	Linear continuous-time 2D dynamical systems	2
1.1	Introduction	2
1.2	General form	2
2	Nonlinear CT, 2D systems	4
2.1	General form	4
2.2	Methodology	4
2.3	Linearization	4
2.4	Population dynamics	5
2.5	Conservative systems	5
2.6	Reversible systems	6

Linear continuous-time 2D dynamical systems

1.1 Introduction

Consider the 2D dynamical system $\dot{x} = f(x)$, $x \in \mathbb{R}^n$. Let $\Omega \subseteq \mathbb{R}^n$ that is compact and positively invariant. Assume that $f \in \mathcal{C}^1$ on Ω , i.e. it is continuously differentiable on Ω . Let $x(0) \in \Omega$. Then the system has one and only one solution for all positive times.

Ω is positively invariant if $x(0) \in \Omega \implies x(t) \in \Omega \forall t \geq 0$.

1.2 General form

The general form of a linear dynamical system is

$$\dot{x} = Ax \quad x \in \mathbb{R}^n \quad (1.1)$$

and its solution is of the form $x(t) = \exp(At)x(0)$. We will now study the different possibilities of stability, based on the matrix A :

1. A is diagonalizable:
 - (a) $\lambda_1 > \lambda_2 > 0$: unstable (repelling) node (see LINMA2370).
 - (b) $\lambda_1 > \lambda_2 = 0$:
 - (c) $\lambda_1 > 0 > \lambda_2$: saddle point.
 - (d) $0 = \lambda_1 > \lambda_2$:
 - (e) $0 > \lambda_1 > \lambda_2$: attracting node.
 - (f) $\lambda_1 = \lambda_2 > 0$: unstable star, the eigenvectors are perpendicular and the direction is repelling from the origin.
 - (g) $\lambda_1 = \lambda_2 = 0$: every point is an equilibrium.
 - (h) $\lambda_1 = \lambda_2 < 0$: stable star, the eigenvectors are perpendicular and the direction is going to the origin.
2. A has two equal eigenvalues with only one eigenvector, i.e. is not diagonalizable:
 - (a) $\lambda < 0$: convergence to the equilibrium point.
 - (b) $\lambda = 0$:

(c) $\lambda > 0$:

3. A has complex conjugate eigenvalues:

- (a) $\operatorname{Re}(\lambda) > 0$: diverges from the equilibrium in a spiral form.
- (b) $\operatorname{Re}(\lambda) = 0$: the trajectory is periodic, does not converge nor diverge.
- (c) $\operatorname{Re}(\lambda) < 0$: converges to the equilibrium in a spiral form.

Nonlinear CT, 2D systems

2.1 General form

The general form of a nonlinear CT 2D dynamical system is

$$\dot{x} = f(x) \quad x \in \mathbb{R}^2, f : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \quad (2.1)$$

A nullcline is a curve such that for an element x_i of x , $\dot{x}_i = 0$. There are two in a 2D system and their intersections are the equilibrium points. The vector field has a horizontal or vertical direction on those curves, and their sense is given by the sign of \dot{x} or \dot{y} .

→ N.B.: The nullclines cross trajectories, as they are not trajectories themselves.

An equilibrium point is stable if all eigenvalues of the jacobian matrix evaluated at that point have nonpositive real part. It is said to be hyperbolic if its real part is nonzero.

Theorem 2.1. Consider the initial value problem $\dot{x} = f(x), x(0) = x_0$. Suppose that f is continuous and all its partial derivatives are continuous for x in some open connected set $D \subseteq \mathbb{R}^n$. Then for $x_0 \in D$, the initial value problem has a solution $x(t)$ and the solution is unique.

This means that existence and uniqueness of solutions are guaranteed if f is continuously differentiable.

2.2 Methodology

1. Nullclines: Find the graph of the functions such that $\dot{x} = 0, \dot{y} = 0$.
2. Equilibrium points: Those are the intersections of both nullclines.
3. Stability: Analyze the eigenvalues of the Jacobian matrix at each equilibrium point.

2.3 Linearization

The system here is

$$\begin{aligned} \dot{x} &= f(x, y) \\ \dot{y} &= g(x, y) \end{aligned} \quad (2.2)$$

If (x^*, y^*) is one of its equilibrium points, then the Jacobian matrix for that point is

$$A := \begin{pmatrix} \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} \\ \frac{\partial g}{\partial x} & \frac{\partial g}{\partial y} \end{pmatrix}_{(x^*, y^*)} \quad (2.3)$$

and the linearized system is

$$\begin{pmatrix} \dot{u} \\ \dot{v} \end{pmatrix} = A \begin{pmatrix} u \\ v \end{pmatrix} \quad (2.4)$$


with $u = x - x^*$ and $v = y - y^*$.

2.4 Population dynamics

The Lotka-Volterra model of competition is used to model the population growth of two species fighting for the same finite resources, but not eating each other, e.g. sheeps and rabbits. The state-space model is

$$\begin{aligned} \dot{x} &= x(3 - x - 2y) \\ \dot{y} &= y(2 - y - x) \end{aligned} \quad (2.5)$$

Its equilibrium points are $(0,0)$, $(0,2)$, $(3,0)$, $(1,1)$. The first is an unstable node, the next two are stable and the last is a saddle point.



→ N.B.: the stable manifold is the boundary between the regions of the two stable equilibria. On one side of the stable manifold, every trajectory converges to one equilibrium, and to the other one on the other side. Due to this characteristic, it is also called the basin boundary.

→ N.B.: if the real part of the eigenvalues is zero, additional information is needed to conclude on the stability.

2.5 Conservative systems

A conservative system is such that there exists a quantity $V : \mathbb{R} \rightarrow \mathbb{R}$ such that it is constant, according to the time, along every trajectory. It often has the dimensions of an energy.

Theorem 2.2. Consider the system $\dot{x} = f(x)$, $x \in \mathbb{R}^2$ and $f \in \mathcal{C}^1$. Suppose that there exists a conserved quantity $E(x)$ and suppose that x^* is an isolated fixed point. If x^* is a local minimum of E , then all trajectories sufficiently close to x^* are closed.


2.6 Reversible systems

Definition 2.3. If the system $\begin{cases} \dot{x} = f(x, y) \\ \dot{y} = g(x, y) \end{cases}$ has the same equations under the transformation of coordinates

$$\begin{cases} \tau := -t \\ X := x \\ Y := -y \end{cases} \quad (2.6)$$

then the system is reversible. In 2D, this is saying that the system is reversible if f is odd and g is even.

Theorem 2.4. If (x^*, y^*) is an equilibrium point, center for the linearized system, and the system is reversible, then, when close enough to the equilibrium point, all trajectories are closed curves.



→ N.B.: Trajectories that start and end at the same fixed point are called homoclinic orbits.