

LINMA2345 Game Theory

SIMON DESMIDT

Academic year 2024-2025 - Q2



Contents

1	Basic models					
	1.1	Extensive form games	2			
	1.2	Strategic form games	3			
		Equivalent models				
	1.4	Bayesian game	4			
	1.5	Nash Equilibrium	4			
	1.6	Properties	5			
	1.7	How do find them	5			


Basic models

In a model, we need to define the actions, the knowledge, the randomness and the outcomes.

A private information is any knowledge on the state of the world that is not a common knowledge. An information X is said to be common knowledge if the statement (everybody knows that) K everybody knows X holds for any integer K

1.1 Extensive form games

The extensive form is a tree. The nodes trigger something: they can be triggered by a player of by chance. The edges are the actual events of the game, from a player or from chance. The leaves are the payoffs, containing one item per player.



The extensive form contains the following elements:

- *N*: the set of players (e.g. by name);
- *S*: the set of information states, i.e. the information to which each player has access;
- *D*: the set of moves for a player that is at a certain information states (meet or pass, raise or stop, etc.);
- *V*: the set of nodes (e.g. numbered);
- V_0 : the set of chance nodes;
- Y: the set of action nodes, i.e. where someone does something;

- *X*: the set of leaves;
- t: the transitions, i.e. the outgoing edges from a node;
- *p*: the probability distribution for the random transitions;
- *w*: the payoffs for each player at each leaf.

1.1.1 Concepts

- Perfect recall: Players remember their past states and moves;
- Perfect information: Every player knows where they are in the game;
- Pure strategy: A function associates a move to each information state, i.e. if something happens, the strategy says what to do next;
- Payoff of a strategy: It is the expected final value, i.e. the probability of an result times its payoff;
- Randomized strategy: We assign, at each information state, a probability distribution on the moves.

1.2 Strategic form games

A strategic game is a tuple $\Gamma = (N, C, u)$ where N is the set of players, C the set of pure strategies for all players, and u the payoff function. Using a randomized strategy, we have the following expected payoff formula, with the probability distribution σ_i :

$$E_{\sigma}(u_{i}(c)) = \sum_{c \in C} \sigma(x)u_{i}(x) = \sum_{c_{i} \in C_{i}} \sigma_{i}(c_{i}) \sum_{c_{-i} \in C_{-i}} \sigma_{-i}(c_{-i})u_{i}(c_{i}, c_{-i})$$
(1.1)

where -i denotes the set of all players except i, and $\sigma_{-i}(c_{-i}) = \prod_{j \neq i} \sigma_j(c_j)$. Consider a player $i \in N$. If the players -i play following the strategy profile σ_{-i} , then the set of best response strategies for player i is given by

$$\arg \max_{c_i \in C_i} \sum_{c_{-i} \in C_{-i}} \sigma_{-i}(c_{-i}) u_i(c_i, c_{-i})$$
(1.2)

1.2.1 Strong domination

A strategy is strongly dominated if it cannot be a best response.

Corentin vs François		В		D
a	8, 5	4, 6	1, 6	7, 8
b	5, 6	7, 7	4, 4	2, 9
c	6, 7	5, 5	2, 6	3, 6
d	5, 4	5, 2	1, 6 4, 4 2, 6 6, 5	7, 5

It means that, for a given column, there exists another one where all the second numbers are strictly bigger than in the considered one. This works too for the rows, using the first numbers. In that case, the column or row can be completely removed, as the player never has an interest in using that strategy. In the example above, column B is strongly dominated by column D.

→ Note: once a column or row has been removed, we can continue to iterate on the smaller table until no row or column can be removed again. The convergence to a certain table is guaranteed, whatever the order of removal.

1.2.2 Weak domination

Weak domination changes the strictly bigger into a bigger or equal. In that case, it is not always a good move to remove the strategy, particularly if the game includes randomness.

→ Note: Removing the weakly dominated strategies does not conserve convergence.

1.3 Equivalent models

Two games are fully equivalent if

$$\forall i \in N, \ \exists A_i > 0, B_i : \ \hat{u}_i(c) = A_i u_i(c) + B_i \qquad \forall c \in C$$
 (1.3)

This means that the scale of the payoffs changes nothing.

1.3.1 Best-reponse equivalence

Two games are best-response equivalent iff they have the best response sets, i.e.

$$c_{i}' \in \arg\max_{c_{i} \in C_{i}} \sum_{c_{-i} \in C_{-i}} \sigma_{-i}(c_{-i}) u_{i}(c_{i}, c_{-i}) \iff c_{i}' \in \arg\max_{c_{i} \in C_{i}} \sum_{c_{-i} \in C_{-i}} \sigma_{-i}(c_{-i}) \hat{u}_{i}(c_{i}, c_{-i})$$

$$(1.4)$$

1.4 Bayesian game

In a Bayesian game, we add the set of player types *T* and the belief functions. What was before a set of players is now a set of agents, each of which is a player in a certain state depending on the randomness of some piece of information.

1.5 Nash Equilibrium

1.5.1 Definitions

Given a game in extensive form Γ_e, a pure strategy for a player i ∈ N is an element
 c ∈ C_i = ×_{s∈S_i}D_s, acting as a function that associates to each information state of
 the player a move to be played at that state. A pure strategy is such that, for any
 move of the opponent(s) and at any time in the game, it prescribes a move to be
 played.

• A randomized strategy $(\sigma_1, \ldots, \sigma_N)$ consists in choosing one of the pure strategies with a certain probability. For example, in rock paper scissors, a strategy is to do one of the three possibilities with probability 1/3 for each. They verify the following conditions:

$$\sum_{c_i \in D_i} \sigma_i(c_i) = 1$$

$$\forall d_i \in D_i : \qquad \sigma_i(d_i) \ge 0$$

$$\forall e_i \in C_i \setminus D_i : \qquad \sigma_i(e_i) = 0$$

$$(1.5)$$

• A Nash equilibrium is such that no player alone has an interest to deviate from his current situation. For example, in the prisoner's dilemma, the Nash equilibrium is that both players defect, because none of them can decide to cooperate and have a higher payoff.

In a Nash equilibrium, all strategies are best responses to σ_{-i} , and thus they payoffs are equal:

$$\forall d_{i} \in D_{i} : \sum_{c_{-i} \in D_{-i}} \left(\prod_{j \in N-i} \sigma_{j}(c_{j}) \right) u_{i}(c_{-i}, d_{i}) = w_{i}$$

$$\forall e_{i} \in C_{i} \setminus D_{i} : \sum_{c_{-i} \in D_{-i}} \left(\prod_{j \in N-i} \sigma_{j}(c_{j}) \right) u_{i}(c_{-i}, e_{i}) \leq w_{i}$$

$$(1.6)$$

- \rightarrow Note: the product in those equations corresponds to the probabilities of the other players strategies, and the u_i next to it correspond to the rewards considering each player's strategy. The first equation means that all selected strategies have the same payoff, and the second one means that all strategies are the best response.
- The support of a game consists in the set of all possible combinations of strategies for all players. It can contain pure strategies as well as randomized strategies.

1.6 Properties

- There always exists at least one Nash equilibrium.
- If there exists more than one Nash equilibrium in pure strategies, then there exists an infinity of them in randomized strategies, as any linear combination of the Nash equilibria is also a Nash equilibrium.

1.7 How do find them

- Remove all strongly dominated strategies;
- Check if some of the remaining strategies will never be played, whatever the opponents choose.