



---

# LINMA2460 Nonlinear Programming

---

SIMON DESMIDT  
ISSAMBRE L'HERMITE DUMONT

Academic year 2024-2025 - Q2



UCLouvain

# Contents

<b>1</b>	<b>Definitions, notations and random properties</b>	<b>2</b>
1.1	Properties . . . . .	3
1.2	Complexity table . . . . .	3
<b>2</b>	<b>TODO</b>	<b>4</b>
<b>3</b>	<b>Gradient descent without gradient</b>	<b>6</b>

# Definitions, notations and random properties

- The Taylor expansion of order  $p$  of the function  $f$  around  $x_k$  and evaluated at  $y$  is:

$$T_p(y; x_k) = f(x_k) + \sum_{i=1}^p \frac{1}{i!} D^i f(x_k) (y - x_k)^i \quad (1.1)$$

- We can thus define the gradient w.r.t.  $y$  of the Taylor expansion of order  $p$  of  $f$  around  $x_k$  and evaluated at  $x_{k+1}$ :

$$\nabla_y T_p(x_{k+1}; x_k) = \nabla_y T_p(y; x_k) \big|_{y=x_{k+1}} \quad (1.2)$$

- An oracle is a "black box" that gives information about the derivatives based on  $x$ . The general form of an oracle is:

$$\text{p-order oracle: } x \mapsto \{D^i f(x)\}_{i=0}^p \quad (1.3)$$

And so we have the following simple oracles examples:

$$\begin{aligned} \text{Zero}^{th}\text{-order oracle: } x &\mapsto \{f(x)\} \\ \text{First-order oracle: } x &\mapsto \{f(x), \nabla f(x)\} \\ \text{Second-order oracle: } x &\mapsto \{f(x), \nabla f(x), \nabla^2 f(x)\} \end{aligned} \quad (1.4)$$

- $\mathcal{C}_L^p(\mathbb{R}^n)$ : Class of functions  $p$ -times continuously differentiable with  $L$ -Lipschitz continuous  $p$ -order derivative, i.e.  $\|D^p f(x) - D^p f(y)\| \leq L\|x - y\|, \forall x, y \in \mathbb{R}^n$ . And so we have the following simple classes of problems:

- $\mathcal{C}_L^1(\mathbb{R}^n)$ : Class of continuously differentiable functions with  $L$ -Lipschitz gradient;
- $\mathcal{C}_L^2(\mathbb{R}^n)$ : Class of continuously differentiable functions with  $L$ -Lipschitz hessian.

- $p$  order method (generalization of GM):

$$x_{k+1} = \arg \min_{y \in \mathbb{R}^n} \Omega_{x_k, y, p}(y) \equiv T_{x_k, p}(y) + \frac{M}{(p+1)!} \|y - x_k\|^{p+1} \quad (1.5)$$

## 1.1 Properties

- For a function  $f \in \mathcal{C}^1(\Omega)$  and  $\Omega$  is bounded, the following holds:  $\|\nabla f(x)\| \leq L$  for all  $x \in \Omega$  for some  $L \geq 0$ .
- By the mean value theorem, for a continuously differentiable function  $f$ ,  $\forall x, y \in \Omega$ ,  $\exists z \in \Omega : f(y) - f(x) = \langle \nabla f(z), y - x \rangle$ .
- For a matrix  $A$  and a scalar  $b$ ,  $\|A\| \leq b \implies |\lambda(A)| \leq b \implies |A| \preceq bI_n$ , where the absolute value of the matrix is taken component wise.

## 1.2 Complexity table

Method	Lipschitz	$\nabla f$	$\nabla^2 f$	...	$\nabla^p f$
Zero order		$O(n\epsilon^{-2})$			
First order	$p = 1$	$O(\epsilon^{-2})$			
Second order	$p = 2$	$\times$	$O(\epsilon^{-3/2})$		
$\vdots$		$\times$	$\times$	$\ddots$	
p order		$\times$	$\times$	$\times$	$O(\epsilon^{-\frac{p+1}{p}})$

# TODO

We can generalise the property of a  $L$ -Lipschitz function to  $f \in \mathcal{C}_L^p(\mathbb{R}^n)$ . For  $p = 1$ , we had

$$f(y) \leq f(x_k) + \langle \nabla f(x_k), y - x_k \rangle + \frac{L}{2} \|y - x_k\|^2 \quad \forall y \in \mathbb{R}^n \quad (2.1)$$

For a general value of  $p$ , it becomes

$$f(y) \leq T_p(y; x_k) + \frac{L}{(p+1)!} \|y - x_k\|^{p+1} \quad \forall y \in \mathbb{R}^n \quad (2.2)$$

Using this, [we need a  \$p\$ -th order oracle](#) for the method to work.

To solve  $\min_{x \in \mathbb{R}^n} f(x)$ , we can use the iteration

$$x_{k+1} = \arg \min_{y \in \mathbb{R}^n} T_p(y; x_k) + \frac{M}{(p+1)!} \|y - x_k\|^{p+1} \quad (2.3)$$

where the constant  $M$  is an approximation of the Lipschitz constant  $L$ . [Assuming  \$f \in \mathcal{C}\_L^p\(\mathbb{R}^n\)\$](#) , we have

$$\begin{aligned} f(x_{k+1}) &\leq T_p(x_{k+1}; x_k) + \frac{L}{(p+1)!} \|x_{k+1} - x_k\|^{p+1} \\ &= \underbrace{T_p(x_{k+1}; x_k) + \frac{M}{(p+1)!} \|x_{k+1} - x_k\|^{p+1}}_{\leq f(x_k)} + \frac{(L-M)}{(p+1)!} \|x_{k+1} - x_k\|^{p+1} \end{aligned} \quad (2.4)$$

where the inequality  $\leq f(x_k)$  is due to the decrease of  $f$  and equation (2.3). [Suppose that  \$M > 2L\$](#) . After some algebraic manipulations, we get

$$f(x_k) - f(x_{k+1}) \geq \frac{L}{(p+1)!} \|x_{k+1} - x_k\|^{p+1} \quad (2.5)$$

On the other hand, using the triangular inequality,

$$\begin{aligned} \|\nabla f(x_{k+1})\| &\leq \|\nabla f(x_{k+1}) - \nabla_y T_p(x_{k+1}; x_k)\| \\ &\quad + \underbrace{\left\| \nabla_y T_p(x_{k+1}; x_k) + \nabla \left( \frac{M}{(p+1)!} \|\cdot - x_k\|^{p+1} \right) \right\|_{y=x_{k+1}}}_{=0} \\ &\quad + \left\| \nabla \left( \frac{M}{(p+1)!} \|\cdot - x_k\|^{p+1} \right) \right\|_{y=x_{k+1}} \\ &\leq \frac{L}{p!} \|x_{k+1} - x_k\|^p + \frac{M}{p!} \|x_{k+1} - x_k\|^p \end{aligned} \quad (2.6)$$

Le + rouge doit être un -?

$$\implies \|x_{k+1} - x_k\| \geq \left( \frac{p!}{L+M} \right)^{1/p} \|\nabla f(x_{k+1})\|^{1/p} \quad (2.7)$$

Combining equations (2.5) and (2.7),

$$f(x_k) - f(x_{k+1}) \geq \underbrace{\frac{L}{(p+1)!} \left( \frac{p!}{L+M} \right)^{\frac{p+1}{p}}}_{=: C(L)} \|\nabla f(x_{k+1})\|^{\frac{p+1}{p}} \quad (2.8)$$

Let  $T(\varepsilon) = \inf\{k \in \mathbb{N} : \|\nabla f(x_k)\| \leq \varepsilon\}$ . Assume that  $T(\varepsilon) \geq 2$  and  $f(x) \geq f_{low} \forall x \in \mathbb{R}^n$ . Summing up (2.8) for  $k = 0, \dots, T(\varepsilon) - 2$ ,

$$\begin{aligned} f(x_0) - f_{low} &\geq f(x_0) - f(x_{T(\varepsilon)-1}) = \sum_{k=0}^{T(\varepsilon)-2} f(x_k) - f(x_{k+1}) \\ &\geq (T(\varepsilon) - 1) C(L) \varepsilon^{\frac{p+1}{p}} \\ \implies T(\varepsilon) &\leq 1 + \frac{f(x_0) - f_{low}}{C(L)} \varepsilon^{-\frac{p+1}{p}} \equiv \mathcal{O} \left( \varepsilon^{-\frac{p+1}{p}} \right) \end{aligned} \quad (2.9)$$

# Gradient descent without gradient

For this problem consider an adversarial attack on block-based image classifier. We have a machine learning model that given an image  $a \in \mathbb{R}^p$  it return  $c(a) \in \mathbb{R}^m$ , where  $c_j(a) \in [0, 1]$  is the probability of image  $a$  to be in class  $j$ . The classifier prediction is:  $j(a) = \arg \max_{j \in [1, \dots, m]} c_j(a)$ .