



LMECA2660 - Numerical Methods in Fluid Mechanics

ALEXANDRE ORÉKHOFF
SIMON DESMIDT

This summary may not be up-to-date, the newer version is available at this address:
<https://github.com/SimonDesmidt/Syntheses>

Academic year 2025-2026 - Q2



UCLouvain

Contents

1	Finite differences with uniform grid	2
1.1	Classical finite differences	2
1.2	Richardson extrapolation	2
1.3	Operators	3

Finite differences with uniform grid

1.1 Classical finite differences

Let us define a function $u(\cdot)$ that depends on a variable x . Suppose that in the dimension x , we discretize the function uniformly with a step h and the values at the nodes are written u_i . Then, by a Taylor development series,

$$\begin{cases} u_{i+1} = u_i + h \left(\frac{\partial u}{\partial x} \right)_i + \frac{h^2}{2!} \left(\frac{\partial^2 u}{\partial x^2} \right)_i + \frac{h^3}{3!} \left(\frac{\partial^3 u}{\partial x^3} \right)_i + \frac{h^4}{4!} \left(\frac{\partial^4 u}{\partial x^4} \right)_i + \dots \\ u_{i-1} = u_i - h \left(\frac{\partial u}{\partial x} \right)_i + \frac{h^2}{2!} \left(\frac{\partial^2 u}{\partial x^2} \right)_i - \frac{h^3}{3!} \left(\frac{\partial^3 u}{\partial x^3} \right)_i + \frac{h^4}{4!} \left(\frac{\partial^4 u}{\partial x^4} \right)_i - \dots \end{cases} \quad (1.1)$$

This gives three possible finite-difference approximations:

$$\begin{aligned} \left(\frac{\partial u}{\partial x} \right)_i &= \frac{u_{i+1} - u_i}{h} + \mathcal{O}(h) && \text{(Forward differences)} \\ \left(\frac{\partial u}{\partial x} \right)_i &= \frac{u_i - u_{i-1}}{h} + \mathcal{O}(h) && \text{(Backward differences)} \\ \left(\frac{\partial u}{\partial x} \right)_i &= \frac{u_{i+1} - u_{i-1}}{2h} + \mathcal{O}(h^2) && \text{(Centered differences)} \end{aligned} \quad (1.2)$$

This also gives, for the second order,

$$\left(\frac{\partial^2 u}{\partial x^2} \right)_i = \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} - \frac{h^2}{12} \left(\frac{\partial^4 u}{\partial x^4} \right)_i + \dots \quad (1.3)$$

→ Note: in general, discentered differences are only used for stability reasons.

1.2 Richardson extrapolation

Richardson extrapolation combines centered finite differences at different scales to get a better error:

$$\begin{aligned} \frac{4}{3} \left[\left(\frac{\partial u}{\partial x} \right)_i = \frac{u_{i+1} - u_{i-1}}{2h} - \frac{h^2}{6} \left(\frac{\partial^3 u}{\partial x^3} \right)_i - \frac{h^4}{120} \left(\frac{\partial^5 u}{\partial x^5} \right)_i - \dots \right] \\ \frac{-1}{3} \left[\left(\frac{\partial u}{\partial x} \right)_i = \frac{u_{i+2} - u_{i-2}}{2(2h)} - \frac{(2h)^2}{6} \left(\frac{\partial^3 u}{\partial x^3} \right)_i - \frac{(2h)^4}{120} \left(\frac{\partial^5 u}{\partial x^5} \right)_i - \dots \right] \\ \implies \left(\frac{\partial u}{\partial x} \right)_i = \frac{8(u_{i+1} - u_{i-1}) - (u_{i+2} - u_{i-2})}{12h} + \frac{h^4}{30} \left(\frac{\partial^5 u}{\partial x^5} \right)_i - \dots \end{aligned} \quad (1.4)$$

With this method, the truncation error is of order $\mathcal{O}(h^4)$. In the same way, for second order,

$$\left(\frac{\partial^2 u}{\partial x^2}\right)_i = \frac{4}{3} \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} - \frac{1}{3} \frac{u_{i+2} - 2u_i + u_{i-2}}{(2h)^2} + \mathcal{O}(h^4) \quad (1.5)$$

1.3 Operators

Let us define the following operators:

- Forward difference: $\Delta u_i = u_{i+1} - u_i$;
- Backward difference: $\nabla u_i = u_i - u_{i-1}$;
- Centered difference: $\delta u_i = u_{i+1/2} - u_{i-1/2}$;
- Mean: $\mu u_i = \frac{1}{2}(u_{i+1/2} + u_{i-1/2})$;

→ Note: $u_{i+1/2}$ and $u_{i-1/2}$ are not computable because they are not grid values, but can be used for derivations of other formulae.

- Identity operator: $Iu_i = u_i$;
- Forward operator: $E u_i = u_{i+1}$;
- Backward operator: $E^{-1} u_i = u_{i-1}$;

→ Note: $E^{-1}E = I$.

Those operators have the following properties:

- $\mu\delta = \frac{1}{2}(E - E^{-1})$;
- $\mu^2 = I + \delta^2/4$;

The forward operator can be re-expressed using a Taylor development series:

$$\begin{aligned} E u_i &= u_{i+1} = u_i + h \frac{\partial}{\partial x} u_i + \frac{h^2}{2!} \frac{\partial^2}{\partial x^2} u_i + \frac{h^3}{3!} \frac{\partial^3}{\partial x^3} u_i + \dots \\ &= \left(I + hD + \frac{(hD)^2}{2!} + \frac{(hD)^3}{3!} + \dots \right) u_i = \exp(hD) u_i \end{aligned} \quad (1.6)$$

From this, using a second Taylor development series,

$$hD = \log(I + \Delta) = \Delta - \frac{\Delta^2}{2} + \frac{\Delta^3}{3} - \frac{\Delta^4}{4} + \dots \quad (1.7)$$

And, in the same way,

$$hD = -\log(I - \nabla) = \nabla + \frac{\nabla^2}{2} + \frac{\nabla^3}{3} + \frac{\nabla^4}{4} + \dots \quad (1.8)$$

We can do the same for another operator:

$$\mu\delta = \frac{1}{2}(E - E^{-1}) = \frac{1}{2}(\exp(hD) - \exp(-hD)) = \sinh(hD) \quad (1.9)$$

and we can use the Taylor series for $\operatorname{arc\,sinh}(x)$ but it is not very useful. By the property that $\mu^2 = I + \delta^2/4$, we get another form:

$$hD = \mu\delta \left(I - \frac{1}{6}\delta^2 + \frac{1}{30}\delta^4 - \frac{140}{\delta^6} + \dots \right) \quad (1.10)$$

If we keep only the first order term, we find the centered-difference scheme, and the terms up to second order give the Richardson extrapolation.

→ Note: in any scheme, using more information (more values, e.g. u_{i+2}, u_{i+3}, \dots) gives a more accurate solution and the order of the truncation error increases (e.g. to $\mathcal{O}(h^3)$).

page 8.