



Quantum Mechanics Assignment #1 part 5



TIF290

Simon Stefanus Jacobsson

September 2019

Feynman propagator

Take $\hbar = 1$.

The Schrödinger equation,

$$i \frac{\partial}{\partial t} |\alpha\rangle = H |\alpha\rangle, \quad (\text{SE})$$

has the solution

$$|\alpha(t)\rangle = e^{-iHt} |\alpha(0)\rangle = \sum_a |a\rangle \langle a|\alpha(0)\rangle e^{-iE_a t} \quad (1)$$

for an H -eigenbasis $\{|a\rangle\}$. We can project (1) onto the position basis by

$$\psi(\mathbf{x}, t) = \langle \mathbf{x}|\alpha(t)\rangle = \sum_a \langle \mathbf{x}|a\rangle \langle a|\alpha(0)\rangle e^{-iE_a t}. \quad (2)$$

Denote $\langle \mathbf{x}|a\rangle$ by $u_a(\mathbf{x})$ & $\langle a|\alpha(0)\rangle$ by $c_a(0)$. Using resolution of identity

$$1 = \int d^3x |\mathbf{x}\rangle \langle \mathbf{x}|, \quad (\text{RES})$$

we can express

$$c_a(0) = \int d^3x \langle a|\mathbf{x}\rangle \langle \mathbf{x}|\alpha(0)\rangle = \int d^3x u_a^*(\mathbf{x}) \psi(\mathbf{x}, 0).$$

Using (RES) again on (2) we get that

$$\begin{aligned} \psi(\mathbf{x}, t) &= \int d^3x' \sum_a \langle \mathbf{x}|a\rangle \langle a|\mathbf{x}'\rangle \langle \mathbf{x}'|\alpha(0)\rangle e^{-iE_a t} \\ &= \int d^3x' K(\mathbf{x}, \mathbf{x}', t) \psi(\mathbf{x}', 0) \end{aligned} \quad (3)$$

where

$$K(\mathbf{x}, \mathbf{x}', t) = \sum_a \langle \mathbf{x} | a \rangle \langle a | \mathbf{x}' \rangle e^{-iE_a t} = \langle \mathbf{x} | e^{-iHt} | \mathbf{x}' \rangle. \quad (4)$$

K is called *propagator* and is the Green's function for (SE) in the position basis.

Transition amplitudes

Switching to the Heisenberg picture:

$$\begin{aligned} |\alpha(t)\rangle &\mapsto |\alpha\rangle \\ \mathbf{x} &\mapsto \mathbf{x}(t) \\ e^{iHt} |\mathbf{x}\rangle &\mapsto |\mathbf{x}(t)\rangle. \end{aligned} \quad (5)$$

The last mapping follows from that operators time-evolve according to

$$\mathcal{O}(t) = e^{iHt} \mathcal{O}(0) e^{-iHt},$$

so $|\mathbf{x}\rangle$ being an eigenvector to \mathbf{x} implies $e^{iHt} |\mathbf{x}\rangle$ being an eigenvector of $\mathbf{x}(t)$.¹

We now have, by (5), that

$$K(\mathbf{x}, \mathbf{x}', t) = \langle \mathbf{x}(t) | \mathbf{x}'(0) \rangle \quad (6)$$

which is the probability amplitude of a particle starting at position $\mathbf{x}(0)$ at time 0 being observed at position $\mathbf{x}(t)$ at time t .

Path integrals

Since, for any t , $\{|x(t)\rangle\}$ form a complete eigenbasis of the state space, we can decompose K as

$$K(\mathbf{x}, \mathbf{x}'', t) = \langle \mathbf{x}(t) | \mathbf{x}'(0) \rangle = \int d^3\mathbf{x}' \langle \mathbf{x}(t) | \mathbf{x}'(t') \rangle \langle \mathbf{x}'(t') | \mathbf{x}''(0) \rangle. \quad (7)$$

Dividing some time interval $[t_1, t_N]$ into N equal parts, we can inductively use (7) to obtain

$$K(\mathbf{x}_1, \mathbf{x}_N, t_N - t_1) = \int d^3\mathbf{x}_{N-1} \int d^3\mathbf{x}_{N-2} \cdots \int d^3\mathbf{x}_2 \langle \mathbf{x}_N(t_N) | \mathbf{x}_{N-1}(t_{N-1}) \rangle \langle \mathbf{x}_{N-1}(t_{N-1}) | \mathbf{x}_{N-2}(t_{N-2}) \rangle \cdots \langle \mathbf{x}_2(t_2) | \mathbf{x}_1(t_1) \rangle.$$

This is an integral over all possible paths from \mathbf{x}_1 to \mathbf{x}_N .

Let \mathcal{L} be the classical Lagrangian & denote by Δt the difference between t_n & t_{n-1} . Then, by what Feynman discovered,

$$\langle \mathbf{x}_n(t_n) | \mathbf{x}_{n-1}(t_{n-1}) \rangle = \frac{1}{w(\Delta t)} \exp \left\{ i \int_{t_{n-1}}^{t_n} \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}) dt \right\} \quad (9)$$

¹I am here implicitly using the fact that exponents of operators can be added like $e^A e^B = e^{A+B}$ if A & B commute. Since any operator commutes with itself, this is clearly allowed here.

where $\int_{t_{n-1}}^{t_n} \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}) dt$ is an integral taken over some specific path, thus it depends implicitly on \mathbf{x}_n & \mathbf{x}_{n-1} . If we apply (9) to (8) we get

$$K(\mathbf{x}_1, \mathbf{x}_N, t_N - t_1) = \left(\frac{1}{w(\Delta t)} \right)^{N-1} \int d^3 \mathbf{x}_{N-1} \int d^3 \mathbf{x}_{N-2} \cdots \int d^3 \mathbf{x}_2 \exp \left\{ i \int_{t_1}^{t_N} \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}) dt \right\}. \quad (10)$$

Defining

$$\int_{\mathbf{x}_1}^{\mathbf{x}_N} \mathcal{D}[\mathbf{x}(t)] = \lim_{N \rightarrow \infty} \left(\frac{1}{w(\Delta t)} \right)^{N-1} \int d^3 \mathbf{x}_{N-1} \int d^3 \mathbf{x}_{N-2} \cdots \int d^3 \mathbf{x}_2$$

we have

$$K(\mathbf{x}_1, \mathbf{x}_N, t_N - t_1) = \int_{\mathbf{x}_1}^{\mathbf{x}_N} \mathcal{D}[\mathbf{x}(t)] \exp \left\{ i \int_{t_1}^{t_N} \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}) dt \right\}. \quad (11)$$

which is known as *Feynman's path integral*.

By dimensional analysis, we see that there must be a $\frac{1}{\hbar}$ factor in the exponent in (11). As $\hbar \rightarrow 0$ we can then see that any path not satisfying

$$\delta \left(\int_{t_1}^{t_N} \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}) dt \right)$$

will not have a significant contribution to K due to interference. This implies that classical mechanics is retrieved in the limit of (11) as $\hbar \rightarrow 0$.

Putting K back into SE

We are now in a position to check whether Feynman's discovery is consistent with (SE). First, consider a short time interval $[t_{n-1}, t_n]$, then (taking $m = 1$)

$$\int_{t_{n-1}}^{t_n} \mathcal{L}(\mathbf{x}, \dot{\mathbf{x}}) dt = \int_{t_{n-1}}^{t_n} \frac{1}{2} \dot{\mathbf{x}}^2 - V(\mathbf{x}) dt = \Delta t \left[\frac{1}{2} \left(\frac{\mathbf{x}_n - \mathbf{x}_{n-1}}{\Delta t} \right)^2 - V(\mathbf{x}_n) \right].$$

Now

$$\langle \mathbf{x}_n(t_n) | \mathbf{x}_{n-1}(t_{n-1}) \rangle = \frac{1}{w(\Delta t)} \exp \left\{ i \Delta t \left[\frac{1}{2} \left(\frac{\mathbf{x}_n - \mathbf{x}_{n-1}}{\Delta t} \right)^2 - V(\mathbf{x}_n) \right] \right\}. \quad (12)$$

Using $\langle \mathbf{x}_n(t_n) | \mathbf{x}_{n-1}(t_{n-1}) \rangle = \delta^3(\mathbf{x}_n - \mathbf{x}_{n-1})$ & letting $V(\mathbf{x}) = 0$ momentarily, we can integrate (12) over \mathbb{R}^3 to obtain

$$1 = \int_{\mathbb{R}^3} d^3 \mathbf{x}_n \frac{1}{w(\Delta t)} \exp \left\{ i \frac{(\mathbf{x}_n - \mathbf{x}_{n-1})^2}{2 \Delta t} \right\} = \frac{1}{w(\Delta t)} (2\pi i \Delta t)^{\frac{3}{2}} \quad (13)$$

$$w(\Delta t) = (2\pi i \Delta t)^{\frac{3}{2}}. \quad (14)$$

Assuming w is independent of V , we restate (12) as

$$\langle \mathbf{x}_n(t_n) | \mathbf{x}_{n-1}(t_{n-1}) \rangle = \frac{1}{(2\pi i \Delta t)^{\frac{3}{2}}} \exp \left\{ i \frac{(\mathbf{x}_n - \mathbf{x}_{n-1})^2}{2 \Delta t} - i \Delta t V(\mathbf{x}_n) \right\}. \quad (15)$$

Consider $\langle \mathbf{x}(t + \Delta t) | \mathbf{x}_1(t_1) \rangle$, using decomposition we can write this as

$$\langle \mathbf{x}(t + \Delta t) | \mathbf{x}_1(t_1) \rangle = \int d^3(\Delta \mathbf{x}) \frac{1}{(2\pi i \Delta t)^{\frac{3}{2}}} \exp \left\{ i \frac{(\Delta \mathbf{x})^2}{2\Delta t} - i \Delta t V(\mathbf{x}) \right\} \langle (\mathbf{x} - \Delta \mathbf{x})(t) | \mathbf{x}_1(t_1) \rangle. \quad (16)$$

Expanding (16) in power series in $\Delta \mathbf{x} = (\Delta x, \Delta y, \Delta z)$ & Δt we get

$$\begin{aligned} & \langle (\mathbf{x})(t) | \mathbf{x}_1(t_1) \rangle + \Delta t \frac{\partial}{\partial t} \langle (\mathbf{x})(t) | \mathbf{x}_1(t_1) \rangle + \mathcal{O}((\Delta t)^2) \\ &= \int d^3(\Delta \mathbf{x}) \frac{1}{(2\pi i \Delta t)^{\frac{3}{2}}} \exp \left\{ i \frac{(\Delta \mathbf{x})^2}{2\Delta t} \right\} \left[1 - i \Delta t V(\mathbf{x}) + \mathcal{O}((\Delta t)^2) \right] \langle (\mathbf{x} - \Delta \mathbf{x})(t) | \mathbf{x}_1(t_1) \rangle \\ &= \int d^3(\Delta \mathbf{x}) \frac{1}{(2\pi i \Delta t)^{\frac{3}{2}}} \exp \left\{ i \frac{(\Delta \mathbf{x})^2}{2\Delta t} \right\} \left[1 - i \Delta t V(\mathbf{x}) + \mathcal{O}((\Delta t)^2) \right] \left[1 + \frac{(\Delta \mathbf{x})^2}{2} \frac{\partial^2}{\partial \mathbf{x}^2} + \mathcal{O}(|\Delta \mathbf{x}|^3) \right] \langle \mathbf{x}(t) | \mathbf{x}_1(t_1) \rangle. \end{aligned} \quad (17)$$

Here, in the last parentheses, we have neglected any odd power of Δx , Δy , or Δz which vanishes when integrated over \mathbb{R}^3 due to symmetry (all other factors are even in $\Delta \mathbf{x}$). By $\frac{\partial^2}{\partial \mathbf{x}^2}$, we mean the Laplacian.

Using

$$\int d^3(\Delta \mathbf{x}) (\Delta \mathbf{x})^2 \exp \left\{ i \frac{(\Delta \mathbf{x})^2}{2\Delta t} \right\} = 6\sqrt{2}\pi^{3/2} (i\Delta t)^{5/2} \quad (18)$$

& pairing the first order Δt terms, we have

$$\begin{aligned} \Delta t \frac{\partial}{\partial t} \langle \mathbf{x}(t) | \mathbf{x}_1(t_1) \rangle &= \left[-\frac{(2\pi i \Delta t)^{\frac{3}{2}}}{(2\pi i \Delta t)^{\frac{3}{2}}} i \Delta t V(\mathbf{x}) + \frac{6\sqrt{2}\pi^{3/2} (i\Delta t)^{5/2}}{2(2\pi i \Delta t)^{\frac{3}{2}}} \frac{\partial^2}{\partial \mathbf{x}^2} \right] \langle \mathbf{x}(t) | \mathbf{x}_1(t_1) \rangle \\ &= \left[-i \Delta t V(\mathbf{x}) + \frac{3}{2} i \Delta t \frac{\partial^2}{\partial \mathbf{x}^2} \right] \langle \mathbf{x}(t) | \mathbf{x}_1(t_1) \rangle. \end{aligned} \quad (19)$$

Multiplying (19) by $-\frac{i}{\Delta t}$ on each side, we get

$$i \frac{\partial}{\partial t} \langle \mathbf{x}(t) | \mathbf{x}_1(t_1) \rangle = \left[V(\mathbf{x}) - \frac{3}{2} \frac{\partial^2}{\partial \mathbf{x}^2} \right] \langle \mathbf{x}(t) | \mathbf{x}_1(t_1) \rangle, \quad (20)$$

which is the (SE) in positional basis.

Remarks

When evaluating integrals such as (13) & (18), we have pushed Δt down into the lower half of the complex plane by some amount ϵ in order to achieve convergence. After having evaluated such a convergent integral, we can let $\epsilon \rightarrow 0$ to retrieve an ϵ -independent expression.