

Network Biology Analysis of Alzheimer's Disease

CXCR4 as a Hub-Bottleneck Linking Neuroinflammation and GABAergic Interneuron Vulnerability

Executive Summary

This project applies a network biology framework to RNA-seq data from post-mortem human brain tissue (fusiform gyrus, GSE125583, Genentech; 289 samples: 219 Alzheimer's disease [AD], 70 age-matched controls) to identify central genes in the molecular architecture of AD. Differential expression analysis, protein-protein interaction (PPI) network topology, and functional module detection converge on a single coherent network module linking chemokine-mediated neuroinflammation to suppressed GABAergic neurotransmission. Within this module, CXCR4 emerges as a hub-bottleneck node — a result independently corroborated by at least three previously published PPI network studies on AD using different datasets and methods.

Dataset

- Source: GSE125583 (Gene Expression Omnibus), originating from Genentech
- Samples: 289 total — 219 AD, 70 age-matched controls
- Tissue: fusiform gyrus, post-mortem
- Technology: bulk RNA-sequencing

Methodology

- 1. Differential expression analysis** — DESeq2 (R/Bioconductor). Reference-level directionality was explicitly validated against a known AD-associated marker (GFAP) prior to downstream interpretation.
- 2. Quality control and confound correction** — principal component analysis (PCA) on the initial model revealed sex, not disease status, as the dominant source of transcriptional variance in this heterogeneous 289-sample cohort. Sex was therefore included as a covariate (design: $\sim \text{sex} + \text{diagnosis}$). Key findings (GFAP, CXCR4) remained stable and significant after this correction, confirming they are not artifacts of sex imbalance between groups.
- 3. Protein-protein interaction network** — constructed via the STRING database from significant DEGs ($\text{padj} < 0.05$, $|\log_2\text{FC}| > 1$; 429 genes, 184 network nodes).
- 4. Network topology and hub identification** — Cytoscape, CytoHubba plugin (Degree and Betweenness Centrality algorithms).
- 5. Functional module detection** — MCODE clustering algorithm.
- 6. Functional enrichment** — Gene Ontology (Biological Process) and Reactome pathway analysis.
- 7. Druggability assessment** — DGIdb (Drug-Gene Interaction Database).

Key Findings

A single, functionally coherent network module

MCODE identified one dominant module comprising two interconnected, oppositely regulated sub-programs:

- Upregulated — chemokine / neuroinflammatory signaling: CXCL8, CXCR4, CCR2, CXCR3, IL7R

- Downregulated — GABAergic neurotransmission and neurotrophic support: GAD1, SST, SLC32A1, BDNF, CRH

CXCR4 as hub-bottleneck

CXCR4 shows both high degree and high betweenness centrality, identifying it as the topological bridge between the neuroinflammatory and GABAergic sub-modules — structurally analogous to EGFR/FN1/TP53 identified as hub-bottlenecks in a parallel glioblastoma network analysis, and to PPARG in a dexamethasone/airway smooth muscle model.

Independent literature validation

CXCR4 (or its ligand CXCL12) has been independently identified as a hub gene in at least three published PPI network studies of AD, using distinct datasets and analytical pipelines — a convergence that substantially strengthens confidence in this result beyond a single-dataset finding.

Note on effect size

Log2 fold-changes for validated markers (e.g., GFAP = 0.6) are modest compared to acute disease models (e.g., glioblastoma) despite high statistical significance. This is consistent with AD's nature as a chronic, progressive condition affecting a clinically heterogeneous post-mortem cohort, rather than a technical limitation of the analysis.

Biological Interpretation

The chemokine receptor CXCR4 and its ligand CXCL12 have an established, well-characterized role in the developmental migration and laminar positioning of somatostatin-, calretinin-, and neuropeptide Y-expressing cortical GABAergic interneurons, guiding their trajectory from the medial ganglionic eminence to the neocortex. A parallel body of work in schizophrenia has linked disrupted CXCR4/CXCL12 signaling to cortical GABAergic interneuron dysfunction.

Proposed model: the CXCR4/CXCL12 axis — physiologically responsible for guiding the developmental migration and laminar positioning of cortical GABAergic interneurons — is hypothesized to become aberrantly reactivated in the chronic neuroinflammatory context of AD, together with immune cell recruitment mediated by CCR2/CXCR3. This reactivation is proposed to contribute to the selective vulnerability of the very interneuron population this axis helps establish during development, rather than being a mere secondary consequence of neurodegeneration. This model is directly supported by the topological co-localization of the chemokine and GABAergic sub-modules around CXCR4 as a bridging node.

This connection between CXCR4/CXCL12 signaling and GABAergic interneuron biology is grounded in established developmental neuroscience; its extension to an adult neuroinflammatory/neurodegenerative context does not appear to have been explicitly tested in the literature reviewed for this project, and is proposed here as a data-driven, testable hypothesis rather than a demonstrated mechanism.

Functional consequence of GABAergic interneuron loss

Loss of GABAergic inhibitory tone — consistent with the downregulation of GAD1, SST, SLC32A1, and BDNF observed here — disrupts cortical excitation/inhibition balance, a mechanism increasingly recognized as an early event in AD pathogenesis, preceding overt neuronal death. Consequences include network hyperexcitability, loss of oscillatory synchronization required for memory processes, and reduced neurotrophic support via BDNF, compounding neuronal vulnerability.

Translational Relevance

Druggability assessment (DGIdb)

CXCR4 shows 60 documented drug-gene interactions in DGIdb, reflecting its maturity as a pharmacological target. Three CXCR4 antagonists are already clinically approved, though for indications unrelated to neurodegeneration:

Gene	Drug	Status	Original indication
CXCR4	Mavorixafor	Approved	HIV (antiviral)
CXCR4	Motixafortide	Approved	Antineoplastic
CXCR4	Plerixafor	Approved	Antineoplastic (HSC mobilization)

Drug repurposing opportunity

The availability of CXCR4 antagonists with an already characterized clinical safety profile represents a concrete drug repurposing opportunity: rather than developing a novel compound, these existing drugs could in principle be redirected toward testing the neuroinflammatory hypothesis proposed here, offering a substantially faster and less costly path from hypothesis to clinical testing than de novo drug development.

References

1. Frontiers in Genetics (2023) — Meta-analysis of microarray datasets identifying biological regulatory networks in Alzheimer's disease. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10497115/>
2. Genomics & Informatics (2020) — Prediction of hub genes of Alzheimer's disease using a protein interaction network and functional enrichment analysis. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7808865/>
3. Multi-algorithm topological PPI analysis identifying CXCR4 as AD hub gene (2025). <https://pmc.ncbi.nlm.nih.gov/articles/PMC12307957>
4. Tanaka et al. (2010) — CXCR4 is required for proper regional and laminar distribution of cortical somatostatin-, calretinin-, and neuropeptide Y-expressing GABAergic interneurons. Cerebral Cortex. <https://academic.oup.com/cercor/article/20/12/2810/362255>
5. Volk & Lewis (2015) — Chemokine receptors and cortical interneuron dysfunction in schizophrenia. <https://pmc.ncbi.nlm.nih.gov/articles/PMC4427549/>

Author

[Simone Marroni] — BSc in Biological Sciences, currently MSc student in Advanced Biotechnologies.

[simone.marroni@outlook.it]

Full analysis code and figures: [link al repository GitHub]