

Thermo Manual

Simone Conti

July 13, 2017

Contents

1	Introduction to Thermo	1
1.1	License and Copyright	2
1.2	Citation References	2
2	Install instructions	2
3	Usage	3
3.1	Thermo input	4
3.2	Command line options	5
4	Examples	5
4.1	Water	5
4.2	Dimerization of Insulin	6
4.3	Free Energy Difference between Peptides and Proteins Conformers	7
5	Theory	8
5.1	Partition Function	8
5.2	Helmholtz Free energy	9
5.3	Molar Helmholtz free energy	10
5.4	Internal energy	10
5.5	Molar internal energy	11
5.6	Entropy	12
5.7	Molar Entropy	12
	References	13

1 Introduction to Thermo

Thermo calculates an approximation of thermodynamics quantities like entropy and free energy for molecular systems. The calculation is based on closed-form expressions derived from statistical mechanics, essentially obtained under the rigid rotor and harmonic oscillator (RRHO) approximations.

Thermo evaluates the internal energy, the entropy and the free energy of molecules from their molecular properties like the mass, the moments of inertia, or the vibrational frequencies.

In the following, it will be described how to download and install the code and how to use it with some examples. Next, a Theory chapter aims at describing the underlying statistical mechanics to understand how this works and under which approximations and limits the obtained results can be interpreted.

1.1 License and Copyright

This code and manual has been written by Simone Conti, partially while at work at the University of Strasbourg:

© 2014-2017 Simone Conti

© 2015-2016 Université de Strasbourg

Thermo is free software: you can redistribute it and/or modify it under the terms of the GNU General Public License as published by the Free Software Foundation, either version 3 of the License, or (at your option) any later version.

Thermo is distributed in the hope that it will be useful, but WITHOUT ANY WARRANTY; without even the implied warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE. See the GNU General Public License for more details.

You should have received a copy of the GNU General Public License along with this program. If not, see <http://www.gnu.org/licenses/>.

The Thermo Manual is licensed under the Creative Commons Attribution-ShareAlike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-sa/4.0/>.



1.2 Citation References

If you use the Thermo software in scientific publications, please cite¹

S. Conti and M. Cecchini “Predicting molecular self-assembly at surfaces: a statistical thermodynamics and modeling approach”, PCCP, 2016, 18, pp. 1480-31493

2 Install instructions

The last version of the code can be downloaded from git via:

```
git clone https://github.com/SimoneCnt/thermo.git thermo
```

which will copy the latest available version inside the **thermo** directory. If you do not have git installed and you want to install it, take a look at <http://git-scm.com> Otherwise you can get the full code as a zip archive directly from the git project page at <https://github.com/SimoneCnt/thermo>

Once downloaded the installation is straightforward. The code is written in standard C and the compilation is managed by a CMake script. So to compile the code the following list of commands will easily do the work:

```
cd thermo
mkdir build
cd build
cmake ..
make release
```

After these commands, you should find the **thermo** binary inside the **build** directory. To check if the compilation went fine, you can issue the **make check** command (always from inside the build directory).

One option of Thermo is to give as input a hessian matrix to evaluate the normal mode frequencies. To activate this option you need to link Thermo to a linear algebra (lapack) library. On MacOS the linking should be automatic (the developer tools in MacOS already contains an efficient lapack library). On Linux you need first to install an efficient lapack library (e.g. MKL from Intel, freeware), and second to specify the include directory and the link options to the Thermo build script. You have two options: hope it will work automatically (unlikely), or explicitly specify the include and link directive through the **LAPACK_INC** and **LAPACK_LINK** environmental variables. For example, on my linux machine to link to MKL I have:

```
LAPACK_LINK=-L/opt/compilers_and_libraries_2017.0.098/linux/mkl/lib/intel64
-Wl,--no-as-needed -lmkl_gf_lp64 -lmkl_gnu_thread -lmkl_core -lgomp
-lpthread -lm -ldl
LAPACK_INCL=-m64 -I/opt/compilers_and_libraries_2017.0.098/linux/mkl/include
```

This is highly dependent on your system and which lapack library you use.

For any problem, feel free to refer to <https://github.com/SimoneCnt/thermo/issues> or directly to simonecnt@gmail.com.

3 Usage

To use Thermo you need to set some command line options and to write a thermo input file. For each molecule you want to study, you need to set up one thermo input file, which contains the molecular properties used to evaluate the entropy, the free energy and the internal energy. The thermo input is described first, the command line options follow.

3.1 Thermo input

The thermo input file is a simple text file which contains **key = value** pairs. Based on the “key” set, different contributions to the total free energy are evaluated. Generally speaking (see the Theory section for more details), the free energy (as the entropy and the internal energy) is split in an energetic, a translational, rotational and vibrational contributions. Usually, a molecule in solution has three rotational and three translational degrees of freedom and $3N-6$ vibrations (N is the number of atoms). This is not valid for linear molecules or for systems with constrained degrees of freedom, like molecules adsorbed on a surface. The keys described below allow you to define the repartition of the $3N$ degrees of freedom and the molecular properties, like mass and moment of inertia, necessary to evaluate the free energy.

The keys you can set in the thermo input file are:

- **temperature**: The temperature in kelvin. Default 300 K.
- **nmoles**: The number of moles in mol. Default 1 mol.
- **volume**: Volume in liters. Together with the number of moles defines the concentration. Default 1 l.
- **energy**: The ground state energy in kcal/mol. Default 0 kcal/mol.
- **mass**: Molecular mass in g/mol. Default zero.
- **translations**: Number of translational degrees of freedom. Default 3.
- **rotations**: Number of rotational degrees of freedom. Default 3. This variable has to be followed by the moments of inertia one per line, in equal number of the specified number of rotational degrees of freedom, expressed in g/mol/\AA^2 .
- **sigma**: Symmetry number. Default 1.
- **vibrations**: Number of vibrational degrees of freedom. Default zero. This variable has to be followed by the vibrational frequencies, one per line, in equal number of the specified number of vibrational degrees of freedom, expressed in cm^{-1} .
- **hessian**: Name of the file where a hessian matrix can be read. The matrix will be internally diagonalized (if Thermo is compiled with lapack support), and the normal mode frequencies calculated. The hessian and vibrations keywords are mutually exclusive. Default none.

For example, if you want to specify the temperature you can write in the input file:

```
temperature = 310
```

or for the moments of inertia:

```
rotations = 3
moment1
moment2
moment3
```

The keywords are identified based only on the first four characters, so these three statements are actually identical:

```
temperature = 300
temp = 300
temperatura = 300
```

3.2 Command line options

The mode of working of Thermo is defined by few command line options. The basic is via the `--A` option followed by the name of the thermo input file. Thus running

```
thermo --A myinput.thermo
```

will calculate all internal energy, entropy and free energy based on the options set in the `myinput.thermo` input file. The `--B` option is essentially the same as `--A`. You can use both `--A` and `--B` in the same run. This is useful in particular with the `--stechio a:b` option. The parameters `a` and `b` are the stoichiometric coefficient for the reaction $aA \rightleftharpoons bB$. Thus via this command line

```
thermo --A monomer.thermo --B dimer.thermo --stechio 2:1
```

thermo will evaluate the free energy of the monomer, the dimer, and the free energy difference for the dimerization reaction.

More classical command line options, `--out outfile.out` redirect the thermo output to the `outfile.out` file, `--help` print an hopefully useful help, and `--version` print the current version of the thermo code.

Still to document: `--cumul`, `--vdos`, `--dnu`. These essentially create and write to file the vibrational density of states (VDOS) and the cumulative vibrational free energy.

4 Examples

Here a small description on how to use the code is given. Inside the `examples` directory it is possible to find some ready to use input files for some more applications. A reference output is normally present with the `.ref` suffix.

4.1 Water

The easiest system we can take as example is water. Here a sample input script to evaluate the thermodynamic properties of an hypothetical box of 22.465 liters which contains 1 mol of gas water at 298.15 kelvin (1 atm pressure). Each water molecule will have three translational degrees of freedom, three rotational (with the associated moments of inertia), and $3N - 6 = 3$ vibrations (and associated frequencies). The symmetry number is also reported: since water symmetry pointgroup is C_{2v} , the symmetry number is 2. The mass is also specified.

```
# Set temperature in Kelvin
temperature = 298.15

# Set number of mols and volume (1 atm pressure)
nmols = 1
volume = 22.465

# Mass in g/mol
mass = 18.01528
```

```

# Translational degrees of freedom
translational = 3

# Rotational degrees of freedom and moments of inertia in g/mol Ang^2
rotational = 3
1.7704
0.6169
1.1535

# Symmetry number
sigma = 2

# Number of vibrations and their frequencies in cm-1
vibrations = 3
1635.618
3849.420
3974.869

```

Save this input as e.g. `water.inp` and run it with `thermo -A water.inp` The output will print all parsed options and at the end all thermodynamic quantities, like energy, entropy, free energy, and chemical potential, are printed. For example, the evaluated translational molar entropy is evaluated to be 32.452 cal mol⁻¹ K⁻¹ and the zero point vibrational energy to be 13.524 kcal mol⁻¹.

Input and reference output can be found in the `examples/water` subdirectory.

4.2 Dimerization of Insulin

A second example is based on the reference paper by B. Tidor and M. Karplus², published in 1994:

B. Tidor and M. Karplus, "The Contribution of Vibrational Entropy to Molecular Association: The Dimerization of Insulin", *Journal of Molecular Biology*, 238(3) 1994 pp. 405-414

In this work the effect of vibrations in the stabilization of the insulin dimer is discussed as an example of how the vibrational entropy can partially counterbalance the net loss in translational and rotational degrees of freedom. Translational and rotational contributions are calculated in good agreement with the data reported in the paper, while the vibrational ones are incorrect due to the limited number of frequencies reported in the paper itself.

The input files are inside the `examples/insulin` subdirectory. Since we want to study the dimerization reaction, it is possible to run at the same time both the monomer and the dimer, and, thanks to the `'-stechio'` command line option, the differences for the dimerization reaction are automatically printed. The Thermo command line can thus be:

```
thermo -A monomer.inp -B dimer.inp --stechio 2:1 > insulin.out
```

The reference output is `insulin.out.ref`. At the moment only reactions between two species can be studied (so only `-A` and `-B` commands are available).

4.3 Free Energy Difference between Peptides and Proteins Conformers

Three examples are included to study the difference in free energy between peptides and proteins conformers:

- alanine dipeptide (**examples/diala** directory) in its c7 equatorial (c7eq) and c7 axial (c7ax) conformations;
- beta hairpin from protein G (**examples/bhp**) in the native beta-hairpin (bhp1) and in a three-stranded beta sheet (bhp2) conformations;
- the converter of the biomolecular motor myosin VI (**examples/conv**) in the pre-powerstroke (pps) and rigor-like (rig) conformations.




Figure 1: Representation of the three examples to study the quantum correction to the conformational free energy difference: the alanine dipeptide, the beta hairpin of protein G and the converter of myosin VI. Reproduced from ref. 3.

These examples, see Fig. 1, are taken from the work of M. Cecchini, JCTC 2015³, where they were used to develop a quantum correction to the classical conformational free energy difference.

M. Cecchini, “Quantum Corrections to the Free Energy Difference between Peptides and Proteins Conformers”, Journal of Chemical Theory and Computation, 2015 ASAP

The quantum correction is by default calculated by Thermo; for example, for the myosin converter

```
thermo -A rig.inp -B pps.inp --stechio 1:1
```

Thermo calculates a quantum correction of 1.09 kcal mol⁻¹, in perfect agreement with the Cecchini paper.

5 Theory

This part is still a draft. Please, see the McQuarrie “Statistical Mechanics” book⁴ for more information.

5.1 Partition Function

$$Q = \frac{q^N}{N!} \quad (1)$$

$$\ln Q = N \ln q - N \ln N + N = N \ln \frac{qe}{N} \quad (2)$$

$$q = q_{tr} q_{rot} q_{vib} q_{elec} \quad (3)$$

Translational degrees of freedom:

$$q_{tr} = \left(\frac{2\pi m k_B T}{h^2} \right)^{t/2} V \quad (4)$$

$$\ln q_{tr} = \frac{t}{2} \ln \left(\frac{2\pi m k_B T}{h^2} \right) + \ln V \quad (5)$$

$$\frac{\partial}{\partial T} \ln q_{tr} = \frac{t}{2} \frac{1}{T} \quad (6)$$

Rotational:

$$q_{rot} = \frac{\sqrt{\pi}}{\sigma} \left(\frac{8\pi^2 k_B T}{h^2} \right)^{r/2} \left(\prod_{i=1}^r I_i \right)^{1/2} \quad (7)$$

$$\ln q_{rot} = \frac{1}{2} \ln \frac{\pi}{\sigma^2} + \frac{r}{2} \ln \frac{8\pi^2 k_B T}{h^2} + \frac{1}{2} \sum_{i=1}^r \ln I_i \quad (8)$$

$$\frac{\partial}{\partial T} \ln q_{rot} = \frac{r}{2} \frac{1}{T} \quad (9)$$

Vibrational (classical limit):

$$q_{vib,cl} = \prod_{i=1}^f \frac{k_B T}{h\nu_i} \quad (10)$$

$$\ln q_{vib,cl} = \sum_{i=1}^f \ln \frac{k_B T}{h\nu_i} \quad (11)$$

$$\frac{\partial}{\partial T} \ln q_{vib,cl} = f \frac{1}{T} \quad (12)$$

Vibrational (quantum):

$$q_{vib,qm} = \prod_{i=1}^f \frac{\exp \frac{h\nu}{2k_B T}}{1 - \exp \frac{h\nu}{k_B T}} = \prod_{i=1}^f \left[2 \sinh \left(\frac{h\nu}{2k_B T} \right) \right]^{-1} \quad (13)$$

$$\ln q_{vib,qm} = - \sum_{i=1}^f \ln \left[2 \sinh \left(\frac{h\nu}{2k_B T} \right) \right] \quad (14)$$

$$\frac{\partial}{\partial T} \ln q_{vib,qm} = \frac{1}{T} \sum_{i=1}^f \frac{\frac{h\nu}{2k_B T}}{\tanh \left(\frac{h\nu}{2k_B T} \right)} \quad (15)$$

Electronic:

$$q_{elec} = \exp \left(- \frac{E_m}{k_B T} \right) \quad (16)$$

$$\ln q_{elec} = - \frac{E_m}{k_B T} \quad (17)$$

$$\frac{\partial}{\partial T} \ln q_{elec} = \frac{E_m}{k_B T} \frac{1}{T} \quad (18)$$

5.2 Helmholtz Free energy

$$F = U - TS \quad (19)$$

$$F = -k_B T \ln Q = -Nk_B T \ln \frac{q_e}{N} \quad (20)$$

$$= -Nk_B T \ln \frac{q_{tr}^e}{N} - Nk_B T \ln q_{rot} - Nk_B T \ln q_{vib} - Nk_B T \ln q_{elec} \quad (21)$$

$$= F_{tr} + F_{rot} + F_{vib} + F_{elec} \quad (22)$$

$$F_{tr} = -Nk_B T \ln \frac{q_{tr}^e}{N} = -Nk_B T \ln \left[\left(\frac{2\pi m k_B T}{h^2} \right)^{t/2} \frac{V e}{N} \right] \quad (23)$$

$$= -Nk_B T \left(\frac{t}{2} \ln \frac{2\pi m k_B T}{h^2} + 1 - \ln \frac{N}{V} \right) \quad (24)$$

$$F_{rot} = -Nk_B T \ln q_{rot} = -Nk_B T \left[\frac{1}{2} \ln \frac{\pi}{\sigma^2} + \frac{r}{2} \ln \frac{8\pi^2 k_B T}{h^2} + \frac{1}{2} \sum_{i=1}^r \ln I_i \right] \quad (25)$$

$$F_{vib,cl} = -Nk_B T \ln q_{vib,cl} = -Nk_B T \sum_{i=1}^f \ln \frac{k_B T}{h\nu_i} \quad (26)$$

$$F_{vib,qm} = -Nk_B T \ln q_{vib,qm} = Nk_B T \sum_{i=1}^f \ln \left[2 \sinh \left(\frac{h\nu}{2k_B T} \right) \right] \quad (27)$$

$$F_{elec} = -Nk_B T \ln q_{elec} = NE_m \quad (28)$$

5.3 Molar Helmholtz free energy

$$\mu = F_m = \frac{\partial F}{\partial N} \quad (29)$$

$$= \frac{\partial F_{tr}}{\partial N} + \frac{\partial F_{rot}}{\partial N} + \frac{\partial F_{vib}}{\partial N} + \frac{\partial F_{elec}}{\partial N} \quad (30)$$

$$= \mu_{tr} + \mu_{rot} + \mu_{vib} + \mu_{elec} \quad (31)$$

$$\mu_{tr} = \frac{\partial F_{tr}}{\partial N} = -k_B T \frac{t}{2} \ln \frac{2\pi m k_B T}{h^2} + k_B T \ln \frac{N}{V} = \frac{F_{tr}}{N} + k_B T \quad (32)$$

$$\mu_{rot} = \frac{\partial F_{rot}}{\partial N} = -k_B T \left[\frac{1}{2} \ln \frac{\pi}{\sigma^2} + \frac{r}{2} \ln \frac{8\pi^2 k_B T}{h^2} + \frac{1}{2} \sum_{i=1}^r \ln I_i \right] = \frac{F_{rot}}{N} \quad (33)$$

$$\mu_{vib,cl} = \frac{\partial F_{vib,cl}}{\partial N} = -k_B T \sum_{i=1}^f \ln \frac{k_B T}{h\nu_i} = \frac{F_{vib,cl}}{N} \quad (34)$$

$$\mu_{vib,qm} = \frac{\partial F_{vib,qm}}{\partial N} = k_B T \sum_{i=1}^f \ln \left[2 \sinh \left(\frac{h\nu}{2k_B T} \right) \right] = \frac{F_{vib,qm}}{N} \quad (35)$$

$$\mu_{elec} = \frac{\partial F_{elec}}{\partial N} = E_m = \frac{F_{elec}}{N} \quad (36)$$

5.4 Internal energy

$$U = k_B T^2 \frac{\partial}{\partial T} \ln Q = Nk_B T^2 \frac{\partial}{\partial T} \ln q \quad (37)$$

$$= Nk_B T^2 \frac{\partial}{\partial T} \ln q_{tr} + Nk_B T^2 \frac{\partial}{\partial T} \ln q_{rot} + Nk_B T^2 \frac{\partial}{\partial T} \ln q_{vib} + Nk_B T^2 \frac{\partial}{\partial T} \ln q_{elec} \quad (38)$$

$$= U_{tr} + U_{rot} + U_{vib} + U_{elec} \quad (39)$$

$$U_{tr} = Nk_B T^2 \frac{\partial}{\partial T} \ln q_{tr} = Nk_B T \frac{t}{2} \quad (40)$$

$$U_{rot} = Nk_B T^2 \frac{\partial}{\partial T} \ln q_{rot} = Nk_B T \frac{r}{2} \quad (41)$$

$$U_{vib,cl} = Nk_B T^2 \frac{\partial}{\partial T} \ln q_{vib,cl} = Nk_B T f \quad (42)$$

$$U_{vib,qm} = Nk_B T^2 \frac{\partial}{\partial T} \ln q_{vib,qm} = Nk_B T \sum_{i=1}^f \frac{\frac{h\nu}{2k_B T}}{\tanh\left(\frac{h\nu}{2k_B T}\right)} \quad (43)$$

$$U_{elec} = Nk_B T^2 \frac{\partial}{\partial T} \ln q_{elec} = N E_m \quad (44)$$

5.5 Molar internal energy

$$U_m = \frac{\partial U}{\partial N} = \frac{\partial U_{tr}}{\partial N} + \frac{\partial U_{rot}}{\partial N} + \frac{\partial U_{vib}}{\partial N} + \frac{\partial U_{elec}}{\partial N} = U_{m,tr} + U_{m,rot} + U_{m,vib} + U_{m,elec} \quad (45)$$

$$U_{m,tr} = \frac{\partial U_{tr}}{\partial N} = k_B T \frac{t}{2} = \frac{U_{tr}}{N} \quad (46)$$

$$U_{m,rot} = \frac{\partial U_{rot}}{\partial N} = k_B T \frac{r}{2} = \frac{U_{rot}}{N} \quad (47)$$

$$U_{m,vib,cl} = \frac{\partial U_{vib,cl}}{\partial N} = k_B T f = \frac{U_{vib,cl}}{N} \quad (48)$$

$$U_{m,vib,qm} = \frac{\partial U_{vib,qm}}{\partial N} = k_B T \sum_{i=1}^f \frac{\frac{h\nu}{2k_B T}}{\tanh\left(\frac{h\nu}{2k_B T}\right)} = \frac{U_{vib,qm}}{N} \quad (49)$$

$$U_{m,elec} = \frac{\partial U_{elec}}{\partial N} = E_m = \frac{U_{elec}}{N} \quad (50)$$

5.6 Entropy

$$S = \frac{\partial}{\partial T} (k_B T \ln Q) = k_B \ln Q + k_B T \frac{\partial}{\partial T} \ln Q = Nk_B \ln \frac{q^e}{N} + Nk_B T \frac{\partial}{\partial T} \ln q \quad (51)$$

$$\begin{aligned} &= \left(Nk_B \ln \frac{q_{tr}^e}{N} + Nk_B T \frac{\partial}{\partial T} \ln q_{tr} \right) + \left(Nk_B \ln q_{rot} + Nk_B T \frac{\partial}{\partial T} \ln q_{rot} \right) \\ &\quad + \left(Nk_B \ln q_{vib} + Nk_B T \frac{\partial}{\partial T} \ln q_{vib} \right) + \left(Nk_B \ln q_{elec} + Nk_B T \frac{\partial}{\partial T} \ln q_{elec} \right) \end{aligned} \quad (52)$$

$$= S_{tr} + S_{rot} + S_{vib} + S_{elec} \quad (53)$$

$$S_{tr} = Nk_B \ln \frac{q_{tr}^e}{N} + Nk_B T \frac{\partial}{\partial T} \ln q_{tr} = Nk_B \left(\frac{t}{2} \ln \frac{2\pi m k_B T}{h^2} + \frac{t+2}{2} - \ln \frac{N}{V} \right) \quad (54)$$

$$S_{rot} = Nk_B \ln q_{rot} + Nk_B T \frac{\partial}{\partial T} \ln q_{rot} = Nk_B \left(\frac{1}{2} \ln \frac{\pi}{\sigma^2} + \frac{r}{2} \ln \frac{8\pi^2 k_B T}{h^2} + \frac{1}{2} \sum_{i=1}^r \ln I_i + \frac{r}{2} \right) \quad (55)$$

$$S_{vib,cl} = Nk_B \ln q_{vib,cl} + Nk_B T \frac{\partial}{\partial T} \ln q_{vib,cl} = Nk_B \left(f + \sum_{i=1}^f \ln \frac{k_B T}{h\nu_i} \right) \quad (56)$$

$$S_{vib,qm} = Nk_B \ln q_{vib,qm} + Nk_B T \frac{\partial}{\partial T} \ln q_{vib,qm} = Nk_B \sum_{i=1}^f \left[\frac{\frac{h\nu}{2k_B T}}{\tanh \frac{h\nu}{2k_B T}} - \ln \left(2 \sinh \frac{h\nu}{2k_B T} \right) \right] \quad (57)$$

$$S_{elec} = Nk_B \ln q_{elec} + Nk_B T \frac{\partial}{\partial T} \ln q_{elec} = 0 \quad (58)$$

5.7 Molar Entropy

$$S_m = \frac{\partial S}{\partial N} = \frac{\partial S_{tr}}{\partial N} + \frac{\partial S_{rot}}{\partial N} + \frac{\partial S_{vib}}{\partial N} + \frac{\partial S_{elec}}{\partial N} = S_{m,tr} + S_{m,rot} + S_{m,vib} + S_{m,elec} \quad (59)$$

$$S_{m,tr} = \frac{\partial S_{tr}}{\partial N} = k_B \left(\frac{t}{2} \ln \frac{2\pi m k_B T}{h^2} + \frac{t}{2} - \ln \frac{N}{V} \right) = \frac{S_{tr}}{N} - k_B \quad (60)$$

$$S_{m,rot} = \frac{\partial S_{rot}}{\partial N} = k_B \left(\frac{1}{2} \ln \frac{\pi}{\sigma^2} + \frac{r}{2} \ln \frac{8\pi^2 k_B T}{h^2} + \frac{1}{2} \sum_{i=1}^r \ln I_i + \frac{r}{2} \right) = \frac{S_{rot}}{N} \quad (61)$$

$$S_{m,vib,cl} = \frac{\partial S_{vib,cl}}{\partial N} = k_B \left(f + \sum_{i=1}^f \ln \frac{k_B T}{h\nu_i} \right) = \frac{S_{vib,cl}}{N} \quad (62)$$

$$S_{m,vib,qm} = \frac{\partial S_{vib,qm}}{\partial N} = k_B \sum_{i=1}^f \left[\frac{\frac{h\nu}{2k_B T}}{\tanh \frac{h\nu}{2k_B T}} - \ln \left(2 \sinh \frac{h\nu}{2k_B T} \right) \right] = \frac{S_{vib,qm}}{N} \quad (63)$$

$$S_{m,elec} = \frac{\partial S_{elec}}{\partial N} = 0 \quad (64)$$

References

- (1) Conti, S.; Cecchini, M. Predicting Molecular Self-Assembly at Surfaces: A Statistical Thermodynamics and Modeling Approach. *Physical Chemistry Chemical Physics* **2016**, *18*, 1480–31493.
- (2) Tidor, B.; Karplus, M. The Contribution of Vibrational Entropy to Molecular Association: The Dimerization of Insulin. *Journal of molecular biology* **1994**, *238* (3), 405–414.
- (3) Cecchini, M. Quantum Corrections to the Free Energy Difference Between Peptides and Proteins Conformers. *Journal of Chemical Theory and Computation* **2015**, *11* (9), 4011–4022.
- (4) McQuarrie, D. A. *Statistical Mechanics*; University Science Books, 2000.