

# ETC5410: Nonparametric smoothing methods

July 2008



MONASH University

Rob J Hyndman

<http://www.robjhyndman.com/>

# Outline

- 1 Density estimation
- 2 Kernel regression
- 3 **Splines**
- 4 Additive models
- 5 Functional data analysis

# ETC5410: Nonparametric smoothing methods


## 3. Splines

- 1 Interpolating splines
- 2 Smoothing splines
- 3 Regression splines
- 4 Penalized regression splines
- 5 Other bases

# Outline

- 1 **Interpolating splines**
- 2 Smoothing splines
- 3 Regression splines
- 4 Penalized regression splines
- 5 Other bases

# Interpolating splines



# Interpolating splines

Observations:  $(x_i, y_i)$  for  $i = 1, \dots, n$ .

# Interpolating splines

Observations:  $(x_i, y_i)$  for  $i = 1, \dots, n$ .

A spline is a continuous function  $f(x)$  interpolating all points and consisting of polynomials between each consecutive pair of 'knots'  $x_i$  and  $x_{i+1}$ .

# Interpolating splines

Observations:  $(x_i, y_i)$  for  $i = 1, \dots, n$ .

A spline is a continuous function  $f(x)$  interpolating all points and consisting of polynomials between each consecutive pair of 'knots'  $x_i$  and  $x_{i+1}$ .

- Parameters constrained so that  $f(x)$  is continuous.



# Interpolating splines


Observations:  $(x_i, y_i)$  for  $i = 1, \dots, n$ .

A spline is a continuous function  $f(x)$  interpolating all points and consisting of polynomials between each consecutive pair of 'knots'  $x_i$  and  $x_{i+1}$ .

- Parameters constrained so that  $f(x)$  is continuous.
- Further constraints imposed to give continuous derivatives.


# Interpolating splines

Linear interpolating spline




# Interpolating splines

Linear interpolating spline




# Interpolating splines

Quadratic interpolating spline




# Interpolating splines

Quadratic interpolating spline




# Interpolating splines

Quadratic interpolating spline




# Interpolating splines

Quadratic interpolating spline



# Interpolating splines


Quadratic interpolating spline





# Interpolating splines

Cubic interpolating spline



# Interpolating splines

- To define a cubic spline uniquely, it is necessary to specify two additional constraints.

# Interpolating splines

- To define a cubic spline uniquely, it is necessary to specify two additional constraints.
- Usually, this is done by specifying the form of the function at the two extremes,  $x_1$  and  $x_n$ .

# Interpolating splines

- To define a cubic spline uniquely, it is necessary to specify two additional constraints.
- Usually, this is done by specifying the form of the function at the two extremes,  $x_1$  and  $x_n$ .
- For example, a “natural” spline is obtained by requiring  $f''(x_1) = f''(x_n) = 0$ , thus making the curve linear at the extremes.

# Interpolating splines

- To define a cubic spline uniquely, it is necessary to specify two additional constraints.
- Usually, this is done by specifying the form of the function at the two extremes,  $x_1$  and  $x_n$ .
- For example, a “natural” spline is obtained by requiring  $f''(x_1) = f''(x_n) = 0$ , thus making the curve linear at the extremes.

# Interpolating splines

- To define a cubic spline uniquely, it is necessary to specify two additional constraints.
- Usually, this is done by specifying the form of the function at the two extremes,  $x_1$  and  $x_n$ .
- For example, a “natural” spline is obtained by requiring  $f''(x_1) = f''(x_n) = 0$ , thus making the curve linear at the extremes.

## Implementation in R

```
plot(x,y)  
lines(spline(x,y))
```

# Monotonic interpolation

‘Hyman filter’ ensures spline is monotonic by constraining derivatives while (possibly) sacrificing smoothness.

# Monotonic interpolation

'Hyman filter' ensures spline is monotonic by constraining derivatives while (possibly) sacrificing smoothness.

Second derivative may no longer be continuous at all knots.



# Monotonic interpolation

'Hyman filter' ensures spline is monotonic by constraining derivatives while (possibly) sacrificing smoothness.

Second derivative may no longer be continuous at all knots.


## Implementation in R

```
require(demography)
plot(x,y)
lines(cm.spline(x,y), col=4)
```

Reference: Smith, Hyndman and Wood (JPR, 2001)

# Monotonic interpolation

Cubic monotonic interpolating spline



# Outline

- 1 Interpolating splines
- 2 Smoothing splines**
- 3 Regression splines
- 4 Penalized regression splines
- 5 Other bases

# Spline smoothing

The average squared prediction error is

$$\sum [y_j - r(x_j)]^2.$$

Reduces to zero if  $r$  interpolates the data.

# Spline smoothing

The average squared prediction error is

$$\sum [y_j - r(x_j)]^2.$$

Reduces to zero if  $r$  interpolates the data.

Penalized sum of squares

$$S_\lambda(r) = \sum_{j=1}^n [y_j - r(x_j)]^2 + \lambda \int (r''(x))^2 dx$$

# Spline smoothing

The average squared prediction error is

$$\sum [y_j - r(x_j)]^2.$$

Reduces to zero if  $r$  interpolates the data.

Penalized sum of squares

$$S_\lambda(r) = \sum_{j=1}^n [y_j - r(x_j)]^2 + \lambda \int (r''(x))^2 dx$$

- $\lambda$  denotes a smoothing parameter.

# Spline smoothing

The average squared prediction error is

$$\sum [y_j - r(x_j)]^2.$$

Reduces to zero if  $r$  interpolates the data.

Penalized sum of squares

$$S_\lambda(r) = \sum_{j=1}^n [y_j - r(x_j)]^2 + \lambda \int (r''(x))^2 dx$$

- $\lambda$  denotes a smoothing parameter.
- First term measures closeness to the data.

# Spline smoothing

The average squared prediction error is

$$\sum [y_j - r(x_j)]^2.$$

Reduces to zero if  $r$  interpolates the data.

Penalized sum of squares

$$S_\lambda(r) = \sum_{j=1}^n [y_j - r(x_j)]^2 + \lambda \int (r''(x))^2 dx$$

- $\lambda$  denotes a smoothing parameter.
- First term measures closeness to the data.
- Second term penalizes curvature in the function.



# Cubic smoothing splines

A *cubic smoothing spline* is the function  $\hat{r}_\lambda(x)$  which minimizes  $S_\lambda(r)$  over the class of all twice differentiable functions on the range of  $\{x_j\}$ .

# Cubic smoothing splines

A *cubic smoothing spline* is the function  $\hat{r}_\lambda(x)$  which minimizes  $S_\lambda(r)$  over the class of all twice differentiable functions on the range of  $\{x_j\}$ .

- It consists of piecewise cubic polynomials, with the pieces separated by the  $x_j$  values.

# Cubic smoothing splines

A *cubic smoothing spline* is the function  $\hat{r}_\lambda(x)$  which minimizes  $S_\lambda(r)$  over the class of all twice differentiable functions on the range of  $\{x_j\}$ .

- It consists of piecewise cubic polynomials, with the pieces separated by the  $x_j$  values.
- At the design points,  $x_j$ ,  $\hat{r}_\lambda(x)$  and its first two derivatives are continuous. The third derivative may be discontinuous.

# Cubic smoothing splines

A *cubic smoothing spline* is the function  $\hat{r}_\lambda(x)$  which minimizes  $S_\lambda(r)$  over the class of all twice differentiable functions on the range of  $\{x_j\}$ .

- It consists of piecewise cubic polynomials, with the pieces separated by the  $x_j$  values.
- At the design points,  $x_j$ ,  $\hat{r}_\lambda(x)$  and its first two derivatives are continuous. The third derivative may be discontinuous.
- At the minimum and maximum  $x_j$  values, the second derivative of  $\hat{r}_\lambda(x)$  is zero. Hence, the smoothing spline is linear beyond the extreme data points.

# Cubic smoothing splines

- Large values of  $\lambda$  produce smoother curves while smaller values produce rougher curves.


# Cubic smoothing splines

- Large values of  $\lambda$  produce smoother curves while smaller values produce rougher curves.
- At  $\lambda \rightarrow \infty$ , the penalty term dominates  $S_\lambda(r)$ , forcing  $r''(x) = 0$  for all  $x$ . So the solution is the least squares straight line.

# Cubic smoothing splines


- Large values of  $\lambda$  produce smoother curves while smaller values produce rougher curves.
- At  $\lambda \rightarrow \infty$ , the penalty term dominates  $S_\lambda(r)$ , forcing  $r''(x) = 0$  for all  $x$ . So the solution is the least squares straight line.
- As  $\lambda \rightarrow 0$ , the penalty term becomes negligible and the solution tends to an interpolating function which is twice differentiable.

# Cubic smoothing splines







# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines



# Cubic smoothing splines




# Cubic smoothing splines







# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines



# Cubic smoothing splines




# Cubic smoothing splines







# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines




# Cubic smoothing splines



# Cubic smoothing splines



# Cubic smoothing splines





# Cubic smoothing splines

## Implementation in R

```
plot(price, shipments)
lines(smooth.spline(price, shipments))
lines(smooth.spline(price, shipments,
                    df=5), col=2)
```

# Cubic smoothing splines

## Implementation in R

```
plot(price, shipments)
lines(smooth.spline(price, shipments))
lines(smooth.spline(price, shipments,
  df=5), col=2)
```

- Fits a cubic smoothing spline to the data.

# Cubic smoothing splines

## Implementation in R

```
plot(price, shipments)
lines(smooth.spline(price, shipments))
lines(smooth.spline(price, shipments,
  df=5), col=2)
```

- Fits a cubic smoothing spline to the data.
- The value of  $\lambda$  is selected automatically using cross-validation.

# Cubic smoothing splines

## Implementation in R

```
plot(price, shipments)
lines(smooth.spline(price, shipments))
lines(smooth.spline(price, shipments,
  df=5), col=2)
```

- Fits a cubic smoothing spline to the data.
- The value of  $\lambda$  is selected automatically using cross-validation.
- The argument `df` can be supplied and then  $\lambda$  is chosen to give approximately `df` degrees of freedom.

# Hodrick-Prescott filter

- The Hodrick-Prescott filter is a special case of a cubic smoothing spline.

# Hodrick-Prescott filter

- The Hodrick-Prescott filter is a special case of a cubic smoothing spline.
- HP recommended  $\lambda = 1600$  for quarterly data. There is no theoretical justification for this.

# Hodrick-Prescott filter

- The Hodrick-Prescott filter is a special case of a cubic smoothing spline.
- HP recommended  $\lambda = 1600$  for quarterly data. There is no theoretical justification for this.
- Better to use proper bandwidth selection tools.

# Cross-validation again

Recall: Find smoothing parameter which minimises

$$CV(h) = \frac{1}{n} \sum_{j=1}^n [\hat{r}_j(x_j) - y_j]^2$$

where  $\hat{r}_j(x_j)$  uses all data *except*  $(x_j, y_j)$ .




# Cross-validation again


Recall: Find smoothing parameter which minimises

$$CV(h) = \frac{1}{n} \sum_{j=1}^n [\hat{r}_j(x_j) - y_j]^2$$


where  $\hat{r}_j(x_j)$  uses all data *except*  $(x_j, y_j)$ .



# Cross-validation again



# Cross-validation again



# General cubic splines

Let  $\kappa_1 < \kappa_2 < \cdots < \kappa_k$  be knots in interval  $(a, b)$ .

# General cubic splines

Let  $\kappa_1 < \kappa_2 < \dots < \kappa_k$  be knots in interval  $(a, b)$ .

Let  $h_1(x) = 1$ ,  $h_2(x) = x$ ,  $h_3(x) = x^2$ ,  $h_4(x) = x^3$ ,  
 $h_j(x) = (x - \kappa_{j-4})_+^3$  for  $j = 5, \dots, k + 4$ .

# General cubic splines

Let  $\kappa_1 < \kappa_2 < \dots < \kappa_k$  be knots in interval  $(a, b)$ .

Let  $h_1(x) = 1$ ,  $h_2(x) = x$ ,  $h_3(x) = x^2$ ,  $h_4(x) = x^3$ ,  
 $h_j(x) = (x - \kappa_{j-4})_+^3$  for  $j = 5, \dots, k+4$ .

$\{h_1, \dots, h_{k+4}\}$  form a basis for the set of cubic splines at these knots, called the **truncated power basis**.

# General cubic splines

Let  $\kappa_1 < \kappa_2 < \dots < \kappa_k$  be knots in interval  $(a, b)$ .

Let  $h_1(x) = 1$ ,  $h_2(x) = x$ ,  $h_3(x) = x^2$ ,  $h_4(x) = x^3$ ,  
 $h_j(x) = (x - \kappa_{j-4})_+^3$  for  $j = 5, \dots, k+4$ .

$\{h_1, \dots, h_{k+4}\}$  form a basis for the set of cubic splines at these knots, called the **truncated power basis**.

Any cubic spline  $r(x)$  with these knots can be written as

$$r(x) = \sum_{j=1}^{k+4} \beta_j h_j(x).$$

# General cubic splines

Let  $\kappa_1 < \kappa_2 < \dots < \kappa_k$  be knots in interval  $(a, b)$ .

Let  $h_1(x) = 1$ ,  $h_2(x) = x$ ,  $h_3(x) = x^2$ ,  $h_4(x) = x^3$ ,  
 $h_j(x) = (x - \kappa_{j-4})_+^3$  for  $j = 5, \dots, k+4$ .

$\{h_1, \dots, h_{k+4}\}$  form a basis for the set of cubic splines at these knots, called the **truncated power basis**.

Any cubic spline  $r(x)$  with these knots can be written as

$$r(x) = \sum_{j=1}^{k+4} \beta_j h_j(x).$$

A cubic smoothing spline is obtained by setting  $\kappa_j = x_j$ ,  
 $j = 1, \dots, n$ .



# Matrix form

Let  $B_{ij} = h_j(x_i)$  be the basis matrix. Then for smoothing splines we need to minimize

$$(Y - B\beta)'(Y - B\beta) + \lambda\beta'\Omega\beta$$

where

$$\Omega_{jk} = \int h_j''(x)h_k''(x)dx.$$

# Matrix form

Let  $B_{ij} = h_j(x_i)$  be the basis matrix. Then for smoothing splines we need to minimize

$$(Y - B\beta)'(Y - B\beta) + \lambda\beta'\Omega\beta$$

where

$$\Omega_{jk} = \int h_j''(x)h_k''(x)dx.$$

This gives  $\hat{\beta} = (B'B + \lambda\Omega)^{-1}B'Y$ .

# Inference

Any cubic smoothing spline can be written as a linear smoother with smoothing matrix

$$\mathbf{S} = B(B'B + \lambda\Omega)^{-1}B'.$$

# Inference

Any cubic smoothing spline can be written as a linear smoother with smoothing matrix

$$\mathbf{S} = B(B'B + \lambda\Omega)^{-1}B'.$$

$$\hat{\mathbf{r}} = \mathbf{S}\mathbf{Y} \text{ where } \hat{\mathbf{r}} = [\hat{r}(x_1), \dots, \hat{r}(x_n)]'$$

# Inference

Any cubic smoothing spline can be written as a linear smoother with smoothing matrix

$$\mathbf{S} = B(B'B + \lambda\Omega)^{-1}B'.$$

$$\hat{\mathbf{r}} = \mathbf{S}\mathbf{Y} \text{ where } \hat{\mathbf{r}} = [\hat{r}(x_1), \dots, \hat{r}(x_n)]'$$

- The effect of  $\lambda\Omega$  is to shrink the regression coefficients to give a smoother fit.

# Inference

Any cubic smoothing spline can be written as a linear smoother with smoothing matrix

$$\mathbf{S} = B(B'B + \lambda\Omega)^{-1}B'.$$

$$\hat{\mathbf{r}} = \mathbf{S}\mathbf{Y} \text{ where } \hat{\mathbf{r}} = [\hat{r}(x_1), \dots, \hat{r}(x_n)]'$$

- The effect of  $\lambda\Omega$  is to shrink the regression coefficients to give a smoother fit.
- Because this is in the form of a linear smoother, the inference derived for local polynomial smoothing can be applied here too.

# Inference

Any cubic smoothing spline can be written as a linear smoother with smoothing matrix

$$\mathbf{S} = B(B'B + \lambda\Omega)^{-1}B'.$$

$$\hat{\mathbf{r}} = \mathbf{S}\mathbf{Y} \text{ where } \hat{\mathbf{r}} = [\hat{r}(x_1), \dots, \hat{r}(x_n)]'$$

- The effect of  $\lambda\Omega$  is to shrink the regression coefficients to give a smoother fit.
- Because this is in the form of a linear smoother, the inference derived for local polynomial smoothing can be applied here too.
- $df = \text{trace}(\mathbf{S})$

# Outline

- 1 Interpolating splines
- 2 Smoothing splines
- 3 Regression splines**
- 4 Penalized regression splines
- 5 Other bases



# Regression splines

- Rather than have knots at each data point ( $\kappa_i = x_i$ ), use fewer knots.

# Regression splines

- Rather than have knots at each data point ( $\kappa_i = x_i$ ), use fewer knots.
- Then do ordinary linear regression on the basis function.

# Regression splines

- Rather than have knots at each data point ( $\kappa_i = x_i$ ), use fewer knots.
- Then do ordinary linear regression on the basis function.
- Choice of knots can be difficult and arbitrary.

# Regression splines

- Rather than have knots at each data point ( $\kappa_i = x_i$ ), use fewer knots.
- Then do ordinary linear regression on the basis function.
- Choice of knots can be difficult and arbitrary.
- Automatic knot selection algorithms very slow.

# Regression splines

- Rather than have knots at each data point ( $\kappa_i = x_i$ ), use fewer knots.
- Then do ordinary linear regression on the basis function.
- Choice of knots can be difficult and arbitrary.
- Automatic knot selection algorithms very slow.

# Regression splines

- Rather than have knots at each data point ( $\kappa_i = x_i$ ), use fewer knots.
- Then do ordinary linear regression on the basis function.
- Choice of knots can be difficult and arbitrary.
- Automatic knot selection algorithms very slow.

$$\hat{\beta} = (B'B)^{-1}B'\mathbf{Y}.$$

# Outline

- 1 Interpolating splines
- 2 Smoothing splines
- 3 Regression splines
- 4 Penalized regression splines**
- 5 Other bases

# Penalized spline regression

$$r(x) = \sum_{j=1}^K \beta_j h_j(x).$$

**Idea:** retain all the knots but constrain their influence by

$$\sum \beta_j^2 < C.$$



# Penalized spline regression

$$r(x) = \sum_{j=1}^K \beta_j h_j(x).$$

**Idea:** retain all the knots but constrain their influence by

$$\sum \beta_j^2 < C.$$

Let

$$D = \begin{bmatrix} \mathbf{0}_{2 \times 2} & \mathbf{0}_{2 \times K} \\ \mathbf{0}_{K \times 2} & \mathbf{I}_{K \times K} \end{bmatrix}.$$

Then we want to

minimize  $\|\mathbf{y} - \mathbf{B}\boldsymbol{\beta}\|^2$  subject to  $\boldsymbol{\beta}'\mathbf{D}\boldsymbol{\beta} \leq C$ .

# Penalized regression splines

A Lagrange multiplier argument shows that this is equivalent to minimizing

$$\|\mathbf{y} - \mathbf{B}\boldsymbol{\beta}\|^2 + \lambda^2 \boldsymbol{\beta}' \mathbf{D} \boldsymbol{\beta}$$


for some number  $\lambda \geq 0$ .

**Solution:**  $\hat{\boldsymbol{\beta}}_{\lambda} = (\mathbf{X}'\mathbf{X} + \lambda^2 \mathbf{D})^{-1} \mathbf{X}'\mathbf{y}$ .

**Fitted values:**  $\hat{\mathbf{y}} = \mathbf{X}(\mathbf{X}'\mathbf{X} + \lambda^2 \mathbf{D})^{-1} \mathbf{X}'\mathbf{y}$ .


- A type of ridge regression.

# Penalized regression splines







# Penalized regression splines




# Penalized regression splines



# Penalized regression splines




# Penalized regression splines







# Penalized regression splines




# Penalized regression splines




# Penalized regression splines



# Penalized regression splines



# Penalized regression splines



# Penalized regression splines

## Implementation in R

```
require(SemiPar)
fit <- spm(shipments ~ f(price))
plot(fit)
points(price, shipments)
```

# Penalized regression splines

## Implementation in R

```
require(SemiPar)
fit <- spm(shipments ~ f(price))
plot(fit)
points(price, shipments)
```

For lots of examples and an introduction to the theory:

<http://www.uow.edu.au/~mwand/SPmanu.pdf>

# Outline

- 1 Interpolating splines
- 2 Smoothing splines
- 3 Regression splines
- 4 Penalized regression splines
- 5 Other bases**



# Spline bases

## Truncated power basis of degree $p$

$$1, x, \dots, x^p, (x - \kappa_1)_+^p, \dots, (x - \kappa_K)_+^p$$

# Spline bases

## Truncated power basis of degree $p$

$1, x, \dots, x^p, (x - \kappa_1)_+^p, \dots, (x - \kappa_K)_+^p$

- $p - 1$  continuous derivatives

# Spline bases

## Truncated power basis of degree $p$

$1, x, \dots, x^p, (x - \kappa_1)_+^p, \dots, (x - \kappa_K)_+^p$

- $p - 1$  continuous derivatives
- In penalized regression splines, none of the polynomial coefficients is penalized.

# Spline bases

## Truncated power basis of degree $p$

$1, x, \dots, x^p, (x - \kappa_1)_+^p, \dots, (x - \kappa_K)_+^p$

- $p - 1$  continuous derivatives
- In penalized regression splines, none of the polynomial coefficients is penalized.


# Spline bases

## Truncated power basis of degree $p$

$1, x, \dots, x^p, (x - \kappa_1)_+^p, \dots, (x - \kappa_K)_+^p$

- $p - 1$  continuous derivatives
- In penalized regression splines, none of the polynomial coefficients is penalized.

### Truncated linear spline basis




# Spline bases

## Truncated power basis of degree $p$

$$1, x, \dots, x^p, (x - \kappa_1)_+^p, \dots, (x - \kappa_K)_+^p$$

- $p - 1$  continuous derivatives
- In penalized regression splines, none of the polynomial coefficients is penalized.

### Truncated cubic spline basis



# Spline bases

## B-splines


- *Equivalent* to truncated power bases but with more stable properties.

# Spline bases

## B-splines

- *Equivalent* to truncated power bases but with more stable properties.

**Cubic B-spline basis**





# Spline bases

## Radial basis functions of degree $p$

$$1, x, \dots, x^p, |x - \kappa_1|^p, \dots, |x - \kappa_K|^p$$

# Spline bases

## Radial basis functions of degree $p$

$1, x, \dots, x^p, |x - \kappa_1|^p, \dots, |x - \kappa_K|^p$

- $p - 1$  continuous derivatives

# Spline bases

## Radial basis functions of degree $p$

$1, x, \dots, x^p, |x - \kappa_1|^p, \dots, |x - \kappa_K|^p$

- $p - 1$  continuous derivatives
- In penalized regression splines, none of the polynomial coefficients is penalized.

# Spline bases

## Radial basis functions of degree $p$

$1, x, \dots, x^p, |x - \kappa_1|^p, \dots, |x - \kappa_K|^p$

- $p - 1$  continuous derivatives
- In penalized regression splines, none of the polynomial coefficients is penalized.

# Spline bases

## Radial basis functions of degree $p$

$1, x, \dots, x^p, |x - \kappa_1|^p, \dots, |x - \kappa_K|^p$

- $p - 1$  continuous derivatives
- In penalized regression splines, none of the polynomial coefficients is penalized.

