

Introduction to Differential-Algebraic Equations (DAEs)

Sébastien Gros
NTNU, Eng. Cybernetics

Slides for TTK4130
2021

Objectives of the slides

Learn the basics of DAEs

- ✓ understand what a DAE is
- ✓ identify the different forms of DAEs
- ✓ understand the Tikhonov theorem
- ✓ understand why there are “easy” and “hard” DAEs
- ✓ understand differential index & index reductions

Simulating ODEs

Most common form of ODE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u}) \quad (1)$$

... delivers $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given

State derivative $\dot{\mathbf{x}}$ tells us “where” the state is going from \mathbf{x}, \mathbf{u}

Simulating ODEs

Most common form of ODE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u}) \quad (1)$$

... delivers $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given

State derivative $\dot{\mathbf{x}}$ tells us “where” the state is going from \mathbf{x}, \mathbf{u}

Then we “can” simulate (1), i.e. “follow” $\dot{\mathbf{x}}$
wherever it leads

Simulating ODEs


Most common form of ODE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u}) \quad (1)$$

... delivers $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given

State derivative $\dot{\mathbf{x}}$ tells us “where” the state is going from \mathbf{x}, \mathbf{u}

Then we “can” simulate (1), i.e. “follow” $\dot{\mathbf{x}}$
wherever it leads



Simulating ODEs

Most common form of ODE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u}) \quad (1)$$

... delivers $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given


State derivative $\dot{\mathbf{x}}$ tells us “where” the state is going from \mathbf{x}, \mathbf{u}

Then we “can” simulate (1), i.e. “follow” $\dot{\mathbf{x}}$ wherever it leads

Implicit ODEs

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$$

...simulation similarly requires one to be able to compute $\dot{\mathbf{x}}$ for any \mathbf{x}, \mathbf{u}



Implicit ODEs

How to “solve” an implicit ODE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

Implicit ODEs

How to “solve” an implicit ODE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

- In order to simulate (2) we need $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given)

Implicit ODEs

How to “solve” an implicit ODE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

- In order to simulate (2) we need $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given)
- Equation (2) provides $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given) **implicitly**

Implicit ODEs

How to “solve” an implicit ODE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

- In order to simulate (2) we need $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given)
- Equation (2) provides $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given) **implicitly**
- In order to get $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given, we need to **solve** (2) for $\dot{\mathbf{x}}$

Implicit ODEs

How to “solve” an implicit ODE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

- In order to simulate (2) we need $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given)
- Equation (2) provides $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given) **implicitly**
- In order to get $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given, we need to **solve** (2) for $\dot{\mathbf{x}}$
- Pen-and-paper solution may not exist! E.g. try $e^{\dot{x}} + \dot{x}^3 - u = 0$
 - ▶ has a well-defined solution \dot{x} for any u
 - ▶ cannot write an expression $\dot{x} = \dots$

Implicit ODEs

How to “solve” an implicit ODE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

- In order to simulate (2) we need $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given)
- Equation (2) provides $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given) **implicitly**
- In order to get $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given, we need to **solve** (2) for $\dot{\mathbf{x}}$
- Pen-and-paper solution may not exist! E.g. try $e^{\dot{x}} + \dot{x}^3 - u = 0$
 - ▶ has a well-defined solution \dot{x} for any u
 - ▶ cannot write an expression $\dot{x} = \dots$
- Numerical methods are often needed. More on that soon!

Implicit ODEs

How to “solve” an implicit ODE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

- In order to simulate (2) we need $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given)
- Equation (2) provides $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given) **implicitly**
- In order to get $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given, we need to **solve** (2) for $\dot{\mathbf{x}}$
- Pen-and-paper solution may not exist! E.g. try $e^{\dot{x}} + \dot{x}^3 - u = 0$
 - ▶ has a well-defined solution \dot{x} for any u
 - ▶ cannot write an expression $\dot{x} = \dots$
- Numerical methods are often needed. More on that soon!
- But **if** (2) defines $\dot{\mathbf{x}}$ for any \mathbf{x}, \mathbf{u} , then *there exists a function \mathbf{f} such that:*

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u})$$

even if we cannot *write* \mathbf{f} in terms of *mathematical expressions* (e.g. $\sin, \cos, .^2$),

Implicit ODEs

How to “solve” an implicit ODE

$$F(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (2)$$

- In order to simulate (2) we need $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given)
- Equation (2) provides $\dot{\mathbf{x}}$ (for \mathbf{x}, \mathbf{u} given) **implicitly**
- In order to get $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given, we need to **solve** (2) for $\dot{\mathbf{x}}$
- Pen-and-paper solution may not exist! E.g. try $e^{\dot{x}} + \dot{x}^3 - u = 0$
 - ▶ has a well-defined solution \dot{x} for any u
 - ▶ cannot write an expression $\dot{x} = \dots$
- Numerical methods are often needed. More on that soon!
- But **if** (2) defines $\dot{\mathbf{x}}$ for any \mathbf{x}, \mathbf{u} , then *there exists a function \mathbf{f} such that:*

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u})$$

even if we cannot *write* \mathbf{f} in terms of *mathematical expressions* (e.g. $\sin, \cos, .^2$),

- Then

$$F(\mathbf{f}(\mathbf{x}, \mathbf{u}), \mathbf{x}, \mathbf{u}) = 0, \quad \text{for all } \mathbf{x}, \mathbf{u}$$

Implicit ODEs

How to “solve” an implicit ODE

$$F(\dot{x}, x, u) = 0 \quad (2)$$

- In order to simulate (2) we need \dot{x} (for x, u given)
- Equation (2) provides \dot{x} (for x, u given) **implicitly**
- In order to get \dot{x} for x, u given, we need to **solve** (2) for \dot{x}
- Pen-and-paper solution may not exist! E.g. try $e^{\dot{x}} + \dot{x}^3 - u = 0$
 - ▶ has a well-defined solution \dot{x} for any u
 - ▶ cannot write an expression $\dot{x} = \dots$
- Numerical methods are often needed. More on that soon!
- But **if** (2) defines \dot{x} for any x, u , then *there exists a function f such that:*

$$\dot{x} = f(x, u)$$

even if we cannot *write* f in terms of *mathematical expressions* (e.g. \sin, \cos, \dots),

- Then

$$F(f(x, u), x, u) = 0, \quad \text{for all } x, u$$

When does (2) define \dot{x} properly? I.e. when does f exist?

Solving ODEs - Implicit Function Theorem

Consider function $f(x)$ implicitly given by:

$$F(f(x), x) = 0$$

... when is $f(x)$ well defined?

Solving ODEs - Implicit Function Theorem

Consider function $f(x)$ implicitly given by:

$$F(f(x), x) = 0$$

... when is $f(x)$ well defined?

Implicit Function Theorem (simplified version)

Let $F(y, x)$ be smooth

Solving ODEs - Implicit Function Theorem

Consider function $f(x)$ implicitly given by:

$$F(f(x), x) = 0$$

... when is $f(x)$ well defined?

Implicit Function Theorem (simplified version)

Let $F(y, x)$ be smooth, and suppose that

$$\frac{\partial F(y, x)}{\partial y} \text{ is full rank for all } y, x \text{ s.t. } F(y, x) = 0$$

Solving ODEs - Implicit Function Theorem

Consider function $f(x)$ implicitly given by:

$$F(f(x), x) = 0$$

... when is $f(x)$ well defined?

Implicit Function Theorem (simplified version)

Let $F(y, x)$ be smooth, and suppose that

$$\frac{\partial F(y, x)}{\partial y} \text{ is full rank for all } y, x \text{ s.t. } F(y, x) = 0$$

then there exists $f(x)$ such that

$$F(f(x), x) = 0$$

Solving ODEs - Implicit Function Theorem

Consider function $f(x)$ implicitly given by:

$$F(f(x), x) = 0$$

... when is $f(x)$ well defined?

Implicit Function Theorem (simplified version)

Let $F(y, x)$ be smooth, and suppose that

$$\frac{\partial F(y, x)}{\partial y} \text{ is full rank for all } y, x \text{ s.t. } F(y, x) = 0$$

then there exists $f(x)$ such that

$$F(f(x), x) = 0$$

Solving ODEs - Implicit Function Theorem

Consider function $f(x)$ implicitly given by:

$$F(f(x), x) = 0$$

... when is $f(x)$ well defined?

Implicit Function Theorem a “practical version”

If $F(y, x)$ is smooth, and

$$\frac{\partial F(y, x)}{\partial y} \text{ is always full rank}$$

... then $f(x)$ exists.

Solving ODEs - Implicit Function Theorem (cont')

Can we solve

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (3)$$

... for $\dot{\mathbf{x}}$?

Solving ODEs - Implicit Function Theorem (cont')

Can we solve

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (3)$$

... for $\dot{\mathbf{x}}$?

What about e.g. $u\dot{x}^2 - x = 0$?

Solving ODEs - Implicit Function Theorem (cont')

Can we solve

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (3)$$

... for $\dot{\mathbf{x}}$?

What about e.g. $u\dot{x}^2 - x = 0$?

$$\dot{x} = \pm \sqrt{\frac{x}{u}}, \quad \text{if } u \neq 0$$

Solving ODEs - Implicit Function Theorem (cont')

Can we solve

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (3)$$

... for $\dot{\mathbf{x}}$?

What about e.g. $u\dot{x}^2 - x = 0$?

$$\dot{x} = \pm \sqrt{\frac{x}{u}}, \quad \text{if } u \neq 0$$

Use IFT:

Equation (3) can be solved for $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given if:

$$\frac{\partial \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u})}{\partial \dot{\mathbf{x}}}$$

... is always full rank

Solving ODEs - Implicit Function Theorem (cont')

Can we solve

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (3)$$

... for $\dot{\mathbf{x}}$?

What about e.g. $u\dot{x}^2 - x = 0$?

$$\dot{x} = \pm \sqrt{\frac{x}{u}}, \quad \text{if } u \neq 0$$

Use IFT:

Equation (3) can be solved for $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given if:

$$\frac{\partial \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u})}{\partial \dot{\mathbf{x}}}$$

... is always full rank

What about e.g. $\mathbf{F}(\dot{x}, x, u) = u\dot{x}^2 - x = 0$?

$$\frac{\partial \mathbf{F}}{\partial \dot{x}} = 2\dot{x}u$$

... this is “full rank” (i.e. $\neq 0$ here) if $u \neq 0$ and $\dot{x} \neq 0$

Solving ODEs - Implicit Function Theorem (cont')

Can we solve

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0 \quad (3)$$

... for $\dot{\mathbf{x}}$?

What about e.g. $u\dot{x}^2 - x = 0$?

$$\dot{x} = \pm \sqrt{\frac{x}{u}}, \quad \text{if } u \neq 0$$

Use IFT:

Equation (3) can be solved for $\dot{\mathbf{x}}$ for \mathbf{x}, \mathbf{u} given if:

$$\frac{\partial \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u})}{\partial \dot{\mathbf{x}}}$$

... is always full rank

What about e.g. $\mathbf{F}(\dot{x}, x, u) = u\dot{x}^2 - x = 0$?

$$\frac{\partial \mathbf{F}}{\partial \dot{x}} = 2\dot{x}u$$

... this is “full rank” (i.e. $\neq 0$ here) if $u \neq 0$ and $\dot{x} \neq 0$

Now we know when an ODE has a “well-posed $\dot{\mathbf{x}}$ ”

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

E.g.

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad (4)$$

State is

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix}$$

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

E.g.

State is

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad (4)$$

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix}$$

because $\dot{\mathbf{x}}_2$ does not appear, (4) does not deliver $\dot{\mathbf{x}}_2$...

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

E.g.

State is

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad (4)$$

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix}$$

because $\dot{\mathbf{x}}_2$ does not appear, (4) does not deliver $\dot{\mathbf{x}}_2$...

Jacobian is rank deficient because:

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

→ (4) is a DAE

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

E.g.

State is

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad (4)$$

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix}$$

because $\dot{\mathbf{x}}_2$ does not appear, (4) does not deliver $\dot{\mathbf{x}}_2$...

... however (4) provides well-defined trajectories!

Jacobian is rank deficient because:

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

→ (4) is a DAE

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

E.g.

State is

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad (4)$$

$$\mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix}$$

because $\dot{\mathbf{x}}_2$ does not appear, (4) does not deliver $\dot{\mathbf{x}}_2$...

... however (4) provides well-defined trajectories!

Jacobian is rank deficient because: Actually (4) yields

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\dot{\mathbf{x}}_1 = \mathbf{x}_1$$

$$\mathbf{x}_2 = -\mathbf{x}_1 - u$$

→ (4) is a DAE

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

E.g.

State is

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad (4) \quad \mathbf{x} = \begin{bmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{bmatrix}$$

because $\dot{\mathbf{x}}_2$ does not appear, (4) does not deliver $\dot{\mathbf{x}}_2$...

... however (4) provides well-defined trajectories!

Jacobian is rank deficient because: Actually (4) yields

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \begin{aligned} \dot{\mathbf{x}}_1 &= -\mathbf{x}_2 - u \\ \dot{\mathbf{x}}_2 &= -\mathbf{x}_1 - u \end{aligned}$$

→ (4) is a DAE

In applications, DAEs are often differential equations where **some states derivatives do not appear** as in e.g. (4)

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 - \mathbf{x}_1 + \dot{\mathbf{x}}_2 \\ \dot{\mathbf{x}}_1 + \dot{\mathbf{x}}_2 + \mathbf{x}_2 + u \end{bmatrix} = 0$$

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 - \mathbf{x}_1 + \dot{\mathbf{x}}_2 \\ \dot{\mathbf{x}}_1 + \dot{\mathbf{x}}_2 + \mathbf{x}_2 + u \end{bmatrix} = 0$$

yields

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{x}_1 - x_1 + \dot{x}_2 \\ \dot{x}_1 + \dot{x}_2 + x_2 + u \end{bmatrix} = 0$$

yields

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

This is a DAE even though both \dot{x}_1 and \dot{x}_2 appear

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{x}_1 - x_1 + \dot{x}_2 \\ \dot{x}_1 + \dot{x}_2 + x_2 + u \end{bmatrix} = 0$$

yields

has solution

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$x_1 = -\dot{u}$$

$$x_2 = -x_1 - u$$

DAE well defined only for u continuous!

This is a DAE even though both \dot{x}_1 and \dot{x}_2 appear

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_1 - u \\ (\mathbf{x}_1 - \mathbf{x}_2) \dot{\mathbf{x}}_2 + \mathbf{x}_1 - \mathbf{x}_2 \end{bmatrix} = 0$$

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_1 - u \\ (\mathbf{x}_1 - \mathbf{x}_2) \dot{\mathbf{x}}_2 + \mathbf{x}_1 - \mathbf{x}_2 \end{bmatrix} = 0$$

yields

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 0 \\ 0 & \mathbf{x}_1 - \mathbf{x}_2 \end{bmatrix}$$

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_1 - u \\ (\mathbf{x}_1 - \mathbf{x}_2) \dot{\mathbf{x}}_2 + \mathbf{x}_1 - \mathbf{x}_2 \end{bmatrix} = 0$$

yields

For $\mathbf{x}_2(0) = \mathbf{x}_1(0)$, has solution

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 0 \\ 0 & \mathbf{x}_1 - \mathbf{x}_2 \end{bmatrix}$$

$$\dot{\mathbf{x}}_1 = u - \mathbf{x}_1$$

$$\mathbf{x}_2 = \mathbf{x}_1$$

Is it a DAE or an ODE?

What are DAEs?

Definition

An implicit differential equation $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$ is a DAE if $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ is rank deficient along the trajectory $\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}$.

A state does not appear time differentiated \rightarrow DAE, but also...

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_1 - u \\ (\mathbf{x}_1 - \mathbf{x}_2) \dot{\mathbf{x}}_2 + \mathbf{x}_1 - \mathbf{x}_2 \end{bmatrix} = 0$$

yields

For $\mathbf{x}_2(0) = \mathbf{x}_1(0)$, has solution

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} = \begin{bmatrix} 1 & 0 \\ 0 & \mathbf{x}_1 - \mathbf{x}_2 \end{bmatrix}$$

$$\dot{\mathbf{x}}_1 = u - \mathbf{x}_1$$

$$\mathbf{x}_2 = \mathbf{x}_1$$

Is it a DAE or an ODE? It can be both!

A differential equations can be both an ODE & DAE, even jump back-and-forth.
Avoided in practice, i.e. we like $\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}}$ having fixed rank

A bit of notation

- If some states do not appear time differentiated, we highlight them as “**z**”, e.g.

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad \longrightarrow \quad \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, u) = \begin{bmatrix} \dot{\mathbf{x}} + \mathbf{z} + u \\ \mathbf{x} + \mathbf{z} + u \end{bmatrix} = 0$$

A bit of notation

- If some states do not appear time differentiated, we highlight them as “**z**”, e.g.

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad \longrightarrow \quad \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, u) = \begin{bmatrix} \dot{\mathbf{x}} + \mathbf{z} + u \\ \mathbf{x} + \mathbf{z} + u \end{bmatrix} = 0$$

- Then the DAE definition “works” and is to be understood as:

$$\frac{\partial \mathbf{F}}{\partial \text{states}} = \begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix} = \begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \mathbf{0} \end{bmatrix} \quad \text{is rank deficient}$$

$$\text{because states} = \begin{bmatrix} \mathbf{x} \\ \mathbf{z} \end{bmatrix}$$

A bit of notation

- If some states do not appear time differentiated, we highlight them as “**z**”, e.g.

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad \longrightarrow \quad \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, u) = \begin{bmatrix} \dot{\mathbf{x}} + \mathbf{z} + u \\ \mathbf{x} + \mathbf{z} + u \end{bmatrix} = 0$$

- Then the DAE definition “works” and is to be understood as:

$$\frac{\partial \mathbf{F}}{\partial \text{states}} = \begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix} = \begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \mathbf{0} \end{bmatrix} \quad \text{is rank deficient}$$

$$\text{because states} = \begin{bmatrix} \mathbf{x} \\ \mathbf{z} \end{bmatrix}$$

- Fully-Implicit DAEs

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, u) = 0$$

A bit of notation

- If some states do not appear time differentiated, we highlight them as “z”, e.g.

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, u) = \begin{bmatrix} \dot{\mathbf{x}}_1 + \mathbf{x}_2 + u \\ \mathbf{x}_1 + \mathbf{x}_2 + u \end{bmatrix} = 0 \quad \longrightarrow \quad \mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, u) = \begin{bmatrix} \dot{\mathbf{x}} + \mathbf{z} + u \\ \mathbf{x} + \mathbf{z} + u \end{bmatrix} = 0$$

- Then the DAE definition “works” and is to be understood as:

$$\frac{\partial \mathbf{F}}{\partial \text{states}} = \begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix} = \begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \mathbf{0} \end{bmatrix} \quad \text{is rank deficient}$$

$$\text{because states} = \begin{bmatrix} \mathbf{x} \\ \mathbf{z} \end{bmatrix}$$

- Fully-Implicit DAEs

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$$

- Semi-explicit DAEs (most common form)

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

→ “explicit ODE + algebraic equations”

Conversion semi-explicit \leftrightarrow fully-implicit

Semi-explicit DAEs

Fully-Implicit DAEs

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

\longrightarrow

\longleftarrow

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

Conversion semi-explicit \leftrightarrow fully-implicit

Semi-explicit DAEs

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

\longrightarrow

Fully-Implicit DAEs

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = \begin{bmatrix} \dot{\mathbf{x}} - \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \\ \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \end{bmatrix} = 0$$

\longleftarrow

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

Conversion semi-explicit \leftrightarrow fully-implicit

Semi-explicit DAEs

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

\longrightarrow

Fully-Implicit DAEs

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = \begin{bmatrix} \dot{\mathbf{x}} - \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \\ \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \end{bmatrix} = 0$$

$$\dot{\mathbf{x}} = \mathbf{v}$$

$$0 = \mathbf{F}(\mathbf{v}, \mathbf{z}, \mathbf{x}, \mathbf{u})$$

\longleftarrow

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

Conversion semi-explicit \leftrightarrow fully-implicit

Semi-explicit DAEs

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

Fully-Implicit DAEs

$$\longrightarrow \quad \mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = \begin{bmatrix} \dot{\mathbf{x}} - \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \\ \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \end{bmatrix} = 0$$

$$\dot{\mathbf{x}} = \mathbf{v}$$

$$0 = \mathbf{F}(\mathbf{v}, \mathbf{z}, \mathbf{x}, \mathbf{u})$$


$$\longleftarrow \quad \mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

Fully-implicit or semi-explicit DAEs are not *really* different.

- we like semi-explicit DAEs for their neat structure
- fully-implicit \rightarrow semi-explicit adds variables \mathbf{v} , can be counter-productive

DAEs in mechanics - Example


Pendulum simulation



- Cartesian position $p \in \mathbb{R}^3$, unit mass

DAEs in mechanics - Example

Pendulum simulation




- Cartesian position $\mathbf{p} \in \mathbb{R}^3$, unit mass
- Lagrange function:

$$\mathcal{L} = \frac{1}{2} m \dot{\mathbf{p}}^\top \dot{\mathbf{p}} - mg\mathbf{p}_3 - \mathbf{z}\mathbf{c}(\mathbf{p})$$

DAEs in mechanics - Example

Pendulum simulation



- Cartesian position $\mathbf{p} \in \mathbb{R}^3$, unit mass
- Lagrange function:


$$\mathcal{L} = \frac{1}{2} m \dot{\mathbf{p}}^\top \dot{\mathbf{p}} - mg \mathbf{p}_3 - z\mathbf{c}(\mathbf{p})$$

- Constraint:

$$\mathbf{c}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2$$

DAEs in mechanics - Example

Pendulum simulation



- Cartesian position $\mathbf{p} \in \mathbb{R}^3$, unit mass
- Lagrange function:

$$\mathcal{L} = \frac{1}{2} m \dot{\mathbf{p}}^\top \dot{\mathbf{p}} - mg \mathbf{p}_3 - z \mathbf{c}(\mathbf{p})$$

- Constraint:


$$\mathbf{c}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2$$

- Motion:

$$m \ddot{\mathbf{p}} = -z \mathbf{p} - m \mathbf{g}, \quad \mathbf{g} = \begin{bmatrix} 0 \\ 0 \\ g \end{bmatrix}$$

DAEs in mechanics - Example

Pendulum simulation



Semi-explicit DAE:

$$\dot{p} = v$$

$$\dot{v} = -\frac{1}{m}z p - g$$

- Cartesian position $p \in \mathbb{R}^3$, unit mass
- Lagrange function:

$$\mathcal{L} = \frac{1}{2}m\dot{p}^\top \dot{p} - mgp_3 - zc(p)$$

- Constraint:


$$c(p) = p^\top p - L^2$$

- Motion:

$$m\ddot{p} = -z p - mg, \quad g = \begin{bmatrix} 0 \\ 0 \\ g \end{bmatrix}$$

DAEs in mechanics - Example

Pendulum simulation



Semi-explicit DAE:

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g}$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

- Cartesian position $\mathbf{p} \in \mathbb{R}^3$, unit mass
- Lagrange function:

$$\mathcal{L} = \frac{1}{2} m \dot{\mathbf{p}}^\top \dot{\mathbf{p}} - mg \mathbf{p}_3 - \mathbf{z} \mathbf{c}(\mathbf{p})$$

- Constraint:


$$\mathbf{c}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2$$

- Motion:

$$m \ddot{\mathbf{p}} = -\mathbf{z} \mathbf{p} - m \mathbf{g}, \quad \mathbf{g} = \begin{bmatrix} 0 \\ 0 \\ g \end{bmatrix}$$

DAEs in mechanics - Example

Pendulum simulation



Semi-explicit DAE:

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g}$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

- Cartesian position $\mathbf{p} \in \mathbb{R}^3$, unit mass
- Lagrange function:

$$\mathcal{L} = \frac{1}{2} m \dot{\mathbf{p}}^\top \dot{\mathbf{p}} - mg \mathbf{p}_3 - \mathbf{z} \mathbf{c}(\mathbf{p})$$

- Constraint:

$$\mathbf{c}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2$$


- Motion:

$$m \ddot{\mathbf{p}} = -\mathbf{z} \mathbf{p} - m \mathbf{g}, \quad \mathbf{g} = \begin{bmatrix} 0 \\ 0 \\ g \end{bmatrix}$$

- Algebraic variable \mathbf{z} “adjusts” acceleration to keep \mathbf{p} at distance L from O

DAEs in mechanics - Example

Pendulum simulation



Semi-explicit DAE:

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g}$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

- Cartesian position $\mathbf{p} \in \mathbb{R}^3$, unit mass
- Lagrange function:

$$\mathcal{L} = \frac{1}{2} m \dot{\mathbf{p}}^\top \dot{\mathbf{p}} - mg \mathbf{p}_3 - \mathbf{z} \mathbf{c}(\mathbf{p})$$

- Constraint:

$$\mathbf{c}(\mathbf{p}) = \mathbf{p}^\top \mathbf{p} - L^2$$

- Motion:

$$m \ddot{\mathbf{p}} = -\mathbf{z} \mathbf{p} - m \mathbf{g}, \quad \mathbf{g} = \begin{bmatrix} 0 \\ 0 \\ g \end{bmatrix}$$

- Algebraic variable \mathbf{z} “adjusts” acceleration to keep \mathbf{p} at distance L from O
- DAE must hold this specification as a constraint

Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$

$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-1.00}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-1.22}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-1.44}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-1.67}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-1.89}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-2.11}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-2.33}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:

$$\dot{x} = -x + z$$


$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$

$\epsilon = 10^{-2.56}$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:


$$\dot{x} = -x + z$$

$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:


$$\dot{x} = -x + z$$

$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$



Tikhonov Theorem - What is it about?

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Consider DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

approximation $\epsilon = 0$

Is the DAE a good approximation of the ODE?

Example

ODE:


$$\dot{x} = -x + z$$

$$\epsilon \dot{z} = x - 2z + u$$

DAE approximation:

$$\dot{x} = -x + z$$

$$0 = x - 2z + u$$



For $\epsilon \rightarrow 0$, DAE matches ODE almost everywhere

Tikhonov Theorem - Mechanical example

Systems often have “slow” & “fast” dynamics

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$

ODE:

$$\dot{\theta}_1 = \omega_1$$

$$J_1 \dot{\omega}_1 = T - K(\theta_1 - \theta_2) - \xi K(\omega_1 - \omega_2)$$

$$\dot{\theta}_2 = \omega_2$$

$$J_2 \dot{\omega}_2 = -K(\theta_2 - \theta_1) - \xi K(\omega_2 - \omega_1)$$

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$

ODE:

$$\dot{\theta}_1 = \omega_1$$

$$J_1 \dot{\omega}_1 = T - K(\theta_1 - \theta_2) - \xi K(\omega_1 - \omega_2)$$

$$\dot{\theta}_2 = \omega_2$$

$$J_2 \dot{\omega}_2 = -K(\theta_2 - \theta_1) - \xi K(\omega_2 - \omega_1)$$

- If shaft is highly rigid ($K \gg J_1$), modelling its dynamics is an “overkill”
- If neglected $\omega = \omega_1 = \omega_2$ and

$$\dot{\omega} = \frac{1}{J_1 + J_2} T$$

- Tikhonov provides a systematic way of neglecting fast dynamics

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\text{with } \epsilon \ll 1$$



$$\text{with } J_1 \ll K$$

ODE:

$$\dot{\theta}_1 = \omega_1$$

$$\frac{J_1}{K} \dot{\omega}_1 = \frac{T}{K} + \Delta + \xi \eta$$

$$\dot{\theta}_2 = \omega_2$$

$$J_2 \dot{\omega}_2 = -K\Delta - \xi K \eta$$

where $\Delta = \theta_2 - \theta_1$ and $\eta = \omega_2 - \omega_1$.

- If shaft is highly rigid ($K \gg J_1$), modelling its dynamics is an “overkill”
- If neglected $\omega = \omega_1 = \omega_2$ and

$$\dot{\omega} = \frac{1}{J_1 + J_2} T$$

- Tikhonov provides a systematic way of neglecting fast dynamics

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$

ODE:

$$\dot{\Delta} = \eta$$

$$\frac{J_1}{K} \dot{\eta} = -\frac{T}{K} + \left(1 + \frac{J_1}{J_2}\right) (-\Delta - \xi \eta)$$

$$\dot{\theta}_2 = \omega_2$$

$$J_2 \dot{\omega}_2 = -K\Delta - \xi K \eta$$

where $\Delta = \theta_2 - \theta_1$ and $\eta = \omega_2 - \omega_1$.

- If shaft is highly rigid ($K \gg J_1$), modelling its dynamics is an “overkill”
- If neglected $\omega = \omega_1 = \omega_2$ and

$$\dot{\omega} = \frac{1}{J_1 + J_2} T$$

- Tikhonov provides a systematic way of neglecting fast dynamics

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$

ODE:

$$\dot{\mathbf{x}} = \begin{bmatrix} \omega_2 \\ -\frac{K}{J_2}\Delta - \frac{\xi K}{J_2}\eta \end{bmatrix}$$

$$\epsilon \dot{\mathbf{z}} = \begin{bmatrix} \frac{J_1}{K}\eta \\ -\frac{T}{K} + \left(1 + \frac{J_1}{J_2}\right)(-\Delta - \xi\eta) \end{bmatrix}$$

where $\mathbf{z} = \begin{bmatrix} \Delta \\ \eta \end{bmatrix}$, $\mathbf{x} = \begin{bmatrix} \theta_2 \\ \omega_2 \end{bmatrix}$ and $\epsilon \equiv \frac{J_1}{K}$.

- If shaft is highly rigid ($K \gg J_1$), modelling its dynamics is an “overkill”
- If neglected $\omega = \omega_1 = \omega_2$ and

$$\dot{\omega} = \frac{1}{J_1 + J_2} T$$

- Tikhonov provides a systematic way of neglecting fast dynamics

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$

ODE:

$$\dot{\mathbf{x}} = \begin{bmatrix} -\frac{K}{J_2} \Delta - \frac{\xi K}{J_2} \eta \\ \frac{J_1}{K} \eta \end{bmatrix}$$

$$\epsilon \dot{\mathbf{z}} = \begin{bmatrix} -\frac{T}{K} + \left(1 + \frac{J_1}{J_2}\right) (-\Delta - \xi \eta) \end{bmatrix}$$

where $\mathbf{z} = \begin{bmatrix} \Delta \\ \eta \end{bmatrix}$, $\mathbf{x} = \begin{bmatrix} \theta_2 \\ \omega_2 \end{bmatrix}$ and $\epsilon \equiv \frac{J_1}{K}$.

DAE approximation

$$\dot{\mathbf{x}} = \begin{bmatrix} -\frac{\omega_2}{J_2} \Delta \end{bmatrix} = \begin{bmatrix} \frac{\omega_2}{J_1 + J_2} T \end{bmatrix}$$

$$\eta = 0$$

$$\Delta = -\frac{J_2}{K(J_1 + J_2)} T$$

... approximates “fast” dynamics by pretending they decay to their steady-state

Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:




$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$



Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:




$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$



Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:




$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$



Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:




$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$



Tikhonov Theorem - Mechanical example





Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



Tikhonov Theorem - Mechanical example





Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



Tikhonov Theorem - Mechanical example


Systems often have “slow” & “fast” dynamics, e.g.

Consider ODE:




$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\epsilon \dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

with $\epsilon \ll 1$



with $J_1 \ll K$



Tikhonov Theorem - What does it say?

Tikhonov Theorem

Consider the ordinary differential equation (ODE):

$$\begin{aligned}\dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{z}) \\ \epsilon \dot{\mathbf{z}} &= \mathbf{g}(\mathbf{x}, \mathbf{z})\end{aligned}\quad \text{of solution } \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) \quad (\epsilon > 0)$$

Tikhonov Theorem - What does it say?

Tikhonov Theorem

Consider the ordinary differential equation (ODE):

$$\begin{aligned}\dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{z}) \\ \epsilon \dot{\mathbf{z}} &= \mathbf{g}(\mathbf{x}, \mathbf{z})\end{aligned}\quad \text{of solution } \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) \quad (\epsilon > 0)$$

Suppose:

- the dynamics $\dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z})$ are **stable** $\forall \mathbf{x}$
- matrix $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is **full rank** everywhere

Tikhonov Theorem - What does it say?

Tikhonov Theorem

Consider the ordinary differential equation (ODE):

$$\begin{aligned}\dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{z}) \\ \epsilon \dot{\mathbf{z}} &= \mathbf{g}(\mathbf{x}, \mathbf{z})\end{aligned}\quad \text{of solution } \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) \quad (\epsilon > 0)$$

Suppose:

- the dynamics $\dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z})$ are **stable** $\forall \mathbf{x}$
- matrix $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is **full rank** everywhere

then

$$\lim_{\epsilon \rightarrow 0} \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) = \mathbf{x}_0(t), \mathbf{z}_0(t) \quad \text{almost everywhere}$$

Tikhonov Theorem - What does it say?

Tikhonov Theorem

Consider the ordinary differential equation (ODE):

$$\begin{aligned}\dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{z}) \\ \epsilon \dot{\mathbf{z}} &= \mathbf{g}(\mathbf{x}, \mathbf{z})\end{aligned}\quad \text{of solution } \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) \quad (\epsilon > 0)$$

Suppose:

- the dynamics $\dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z})$ are **stable** $\forall \mathbf{x}$
- matrix $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is **full rank** everywhere

then

$$\lim_{\epsilon \rightarrow 0} \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) = \mathbf{x}_0(t), \mathbf{z}_0(t) \quad \text{almost everywhere}$$

where $\mathbf{x}_0(t), \mathbf{z}_0(t)$ is solution of DAE

$$\begin{aligned}\dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{z}) \\ 0 &= \mathbf{g}(\mathbf{x}, \mathbf{z})\end{aligned}$$

Tikhonov Theorem - What does it say?

Tikhonov Theorem

Consider the ordinary differential equation (ODE):

$$\begin{aligned}\dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{z}) \\ \epsilon \dot{\mathbf{z}} &= \mathbf{g}(\mathbf{x}, \mathbf{z})\end{aligned}\quad \text{of solution } \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) \quad (\epsilon > 0)$$

Suppose:

- the dynamics $\dot{\mathbf{z}} = \mathbf{g}(\mathbf{x}, \mathbf{z})$ are **stable** $\forall \mathbf{x}$
- matrix $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is **full rank** everywhere

then

$$\lim_{\epsilon \rightarrow 0} \mathbf{x}_\epsilon(t), \mathbf{z}_\epsilon(t) = \mathbf{x}_0(t), \mathbf{z}_0(t) \quad \text{almost everywhere}$$

where $\mathbf{x}_0(t), \mathbf{z}_0(t)$ is solution of DAE

$$\begin{aligned}\dot{\mathbf{x}} &= \mathbf{f}(\mathbf{x}, \mathbf{z}) \\ 0 &= \mathbf{g}(\mathbf{x}, \mathbf{z})\end{aligned}$$

In other words:

- Fast dynamics should “decay”
- $\mathbf{g}(\mathbf{x}, \mathbf{z}) = 0$ must be “solvable for \mathbf{z} ” (see Implicit Function Theorem)

Simulating a DAE - “Easy” DAEs

Simulating an ODE $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$, requires solving for $\dot{\mathbf{x}}$ for all \mathbf{x}, \mathbf{u} on the trajectory

What does it mean to be able to simulate a DAE “easily”?

Simulating a DAE - “Easy” DAEs

Simulating an ODE $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$, requires solving for $\dot{\mathbf{x}}$ for all \mathbf{x}, \mathbf{u} on the trajectory

What does it mean to be able to simulate a DAE “easily”?

- for a semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5a)$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5b)$$

(5a) delivers $\dot{\mathbf{x}}$ if \mathbf{z} is known, \mathbf{z} must be provided by (5b)

Simulating a DAE - “Easy” DAEs

Simulating an ODE $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$, requires solving for $\dot{\mathbf{x}}$ for all \mathbf{x}, \mathbf{u} on the trajectory

What does it mean to be able to simulate a DAE “easily”?

- for a semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5a)$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5b)$$

(5a) delivers $\dot{\mathbf{x}}$ if \mathbf{z} is known, \mathbf{z} must be provided by (5b)

- for a fully-implicit DAE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

both $\dot{\mathbf{x}}, \mathbf{z}$ must be provided by $\mathbf{F} = 0$

Simulating a DAE - “Easy” DAEs

Simulating an ODE $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$, requires solving for $\dot{\mathbf{x}}$ for all \mathbf{x}, \mathbf{u} on the trajectory

What does it mean to be able to simulate a DAE “easily”?

- for a semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5a)$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5b)$$

(5a) delivers $\dot{\mathbf{x}}$ if \mathbf{z} is known, \mathbf{z} must be provided by (5b)

- for a fully-implicit DAE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

both $\dot{\mathbf{x}}, \mathbf{z}$ must be provided by $\mathbf{F} = 0$

As long as the DAE delivers $\dot{\mathbf{x}}, \mathbf{z}$, simulating is “just” about following $\dot{\mathbf{x}}$.

Simulating a DAE - “Easy” DAEs

Simulating an ODE $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{x}, \mathbf{u}) = 0$, requires solving for $\dot{\mathbf{x}}$ for all \mathbf{x}, \mathbf{u} on the trajectory

What does it mean to be able to simulate a DAE “easily”?

- for a semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5a)$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) \quad (5b)$$

(5a) delivers $\dot{\mathbf{x}}$ if \mathbf{z} is known, \mathbf{z} must be provided by (5b)

- for a fully-implicit DAE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

both $\dot{\mathbf{x}}, \mathbf{z}$ must be provided by $\mathbf{F} = 0$

As long as the DAE delivers $\dot{\mathbf{x}}, \mathbf{z}$, simulating is “just” about following $\dot{\mathbf{x}}$.

When can the algebraic equation

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

be solved for \mathbf{z} ?

When can the DAE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

be solved for both $\dot{\mathbf{x}}, \mathbf{z}$?

“Easy” DAEs - Semi-explicit case

When can a semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

be solved for $\dot{\mathbf{x}}, \mathbf{z}$?

“Easy” DAEs - Semi-explicit case

When can a semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

be solved for $\dot{\mathbf{x}}, \mathbf{z}$?

Reminder - IFT

If $\mathbf{F}(\mathbf{y}, \mathbf{x})$ is smooth, and $\frac{\partial \mathbf{F}(\mathbf{y}, \mathbf{x})}{\partial \mathbf{y}}$ always full rank then $\mathbf{y} = \mathbf{f}(\mathbf{x})$ exists such that

$$\mathbf{F}(\mathbf{f}(\mathbf{x}), \mathbf{x}) = 0 \quad \text{for all } \mathbf{x}$$

“Easy” DAEs - Semi-explicit case

When can a semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

be solved for $\dot{\mathbf{x}}, \mathbf{z}$?

Reminder - IFT

If $\mathbf{F}(\mathbf{y}, \mathbf{x})$ is smooth, and $\frac{\partial \mathbf{F}(\mathbf{y}, \mathbf{x})}{\partial \mathbf{y}}$ always full rank then $\mathbf{y} = \mathbf{f}(\mathbf{x})$ exists such that


$$\mathbf{F}(\mathbf{f}(\mathbf{x}), \mathbf{x}) = 0 \quad \text{for all } \mathbf{x}$$

Semi-explicit DAE case

- Getting $\dot{\mathbf{x}}$ from the first equation is trivial
- Implicit Function Theorem says that we can solve $\mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u}) = 0$ for \mathbf{z}

if (square) Jacobian $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is full rank

Back to our mechanical example



Semi-explicit DAE:


$$\left. \begin{aligned} \dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \end{aligned} \right\} \equiv \mathbf{f} \quad (6a)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad \left. \right\} \equiv \mathbf{g} \quad (6b)$$

State

$$\mathbf{x} = \begin{bmatrix} \mathbf{p} \\ \mathbf{v} \end{bmatrix} \quad \text{and} \quad \mathbf{z}$$

Back to our mechanical example



Semi-explicit DAE:

$$\left. \begin{aligned} \dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \end{aligned} \right\} \equiv \mathbf{f} \quad (6a)$$


$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \} \equiv \mathbf{g} \quad (6b)$$

State

$$\mathbf{x} = \begin{bmatrix} \mathbf{p} \\ \mathbf{v} \end{bmatrix} \quad \text{and} \quad \mathbf{z}$$

Is that an “easy” DAE?

Back to our mechanical example



Semi-explicit DAE:

$$\left. \begin{aligned} \dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \end{aligned} \right\} \equiv \mathbf{f} \quad (6a)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \} \equiv \mathbf{g} \quad (6b)$$


State

$$\mathbf{x} = \begin{bmatrix} \mathbf{p} \\ \mathbf{v} \end{bmatrix} \quad \text{and} \quad \mathbf{z}$$

Is that an “easy” DAE?

- Algebraic part is $\mathbf{g} = \mathbf{p}^\top \mathbf{p} - L^2$

Back to our mechanical example



Semi-explicit DAE:

$$\left. \begin{aligned} \dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \end{aligned} \right\} \equiv \mathbf{f} \quad (6a)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad \left. \right\} \equiv \mathbf{g} \quad (6b)$$


State

$$\mathbf{x} = \begin{bmatrix} \mathbf{p} \\ \mathbf{v} \end{bmatrix} \quad \text{and} \quad \mathbf{z}$$

Is that an “easy” DAE?

- Algebraic part is $\mathbf{g} = \mathbf{p}^\top \mathbf{p} - L^2$
- Jacobian $\frac{\partial \mathbf{g}}{\partial \mathbf{z}} = 0$. In fact, \mathbf{z} does not enter in \mathbf{g} !

Back to our mechanical example



Semi-explicit DAE:

$$\left. \begin{aligned} \dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \end{aligned} \right\} \equiv \mathbf{f} \quad (6a)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad \left. \right\} \equiv \mathbf{g} \quad (6b)$$


State

$$\mathbf{x} = \begin{bmatrix} \mathbf{p} \\ \mathbf{v} \end{bmatrix} \quad \text{and} \quad \mathbf{z}$$

Is that an “easy” DAE?

- Algebraic part is $\mathbf{g} = \mathbf{p}^\top \mathbf{p} - L^2$
- Jacobian $\frac{\partial \mathbf{g}}{\partial \mathbf{z}} = 0$. In fact, \mathbf{z} does not enter in \mathbf{g} !
- (6b) cannot be solved for \mathbf{z} ...

Back to our mechanical example



Semi-explicit DAE:

$$\left. \begin{aligned} \dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \end{aligned} \right\} \equiv \mathbf{f} \quad (6a)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad \left. \right\} \equiv \mathbf{g} \quad (6b)$$


State

$$\mathbf{x} = \begin{bmatrix} \mathbf{p} \\ \mathbf{v} \end{bmatrix} \quad \text{and} \quad \mathbf{z}$$

Is that an “easy” DAE?

- Algebraic part is $\mathbf{g} = \mathbf{p}^\top \mathbf{p} - L^2$
- Jacobian $\frac{\partial \mathbf{g}}{\partial \mathbf{z}} = 0$. In fact, \mathbf{z} does not enter in \mathbf{g} !
- (6b) cannot be solved for \mathbf{z} ...
- (6a) delivers $\dot{\mathbf{x}}$ if we know \mathbf{z} ...

Back to our mechanical example



Semi-explicit DAE:

$$\left. \begin{aligned} \dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \end{aligned} \right\} \equiv \mathbf{f} \quad (6a)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad \left. \right\} \equiv \mathbf{g} \quad (6b)$$

State

$$\mathbf{x} = \begin{bmatrix} \mathbf{p} \\ \mathbf{v} \end{bmatrix} \quad \text{and} \quad \mathbf{z}$$

Is that an “easy” DAE?

- Algebraic part is $\mathbf{g} = \mathbf{p}^\top \mathbf{p} - L^2$
- Jacobian $\frac{\partial \mathbf{g}}{\partial \mathbf{z}} = 0$. In fact, \mathbf{z} does not enter in \mathbf{g} !
- (6b) cannot be solved for \mathbf{z} ...
- (6a) delivers $\dot{\mathbf{x}}$ if we know \mathbf{z} ...

It is not an “easy” DAE

“Easy” DAEs - Fully-implicit case

When can a fully-implicit DAE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

be solved for both $\dot{\mathbf{x}}, \mathbf{z}$?

“Easy” DAEs - Fully-implicit case

When can a fully-implicit DAE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

be solved for both $\dot{\mathbf{x}}, \mathbf{z}$?

Reminder - IFT

If $\mathbf{F}(\mathbf{y}, \mathbf{x})$ is smooth, and $\frac{\partial \mathbf{F}(\mathbf{y}, \mathbf{x})}{\partial \mathbf{y}}$ always full rank then $\mathbf{y} = \mathbf{f}(\mathbf{x})$ exists such that

$$\mathbf{F}(\mathbf{f}(\mathbf{x}), \mathbf{x}) = 0 \quad \text{for all } \mathbf{x}$$

“Easy” DAEs - Fully-implicit case

When can a fully-implicit DAE

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

be solved for both $\dot{\mathbf{x}}, \mathbf{z}$?

Reminder - IFT

If $\mathbf{F}(\mathbf{y}, \mathbf{x})$ is smooth, and $\frac{\partial \mathbf{F}(\mathbf{y}, \mathbf{x})}{\partial \mathbf{y}}$ always full rank then $\mathbf{y} = \mathbf{f}(\mathbf{x})$ exists such that

$$\mathbf{F}(\mathbf{f}(\mathbf{x}), \mathbf{x}) = 0 \quad \text{for all } \mathbf{x}$$

Fully-Implicit DAE case

Implicit **F**unction **T**heorem says that we can solve $\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$ for $\dot{\mathbf{x}}, \mathbf{z}$

if (square) Jacobian $\begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix}$ is full rank

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver \dot{x} , z ...

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}$, \mathbf{z} ...
... but they (may) deliver well-defined trajectories nonetheless!!

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}, \mathbf{z}, \dots$


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}, \mathbf{z}, \dots$


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}, \mathbf{z}, \dots$


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}$, \mathbf{z} ...


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds
- Chain of “influence”:

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}}$$

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}$, \mathbf{z} ...


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds
- Chain of “influence”:

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} = \dot{\mathbf{p}}$$

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}$, \mathbf{z} ...


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds
- Chain of “influence”:

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} = \dot{\mathbf{p}} \xrightarrow{\int dt} \mathbf{p}$$

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}, \mathbf{z}, \dots$


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds
- Chain of “influence”:

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} = \dot{\mathbf{p}} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} (7c)$$

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}, \mathbf{z}, \dots$


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds
- Chain of “influence”:

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} = \dot{\mathbf{p}} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} (7c)$$

connects \mathbf{z} to (7c) via **2 integrations**

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}, \mathbf{z}, \dots$


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds
- Chain of “influence”:

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} = \dot{\mathbf{p}} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} (7c)$$

connects \mathbf{z} to (7c) via **2 integrations**

- How to reconnect \mathbf{z} to (7c) *algebraically*? (equation we could “solve for \mathbf{z} ” ...)

“Hard” DAEs

Definition (informal): “hard” DAEs do not *readily* deliver $\dot{\mathbf{x}}, \mathbf{z}, \dots$


... but they (may) deliver well-defined trajectories nonetheless!!

Example: how do we get a trajectory from

$$\dot{\mathbf{p}} = \mathbf{v} \quad (7a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \quad (7b)$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2 \quad (7c)$$



Reminder:

- \mathbf{z} “adjusts” $\dot{\mathbf{v}}$ so that (7c) holds
- Chain of “influence”:

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} = \dot{\mathbf{p}} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} (7c)$$

connects \mathbf{z} to (7c) via **2 integrations**

- How to reconnect \mathbf{z} to (7c) *algebraically*? (equation we could “solve for \mathbf{z} ” ...)



Apply $\frac{d}{dt}$ twice on (7c) to “rewind” the chain to \mathbf{z}

Solving “Hard” DAEs

Apply $\frac{d}{dt}$ on $0 = \mathbf{p}^\top \mathbf{p} - L^2$, to “rewind” the chain to \mathbf{z}

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} 0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g}$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\xrightarrow{\frac{d}{dt}}$$

Solving “Hard” DAEs

Apply $\frac{d}{dt}$ on $0 = \mathbf{p}^\top \mathbf{p} - L^2$, to “rewind” the chain to \mathbf{z}

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} 0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{v}\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

Solving “Hard” DAEs

Apply $\frac{d}{dt}$ on $0 = \mathbf{p}^\top \mathbf{p} - L^2$, to “rewind” the chain to \mathbf{z}

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} 0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{v}\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Solving “Hard” DAEs

Apply $\frac{d}{dt}$ on $0 = \mathbf{p}^\top \mathbf{p} - L^2$, to “rewind” the chain to \mathbf{z}

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} 0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}}\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Yields “easy” DAE:

$$\dot{\mathbf{p}} = \mathbf{v} \quad (8a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \quad (8b)$$

$$0 = -\frac{1}{m}\mathbf{p}^\top \mathbf{p}\mathbf{z} - \mathbf{p}^\top \mathbf{g} + \mathbf{v}^\top \mathbf{v} \quad (8c)$$

as (8c) delivers \mathbf{z} (if $\mathbf{p} \neq 0$)

Solving “Hard” DAEs

Apply $\frac{d}{dt}$ on $0 = \mathbf{p}^\top \mathbf{p} - L^2$, to “rewind” the chain to \mathbf{z}

$$\mathbf{z} \xrightarrow{\text{alg.}} \dot{\mathbf{v}} \xrightarrow{\int dt} \mathbf{v} \xrightarrow{\int dt} \mathbf{p} \xrightarrow{\text{alg.}} 0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

 $\xrightarrow{\frac{d}{dt}}$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{v}\end{aligned}$$

 $\xrightarrow{\frac{d}{dt}}$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Yields “easy” DAE:

$$\dot{\mathbf{p}} = \mathbf{v} \quad (8a)$$

$$\dot{\mathbf{v}} = -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \quad (8b)$$

$$0 = -\frac{1}{m}\mathbf{p}^\top \mathbf{p}\mathbf{z} - \mathbf{p}^\top \mathbf{g} + \mathbf{v}^\top \mathbf{v} \quad (8c)$$

as (8c) delivers \mathbf{z} (if $\mathbf{p} \neq 0$)

 $\xrightarrow{\frac{d}{dt}}$

One more $\frac{d}{dt}$ on (8c) turns (8) into an ODE

Differential Index


Definition

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Differential Index

Definition

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE



Pendulum example

Index 3

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

Physical model
"Hard" DAE

$\xrightarrow{\frac{d^2}{dt^2}}$

Index 1

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= 2\mathbf{p}^\top \dot{\mathbf{v}} + 2\mathbf{v}^\top \mathbf{v}\end{aligned}$$

2 time-differentiations
→ "Easy" DAE


$\xrightarrow{\frac{d}{dt}}$

ODE

Differential Index

Definition

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE



Pendulum example

Index 3

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

Physical model
“Hard” DAE

$\xrightarrow{\frac{d^2}{dt^2}}$

Index 1

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= 2\mathbf{p}^\top \dot{\mathbf{v}} + 2\mathbf{v}^\top \mathbf{v}\end{aligned}$$

2 time-differentiations
→ “Easy” DAE

$\xrightarrow{\frac{d}{dt}}$

ODE

The transformation **index-n** $\xrightarrow{\frac{d^{n-1}}{dt^{n-1}}}$ **index-1** is called **index reduction**

Index-1 DAEs

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 DAEs

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

Index-1 DAEs

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{d}{dt}\mathbf{g} = 0$ yields:

$$\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{d}{dt}\mathbf{g} = 0$ yields:

$$\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{g}}{\partial \mathbf{z}}^{-1} \left(\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \mathbf{f} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{d}{dt}\mathbf{g} = 0$ yields:

$$\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{g}}{\partial \mathbf{z}}^{-1} \left(\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ must be full rank.

Index-1 DAEs

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{d}{dt}\mathbf{g} = 0$ yields:

$$\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{g}}{\partial \mathbf{z}}^{-1} \left(\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ must be full rank.

Index-1 fully-implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

Index-1 DAEs

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{d}{dt}\mathbf{g} = 0$ yields:

$$\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{g}^{-1}}{\partial \mathbf{z}} \left(\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ must be full rank.

Index-1 fully-implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

$\frac{d}{dt}\mathbf{F} = 0$ yields:

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} \ddot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{F}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

Index-1 DAEs

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{d}{dt}\mathbf{g} = 0$ yields:

$$\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{g}}{\partial \mathbf{z}}^{-1} \left(\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ must be full rank.

Index-1 fully-implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

$\frac{d}{dt}\mathbf{F} = 0$ yields:

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} \ddot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{F}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{x}} \\ \dot{\mathbf{z}} \end{bmatrix} + \frac{\partial \mathbf{F}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

Index-1 DAEs

The differential index of a DAE is the number of $\frac{d}{dt}$ needed to turn it into an ODE

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$\frac{d}{dt}\mathbf{g} = 0$ yields:

$$\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$\dot{\mathbf{z}} = -\frac{\partial \mathbf{g}}{\partial \mathbf{z}}^{-1} \left(\frac{\partial \mathbf{g}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{g}}{\partial \mathbf{u}} \dot{\mathbf{u}} \right)$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ must be full rank.

Index-1 fully-implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

$\frac{d}{dt}\mathbf{F} = 0$ yields:

$$\frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} \ddot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \dot{\mathbf{z}} + \frac{\partial \mathbf{F}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

and should give us the ODE:

$$\begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix} \begin{bmatrix} \ddot{\mathbf{x}} \\ \dot{\mathbf{z}} \end{bmatrix} + \frac{\partial \mathbf{F}}{\partial \mathbf{x}} \dot{\mathbf{x}} + \frac{\partial \mathbf{F}}{\partial \mathbf{u}} \dot{\mathbf{u}} = 0$$

where $\begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix}$ must be full rank.

Delivers $\ddot{\mathbf{x}}$ and $\dot{\mathbf{z}}$.

Index-1 DAEs are “easy” DAEs - Connection to IFT

Index-1 DAEs readily deliver \dot{x}, z and are therefore “easy”

Index-1 DAEs are “easy” DAEs - Connection to IFT

Index-1 DAEs readily deliver $\dot{\mathbf{x}}$, \mathbf{z} and are therefore “easy”

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is full rank.

Then

- IFT: $\mathbf{g} = 0$ can be solved for \mathbf{z}
- $\dot{\mathbf{x}}$ is delivered by 1st equations
- DAE can be “easily” simulated

Index-1 DAEs are “easy” DAEs - Connection to IFT

Index-1 DAEs readily deliver $\dot{\mathbf{x}}$, \mathbf{z} and are therefore “easy”

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is full rank.

Then

- IFT: $\mathbf{g} = 0$ can be solved for \mathbf{z}
- $\dot{\mathbf{x}}$ is delivered by 1st equations
- DAE can be “easily” simulated

Index-1 DAEs are “easy” DAEs - Connection to IFT

Index-1 DAEs readily deliver $\dot{\mathbf{x}}$, \mathbf{z} and are therefore “easy”

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is full rank.

Then

- IFT: $\mathbf{g} = 0$ can be solved for \mathbf{z}
- $\dot{\mathbf{x}}$ is delivered by 1st equations
- DAE can be “easily” simulated

Index-1 fully-implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

Index-1 DAEs are “easy” DAEs - Connection to IFT

Index-1 DAEs readily deliver $\dot{\mathbf{x}}, \mathbf{z}$ and are therefore “easy”

Index-1 semi-explicit DAE:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

where $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ is full rank.

Then

- IFT: $\mathbf{g} = 0$ can be solved for \mathbf{z}
- $\dot{\mathbf{x}}$ is delivered by 1st equations
- DAE can be “easily” simulated

Index-1 fully-implicit DAE:

$$\mathbf{F}(\dot{\mathbf{x}}, \mathbf{z}, \mathbf{x}, \mathbf{u}) = 0$$

where $\begin{bmatrix} \frac{\partial \mathbf{F}}{\partial \dot{\mathbf{x}}} & \frac{\partial \mathbf{F}}{\partial \mathbf{z}} \end{bmatrix}$ is full rank.

Then

- IFT: $\mathbf{F} = 0$ can be solved for $\dot{\mathbf{x}}, \mathbf{z}$
- DAE can be “easily” simulated

Index reduction for semi-explicit DAEs - A general view

High-index semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

Algorithm

Index reduction for semi-explicit DAEs - A general view

High-index semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

Algorithm

- 1 Check if the DAE system is index 1 (i.e. $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ full rank).
If yes, stop.
- 2 Identify a subset of algebraic equations that **can be solved** for a subset of algebraic variables.
- 3 Apply $\frac{d}{dt}$ on the remaining algebraic equations that contain the differential variables \mathbf{x}_j .
- 4 Terms $\dot{\mathbf{x}}_j$ will appear in these differentiated equations.
- 5 Substitute the $\dot{\mathbf{x}}_j$ with $\mathbf{f}_j(\mathbf{x}, \mathbf{z}, \mathbf{u})$. This leads to new algebraic equations.
- 6 With this new DAE system, go to step 1.

Index reduction for semi-explicit DAEs - A general view

High-index semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

Algorithm

- 1 Check if the DAE system is index 1 (i.e. $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ full rank).
If yes, stop.
- 2 Identify a subset of algebraic equations that **can be solved** for a subset of algebraic variables.
- 3 Apply $\frac{d}{dt}$ on the remaining algebraic equations that contain the differential variables \mathbf{x}_j .
- 4 Terms $\dot{\mathbf{x}}_j$ will appear in these differentiated equations.
- 5 Substitute the $\dot{\mathbf{x}}_j$ with $\mathbf{f}_j(\mathbf{x}, \mathbf{z}, \mathbf{u})$. This leads to new algebraic equations.
- 6 With this new DAE system, go to step 1.

Index reduction for semi-explicit DAEs - A general view

High-index semi-explicit DAE

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

$$0 = \mathbf{g}(\mathbf{x}, \mathbf{z}, \mathbf{u})$$

Algorithm

- 1 Check if the DAE system is index 1 (i.e. $\frac{\partial \mathbf{g}}{\partial \mathbf{z}}$ full rank).
If yes, stop.
- 2 Identify a subset of algebraic equations that **can be solved** for a subset of algebraic variables.
- 3 Apply $\frac{d}{dt}$ on the remaining algebraic equations that contain the differential variables \mathbf{x}_j .
- 4 Terms $\dot{\mathbf{x}}_j$ will appear in these differentiated equations.
- 5 Substitute the $\dot{\mathbf{x}}_j$ with $\mathbf{f}_j(\mathbf{x}, \mathbf{z}, \mathbf{u})$. This leads to new algebraic equations.
- 6 With this new DAE system, go to step 1.

Writing a general-purpose "Index-reduction algorithm" can be very tricky, as one of the steps is not easily automated

Differential index & Index reduction - Examples

$$z = u - x_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_2 - u = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

Differential index & Index reduction - Examples

$$\begin{array}{lcl} \mathbf{z} = u - \mathbf{x}_2 & & \dot{\mathbf{z}} = \dot{u} - \dot{\mathbf{x}}_2 \\ \dot{\mathbf{x}}_2 = \mathbf{x}_1 & \xrightarrow{\frac{d}{dt}} & \dot{\mathbf{x}}_2 = \mathbf{x}_1 \\ \dot{\mathbf{x}}_1 = u - \mathbf{x}_2 & & \dot{\mathbf{x}}_1 = u - \mathbf{x}_2 \end{array} \quad \xrightarrow{\text{Alg.}}$$

$$\begin{array}{lcl} \dot{\mathbf{x}}_1 - z = 0 & & \\ \dot{\mathbf{x}}_2 - \mathbf{x}_1 = 0 & \xrightarrow{\frac{d}{dt}} & \\ \mathbf{x}_2 - u = 0 & & \end{array}$$

Differential index & Index reduction - Examples

$$\begin{array}{lcl} \mathbf{z} = \mathbf{u} - \mathbf{x}_2 & & \dot{\mathbf{z}} = \dot{\mathbf{u}} - \dot{\mathbf{x}}_2 \\ \dot{\mathbf{x}}_2 = \mathbf{x}_1 & \xrightarrow{\frac{d}{dt}} & \dot{\mathbf{x}}_2 = \mathbf{x}_1 \\ \dot{\mathbf{x}}_1 = \mathbf{u} - \mathbf{x}_2 & & \dot{\mathbf{x}}_1 = \mathbf{u} - \mathbf{x}_2 \end{array} \quad \xrightarrow{\text{Alg.}} \quad \begin{array}{l} \dot{\mathbf{z}} = \dot{\mathbf{u}} - \mathbf{x}_1 \\ \dot{\mathbf{x}}_2 = \mathbf{x}_1 \\ \dot{\mathbf{x}}_1 = \mathbf{u} - \mathbf{x}_2 \end{array}$$

$$\begin{array}{l} \dot{\mathbf{x}}_1 - \mathbf{z} = 0 \\ \dot{\mathbf{x}}_2 - \mathbf{x}_1 = 0 \\ \mathbf{x}_2 - \mathbf{u} = 0 \end{array} \quad \xrightarrow{\frac{d}{dt}}$$

Differential index & Index reduction - Examples

DAE index-1

$$z = u - x_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\dot{z} = \dot{u} - \dot{x}_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\text{Alg.}}$$

ODE

$$\dot{z} = \dot{u} - x_1$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_2 - u = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

Differential index & Index reduction - Examples

DAE index-1

$$z = u - x_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\dot{z} = \dot{u} - \dot{x}_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\text{Alg.}}$$

ODE

$$\dot{z} = \dot{u} - x_1$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_2 - u = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_2 - \dot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

Differential index & Index reduction - Examples

DAE index-1

$$z = u - x_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\dot{z} = \dot{u} - \dot{x}_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\text{Alg.}}$$

ODE

$$\dot{z} = \dot{u} - x_1$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_2 - u = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_2 - \dot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_1 - \dot{u} = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

Differential index & Index reduction - Examples

DAE index-1

$$z = u - x_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\dot{z} = \dot{u} - \dot{x}_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\text{Alg.}}$$

ODE

$$\dot{z} = \dot{u} - x_1$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_2 - u = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_2 - \dot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_1 - \dot{u} = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_1 - \ddot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

Differential index & Index reduction - Examples

DAE index-1

$$z = u - x_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\dot{z} = \dot{u} - \dot{x}_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\text{Alg.}}$$

ODE

$$\dot{z} = \dot{u} - x_1$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_2 - u = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_2 - \dot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_1 - \dot{u} = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_1 - \ddot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$z - \ddot{u} = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

Differential index & Index reduction - Examples

DAE index-1

$$z = u - x_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\dot{z} = \dot{u} - \dot{x}_2$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\xrightarrow{\text{Alg.}}$$

ODE

$$\dot{z} = \dot{u} - x_1$$

$$\dot{x}_2 = x_1$$

$$\dot{x}_1 = u - x_2$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_2 - u = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_2 - \dot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$x_1 - \dot{u} = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$\dot{x}_1 - \ddot{u} = 0$$

$$\xrightarrow{\text{Alg.}}$$

$$\dot{x}_1 - z = 0$$

$$\dot{x}_2 - x_1 = 0$$

$$z - \ddot{u} = 0$$

$$\xrightarrow{\frac{d}{dt}}$$

$$\dot{x}_1 = z$$

$$\dot{x}_2 = x_1$$

$$\dot{z} = \ddot{u}$$

Differential index & Index reduction - Examples

DAE index-1

$$\begin{aligned}z &= u - x_2 \\ \dot{x}_2 &= x_1 \\ \dot{x}_1 &= u - x_2\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\begin{aligned}\dot{z} &= \dot{u} - \dot{x}_2 \\ \dot{x}_2 &= x_1 \\ \dot{x}_1 &= u - x_2\end{aligned}$$

$$\xrightarrow{\text{Alg.}}$$

ODE

$$\begin{aligned}\dot{z} &= \dot{u} - x_1 \\ \dot{x}_2 &= x_1 \\ \dot{x}_1 &= u - x_2\end{aligned}$$

DAE index 3

$$\begin{aligned}\dot{x}_1 - z &= 0 \\ \dot{x}_2 - x_1 &= 0 \\ x_2 - u &= 0\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

DAE index 2

$$\begin{aligned}\dot{x}_1 - z &= 0 \\ \dot{x}_2 - x_1 &= 0 \\ \dot{x}_2 - \dot{u} &= 0\end{aligned}$$

$$\xrightarrow{\text{Alg.}}$$

DAE index 2

$$\begin{aligned}\dot{x}_1 - z &= 0 \\ \dot{x}_2 - x_1 &= 0 \\ x_1 - \dot{u} &= 0\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

DAE index 1

$$\begin{aligned}\dot{x}_1 - z &= 0 \\ \dot{x}_2 - x_1 &= 0 \\ \dot{x}_1 - \ddot{u} &= 0\end{aligned}$$

$$\xrightarrow{\text{Alg.}}$$

DAE index 1

$$\begin{aligned}\dot{x}_1 - z &= 0 \\ \dot{x}_2 - x_1 &= 0 \\ z - \ddot{u} &= 0\end{aligned}$$

$$\xrightarrow{\frac{d}{dt}}$$

ODE

$$\begin{aligned}\dot{x}_1 &= z \\ \dot{x}_2 &= x_1 \\ \dot{z} &= \ddot{u}\end{aligned}$$

Consistency conditions - Pendulum example

Pendulum equations:

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g}$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\xrightarrow{\frac{d^2}{dt^2}}$$

Consistency conditions - Pendulum example

Pendulum equations:

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g}$$

$$0 = \mathbf{p}^\top \mathbf{p} - L^2$$

$$\xrightarrow{\frac{d^2}{dt^2}} \rightarrow$$

$$\dot{\mathbf{p}} = \mathbf{v}$$

$$\dot{\mathbf{v}} = -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g}$$

$$0 = \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}$$

Consistency conditions - Pendulum example

Pendulum equations:

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

Original model imposes

$$\mathbf{p}^\top \mathbf{p} = L^2 \quad (9)$$

$$\xrightarrow{\frac{d^2}{dt^2}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Index-1 model imposes

$$\frac{d^2}{dt^2} (\mathbf{p}^\top \mathbf{p}) = 0 \quad (10)$$

Does (10) \Rightarrow (9)? I.e. does index-1 model match original model?

Consistency conditions - Pendulum example

Pendulum equations:

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

Original model imposes

$$\mathbf{p}^\top \mathbf{p} = L^2 \quad (9)$$

$$\xrightarrow{\frac{d^2}{dt^2}} \rightarrow$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Index-1 model imposes

$$\frac{d^2}{dt^2} (\mathbf{p}^\top \mathbf{p}) = 0 \quad (10)$$

Does (10) \Rightarrow (9)? I.e. does index-1 model match original model?

Let's write $c(\mathbf{p}(t)) = \mathbf{p}(t)^\top \mathbf{p}(t) - L^2$

Consistency conditions - Pendulum example

Pendulum equations:

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

Original model imposes

$$\mathbf{p}^\top \mathbf{p} = L^2 \quad (9)$$

$$\xrightarrow{\frac{d^2}{dt^2}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m}\mathbf{z}\mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Index-1 model imposes

$$\frac{d^2}{dt^2} (\mathbf{p}^\top \mathbf{p}) = 0 \quad (10)$$

Does (10) \Rightarrow (9)? I.e. does index-1 model match original model?

Let's write $c(\mathbf{p}(t)) = \mathbf{p}(t)^\top \mathbf{p}(t) - L^2$ then index-1 model yields:

$$\ddot{c} = 0 \quad \Rightarrow \quad c(\mathbf{p}(t)) = c(\mathbf{p}(0)) + \dot{c}(\mathbf{p}(0), \dot{\mathbf{p}}(0))t$$

Consistency conditions - Pendulum example

Pendulum equations:

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

Original model imposes

$$\mathbf{p}^\top \mathbf{p} = L^2 \quad (9)$$

$$\xrightarrow{\frac{d^2}{dt^2}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Index-1 model imposes

$$\frac{d^2}{dt^2} (\mathbf{p}^\top \mathbf{p}) = 0 \quad (10)$$

Does (10) \Rightarrow (9)? I.e. does index-1 model match original model?

Let's write $c(\mathbf{p}(t)) = \mathbf{p}(t)^\top \mathbf{p}(t) - L^2$ then index-1 model yields:

$$\ddot{c} = 0 \quad \Rightarrow \quad c(\mathbf{p}(t)) = c(\mathbf{p}(0)) + \dot{c}(\mathbf{p}(0), \dot{\mathbf{p}}(0))t$$

Hence index-1 model imposes $c(t) = 0$ for all t iff:

$$c(\mathbf{p}(0)) \quad \dot{c}(\mathbf{p}(0), \dot{\mathbf{p}}(0)) = 0$$

Consistency conditions - Pendulum example

Pendulum equations:

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \mathbf{p} - L^2\end{aligned}$$

Original model imposes

$$\mathbf{p}^\top \mathbf{p} = L^2 \quad (9)$$

$$\xrightarrow{\frac{d^2}{dt^2}}$$

$$\begin{aligned}\dot{\mathbf{p}} &= \mathbf{v} \\ \dot{\mathbf{v}} &= -\frac{1}{m} \mathbf{z} \mathbf{p} - \mathbf{g} \\ 0 &= \mathbf{p}^\top \dot{\mathbf{v}} + \mathbf{v}^\top \mathbf{v}\end{aligned}$$

Index-1 model imposes

$$\frac{d^2}{dt^2} (\mathbf{p}^\top \mathbf{p}) = 0 \quad (10)$$

Does (10) \Rightarrow (9)? I.e. does index-1 model match original model?

Let's write $c(\mathbf{p}(t)) = \mathbf{p}(t)^\top \mathbf{p}(t) - L^2$ then index-1 model yields:

$$\ddot{c} = 0 \quad \Rightarrow \quad c(\mathbf{p}(t)) = c(\mathbf{p}(0)) + \dot{c}(\mathbf{p}(0), \dot{\mathbf{p}}(0))t$$

Hence index-1 model imposes $c(t) = 0$ for all t iff:

$$c(\mathbf{p}(0)) \quad \dot{c}(\mathbf{p}(0), \dot{\mathbf{p}}(0)) = 0$$

These are called consistency conditions. Must be satisfied by $\mathbf{x}(0)$.

Consistency conditions