

# Pith Ball Lab

SPH-4UI - Tristan Simpson

December 1, 2022

---

## Purpose

The purpose of this lab was to calculate the electrostatic force of repulsion between a pith ball and an ebonite rod.

## Materials

1. **One** Pith ball electroscope.
2. **One** Pith ball.
3. **One** Protractor.
4. **One** Patch of fox fur.
5. **One** Strand of string.

## Procedure

1. The pith ball was attached to the string. The string's opposing side was attached to the electroscope.
2. The protractor was taped to the front of the electroscope.
3. The pith ball was given a neutral charge by having a group member touch it.
4. The ebonite rod was charged using the patch of fox fur.
5. The charged ebonite rod was held close to the pith ball.
6. The angle that the pith ball had moved to was noted.
7. The above steps were repeated another **Five** times for accuracy purposes.

## Calculations

Using a scale provided to our group in class we were able to measure the weight of the pith ball.

**Result:**

$$m_{ball} \approx 0.177 \times 10^{-4} \text{ kg}$$

By following the above procedures, we documented six angles at which the pith ball had moved to. To get the average of these angles, we added them all, then divided the result by six.

**Result:**


$$\begin{aligned} \theta_{average} &= \frac{(11.5^\circ) + (20^\circ) + (15^\circ) + (23^\circ) + (14^\circ) + (15^\circ)}{6} \\ &\approx 16.4^\circ \end{aligned}$$

To solve for the electrostatic force being exerted on the pith ball ( $F_q$ ) we need to solve for  $F_{T_x}$  since both  $x$  components ( $F_{T_x}$  and  $F_q$ ) are equal. To find  $F_{T_x}$  we solve for  $F_{T_y}$  then use *soh cah toa*

**Result:**

$$\begin{aligned} F_{net_y} &= F_{T_y} + (-F_g) \\ (m\ddot{\theta}^0) &= F_{T_y} + (-F_g) \end{aligned}$$

$$\therefore F_{T_y} = F_g$$



Since we're using the *opposite* and *adjacent* sides of our diagram ( $F_{T_y}$  and  $F_{T_x}$ ) we can rearrange the formula  $\tan \theta = \left( \frac{F_{T_y}}{F_{T_x}} \right)$  to get  $F_{T_x}$

**Final Result:**

$$\begin{aligned} F_q = F_{T_x} &= \frac{F_{T_y}}{\tan \theta_{average}} \\ &= \frac{m_{ball} \times g}{\tan \theta_{average}} \\ &= \frac{(0.177 \times 10^{-4})(9.81)}{\tan(16.4)} \\ &\approx 0.59 \times 10^{-2} \text{ N} \end{aligned}$$

## Sources of Error

### Errors

1. The force of friction was not accounted for.
2. The ebonite rod was not held directly perpendicular to the pith ball, creating more components.
3. The ebonite rod was not charged similarly to the previous tests.

### Solutions

1. Use materials that are very low in friction and/or account for friction in calculations.
2. This can be resolved by having something to support the position of the ebonite rod.
3. This can be resolved by using a timer to measure how long to rub the ebonite rod with fur.

## Conclusion

This lab efficiently demonstrated how to solve for the electrostatic force of repulsion between a pith ball and an ebonite rod. By the completion of the lab, it was determined that the electrostatic force of repulsion ( $F_q$ ) was approximately  $0.59 \times 10^{-2} N$

## Question

From the data collected during the lab, can you calculate the charge of the pithball? If not, why?

### Answer:

Calculating the numerical charge of the pith ball is not possible. This is because we don't have enough variables to support the equation:  $F_q = \left( \frac{kq_a q_b}{(r_{ab})^2} \right)$