SPLIT: Safety **P**rioritization for **L**ong COVID Drug Repurposing via a Causal **I**ntegrated **T**argeting Framework

Supplementary Materials

Contents

Part I: Supplementary Methods **4**

1 Causal Gene Discovery **4**
- 1.1 Transcriptome-Wide Mendelian Randomization (TWMR) . . . 4
- 1.2 Control Theory . . . 4
- 1.3 Differential Causal Effects . . . 4
- 1.4 Integration of Causal Evidence . . . 4

2 Clinical Trial Parsing and Knowledge Graph Integration **5**
- 2.1 Trial Data Acquisition . . . 5
- 2.2 Drug Entity Extraction and Standardization . . . 5
- 2.3 Eligibility Criteria Parsing and Population Characterization . . . 5
- 2.4 Outcome and Disease Normalization . . . 5
- 2.5 Trial Arm Feature Encoding . . . 6
- 2.6 Knowledge Graph Projection of Trial Arms . . . 6
- 2.7 Counterfactual Drug Replacement and Leakage Controls . . . 6

3 PlaNet Model Architecture and Training **6**
- 3.1 PlaNet Encoder and Self-Supervised Pretraining . . . 7
- 3.2 Trial-Arm Embedding Construction . . . 7
- 3.3 Integrating New Trial Arms for Inference . . . 7
- 3.4 Adverse Event and Safety Prediction . . . 7
- 3.5 Comparative Efficacy Prediction . . . 8
- 3.6 Cohort-Anchored Difference Metrics . . . 8

4 AE Burden Scoring and Weighting **9**
- 4.1 Weighted Adverse-Event Burden Score . . . 9
- 4.2 Severity Weights . . . 9
- 4.3 Long COVID Relevance Weights . . . 9
- 4.4 Global Safety Score . . . 10

4.5 Comparative Efficacy Score . . . 10

5 Composite Ranking Equations and Normalization 10
5.1 Min–Max Normalization . . . 10
5.2 Inverse AE Burden Normalization . . . 10
5.3 Composite Score Computation . . . 10

6 Avoid Lists: Definitions, Thresholds, and Stability 11
6.1 Primary Composite Avoid List . . . 11
6.2 Secondary Threshold-Based Screening . . . 11
6.3 Counterfactual-Delta Avoid List . . . 11
6.4 Cross-List Concordance and Sensitivity Analysis . . . 11

7 Rule-Based Filtering Categories 11
7.1 Hard-Block Rules . . . 11
7.2 Soft-Block Rules . . . 12
7.3 Keep-Hint Categories . . . 12

8 Pharmacology Enrichment Protocol 12
8.1 Data Sources . . . 12
8.2 Physicochemical Properties Collected . . . 13
8.3 Lipinski Rule-of-Five Assessment . . . 13

Part II: Supplementary Results 14

9 Key Terms and Definitions 14

10 Causal Driver Gene Evidence 14
10.1 Gene Counts by Method and Evidence Category . . . 14
10.2 Convergent Driver Genes . . . 14

11 Drug–Gene Mapping Details 15

12 PlaNet Counterfactual Predictions 16

13 Adverse Event Signatures by Drug 16

14 Avoid List Structure and Concordance 17
14.1 Avoid List Summary . . . 17
14.2 Concordance Metrics . . . 17

15 Dimensional Risk Profiles 17

16 Secondary Screening: Drugs Failing Multiple Thresholds 18

17 Shared Deprioritized Drugs Across Cohorts 19

18 Full Panel Score Distributions **19**

Part I: Supplementary Methods

1 Causal Gene Discovery

To obtain a robust set of putative Long COVID driver genes, we combined three orthogonal causal inference approaches: Transcriptome-Wide Mendelian Randomization (TWMR), Control Theory (CT), and Differential Causal Effects (DCE). All methods were implemented using harmonized identifiers, consistent multiple-testing correction, and standardized quality-control procedures. The complete annotated gene list with evidence labels is provided in Supplementary Section S1.

1.1 Transcriptome-Wide Mendelian Randomization (TWMR)

TWMR was used to identify genes whose genetically predicted expression levels exert a causal effect on Long COVID risk [3]. We integrated Genome-Wide Association Study (GWAS) summary statistics for Long COVID with quantitative trait locus of cis expression (cis-eQTL) data (GTEx v8) [11]. Genes were retained as drivers supported by TWMR when they showed: (i) consistent direction and magnitude among robust estimators, (ii) false discovery rate (FDR) < 0.05, and (iii) no evidence of substantial pleiotropic distortion based on heterogeneity and intercept tests. This step prioritizes targets with human genetic support, increasing the likelihood that modulating their expression affects Long COVID biology rather than reflecting confounding.

1.2 Control Theory

To capture regulatory importance at the system level, we applied controllability analysis on a curated human protein–protein interaction network following Vinayagam *et al.* [14, 11]. We retained indispensable nodes whose removal increases the number of driver nodes required to control the network. Indispensable nodes have been reported to be enriched among essential genes, disease genes, and drug targets associated with severe adverse events [14, 10].

1.3 Differential Causal Effects

To identify how regulatory pathways are causally altered in Long COVID, we implemented the DCE framework [4, 12]. RNA-seq data from individuals with Long COVID and recovered controls were analyzed within curated pathways (KEGG) to estimate directed causal effects between interacting genes separately in the Long COVID and control groups, while adjusting for confounders. We then computed differential causal effects across groups and ranked genes by their contribution to pathway perturbation. Genes with strong and consistent pathway reorganization signatures across multiple pathways were designated as DCE-supported drivers.

1.4 Integration of Causal Evidence

We integrated the TWMR, CT, and DCE outputs to obtain a consolidated set of Long COVID driver genes. Genes supported by at least one method and not contradicted by sensitivity

analyses were retained. This integration captures genetically anchored risk (TWMR), network-level regulatory leverage (CT), and condition-specific pathway reorganization (DCE).

2 Clinical Trial Parsing and Knowledge Graph Integration

We implemented an automated workflow that (i) parses clinical trial records into structured trial-arm representations, (ii) resolves drugs and cohort descriptors to canonical biomedical identifiers, (iii) projects each arm into a typed-edge knowledge-graph representation, and (iv) supports counterfactual drug substitution while keeping the cohort and protocol constant.

2.1 Trial Data Acquisition

Clinical trial records were retrieved from ClinicalTrials.gov using the public API (version 2). For each trial, we extracted the NCT identifier, arm group definitions, interventions, conditions, eligibility criteria, primary and secondary outcomes, phase, enrollment, and demographic constraints (sex and age ranges). Fields were normalized into a consistent internal schema for downstream parsing and graph construction. Raw clinical trial data for each cohort are provided in Supplementary Section S4 (NCT04809974) and Supplementary Section S5 (NCT04880161).

2.2 Drug Entity Extraction and Standardization

Drug mentions in intervention descriptions were processed using MedEx [15] to extract medication entities and associated attributes when available. Extracted entities were mapped to canonical drug identifiers using a multi-stage strategy supporting exact matching, synonym lookup, and fuzzy matching with configurable similarity thresholds. Candidate replacement drugs were optionally validated against the PlaNet drug vocabulary prior to counterfactual generation; drugs not found were skipped.

2.3 Eligibility Criteria Parsing and Population Characterization

Eligibility criteria were parsed using Criteria2Query [16] to extract structured inclusion/exclusion constraints. Extracted concepts were mapped to the Unified Medical Language System (UMLS) using concept search and Term Frequency-Inverse Document Frequency (TF–IDF) matching. Concepts were propagated through the UMLS hierarchy to enrich cohort representation with clinically relevant parent concepts.

2.4 Outcome and Disease Normalization

Trial conditions were assigned to controlled disease identifiers, including Medical Subject Headings (MeSH) when available, to align trial context with the disease ontology represented in the knowledge graph. Primary outcomes were parsed and normalized to create structured outcome representations compatible with downstream feature encoding.

2.5 Trial Arm Feature Encoding

For each arm, we constructed an arm text document by concatenating the intervention name/description, condition text, primary outcome text, arm label, trial attributes summary, and eligibility criteria text. In parallel, we derived structured trial attributes including phase encoding, logarithmic transformation of enrollment magnitude, sex eligibility, and age-range indicators. These attributes provide consistent low-dimensional covariates that complement text features in downstream embedding models. Parsed trial representations, including arm texts, eligibility criteria, and cohort descriptors, are provided in Supplementary Section S6 (NCT04809974) and Supplementary Section S7 (NCT04880161).

2.6 Knowledge Graph Projection of Trial Arms

Each parsed trial arm was projected into the PlaNet KG by linking intervention, condition, population descriptors, and results to KG entities through typed relations. Arm indexing and labeling were resolved using multiple metadata fields and, when unavailable, by falling back to arm-group order. KG entity merges were handled using union-find over merge relations, and entity-to-KG-ID resolution used the PlaNet entity vocabulary.

2.7 Counterfactual Drug Replacement and Leakage Controls

To evaluate candidate drugs within fixed cohort contexts, we generated counterfactual trial variants by substituting the experimental intervention while preserving all other protocol elements (condition, eligibility criteria, cohort descriptors, enrollment, demographics, and comparator structure). Replacement proceeded by identifying a non-placebo experimental intervention, using canonicalized names and identifiers when available. If a base trial lacked an explicit drug-annotated intervention, a minimal synthetic experimental anchor was created from the inferred non-placebo arm to allow replacement.

To reduce information leakage from real trial outcomes into counterfactual representations, counterfactual trials were cleaned by removing results-derived fields and replacing summaries with a neutral counterfactual statement. Arm labels and descriptions were updated to remove references to the original drug. These steps ensure that predictions are driven by cohort context and by the substituted intervention, rather than by accidental carryover from the original trial.

3 PlaNet Model Architecture and Training

We applied PlaNet, a geometric deep learning framework trained on a large-scale clinical knowledge graph (KG), to predict cohort-anchored adverse-event profiles (AE), safety (S), and comparative efficacy (E) for each parsed trial arm [1]. In PlaNet, clinical trials are represented as trial-arm nodes connected via typed edges to protocol-defining entities, including administered drugs, investigated diseases/conditions, primary outcomes, and structured population descriptors derived from inclusion/exclusion criteria. This KG is integrated with biomedical background subnetworks (drug similarity/class hierarchies, disease hierarchies, drug–protein interactions, protein–protein interactions, and function ontologies), enabling message passing over

higher-order relations among drugs, diseases, populations, and outcomes.

3.1 PlaNet Encoder and Self-Supervised Pretraining

PlaNet learns low-dimensional embeddings for all KG entities using a relational graph convolutional network (R-GCN) encoder operating over typed edges. Entity representations are initialized from type-specific attributes and mapped into a shared embedding space via entity-type-specific linear projections; message passing then propagates information across multi-hop neighborhoods to produce final entity embeddings.

To leverage the large amount of unlabeled KG structure, PlaNet is first pretrained with a self-supervised link-prediction objective: random edges are masked/dropped from k-hop subgraphs and the model is trained to reconstruct missing edges using a bilinear scoring decoder (DistMult) over the learned embeddings. This pretraining encourages embeddings that capture both protocol-level trial structure and biomedical context, improving generalization to unseen drug–disease combinations.

3.2 Trial-Arm Embedding Construction

For each trial arm, PlaNet constructs a protocol-level representation by combining (i) textual arm descriptors (arm title/description, intervention details, trial summary, eligibility criteria, and outcome text) embedded with a biomedical language model (BERT-family encoders), and (ii) structured trial covariates (phase, enrollment, age range, and sex eligibility). These arm attributes serve as input features for trial-arm nodes and are integrated with neighborhood information via the KG network, producing an embedding simultaneously conditioned on cohort context and tested intervention.

3.3 Integrating New Trial Arms for Inference

For inference on newly parsed arms (including counterfactual substitutions), we instantiate trial-arm nodes with their arm text embeddings, structured covariates, and KG edges (drug, disease, outcome, and population links). Predictions are then obtained by applying the pretrained PlaNet encoder and corresponding task heads to the resulting arm embeddings, without retraining.

3.4 Adverse Event and Safety Prediction

PlaNet's safety and adverse-event tasks are defined relative to a placebo (no-treatment) baseline for the same disease and population context. Placebo arms are aggregated to estimate a cohort-anchored prior probability of adverse events for each disease (and associated population), representing expected events in the absence of active intervention. Given an intervention arm, PlaNet predicts whether (and how strongly) an adverse event is enriched compared to this placebo prior, implemented via contingency-table comparisons against the estimated placebo distribution and an odds-ratio enrichment threshold [1].

We query two PlaNet safety-related prediction heads: (i) a binary serious adverse event (SAE) head that outputs a probability of SAE occurrence/enrichment for a given arm; and

(ii) a multi-task AE-category head that outputs calibrated probabilities over a standardized adverse-event vocabulary defined at the Medical Dictionary for Regulatory Activities (MedDRA) Preferred Term (PT) level [8]. For each arm i and adverse-event category k, PlaNet produces:

$$p_{i,k} = \Pr(\text{AE category } k \text{ is enriched in arm } i \text{ relative to placebo}) \in [0, 1].$$

To align with our interpretation where higher values indicate safer profiles, we define the global safety score as the complement of SAE risk:

$$S_{\text{raw},i} = 1 - p_i^{\text{SAE}},$$

where p_i^{SAE} is the SAE-head probability for arm i. A larger S_{raw} indicates safer predicted profiles, while a larger $p_{i,k}$ indicates higher predicted risk of AE enrichment for category k.

3.5 Comparative Efficacy Prediction

PlaNet's efficacy task is trained on survival-type endpoints extracted from completed clinical trials. For trials containing two or more arms with comparable outcome definitions, PlaNet is fine-tuned to predict which arm is expected to achieve a more favorable survival outcome. When a placebo/comparator arm could be reliably identified, we computed cohort-specific comparative efficacy as:

$$E_{\text{raw}} = \Pr(\text{active arm is superior to comparator}), \qquad E_{\text{raw}} \in [0, 1].$$

Values above 0.5 indicate higher predicted probability of benefit under the active intervention, conditional on cohort and protocol context encoded in the trial-arm representation. When placebo identification was ambiguous or a valid pairing could not be established, efficacy was not calculated, and subsequent evaluation relied only on safety and adverse-event components.

3.6 Cohort-Anchored Difference Metrics

For each metric $X \in \{S_{\text{raw}}, AE_{\text{score}}, E_{\text{raw}}\}$:

$$\Delta X = X_{\text{counterfactual}} - X_{\text{reference}}.$$

Negative ΔS or ΔE indicates a worse profile under substitution; positive ΔAE indicates increased toxicity burden.

4 AE Burden Scoring and Weighting

4.1 Weighted Adverse-Event Burden Score

To summarize toxicity while prioritizing clinically severe and Long COVID-relevant events, we computed a weighted probability for each event k in arm i:

$$\hat{p}_{i,k} = p_{i,k} \cdot s_k \cdot \ell_k,$$

where s_k is a CTCAE-aligned severity weight and ℓ_k is a Long COVID relevance weight.

The weighted AE burden score:

$$AE_{\text{score},i} = \sum_k \hat{p}_{i,k}.$$

Larger values indicate worse predicted AE burden.

4.2 Severity Weights

Severity weights (s_k) were assigned based on CTCAE-aligned severity categories [8]:

Severity Category	**Weight (s_k)**
Mild	0.25
Moderate	0.50
Severe	0.75
Life-threatening	1.00
Death	1.20

Table 1: CTCAE-aligned severity weights for adverse event scoring.

4.3 Long COVID Relevance Weights

Relevance weights (ℓ_k) were assigned based on clinical relevance to Long COVID symptom domains:

Relevance Category	**Weight (ℓ_k)**
Core Long COVID symptoms	1.50
Important comorbidity	1.25
General	1.00

Table 2: Long COVID relevance weights for adverse event scoring.

Core Long COVID symptoms include fatigue, cognitive dysfunction ("brain fog"), dyspnea, chest pain, palpitations, postural symptoms, and persistent anosmia/dysgeusia. Important comorbidities include cardiovascular events, thromboembolic events, and metabolic disturbances frequently observed in Long COVID populations.

When a weight could not be confidently assigned, default values of $s_k = 0.50$ and $\ell_k = 1.00$ were applied.

4.4 Global Safety Score

PlaNet provides $S_{\text{raw}} \in [0, 1]$, where higher values indicate safer profiles. For cohort-internal ranking, we optionally applied min–max normalization.

4.5 Comparative Efficacy Score

For trials with valid experimental–comparator pairings, $E_{\text{raw}} \in [0, 1]$, where values above 0.5 indicate predicted benefit. Min–max normalization was optionally applied for ranking.

5 Composite Ranking Equations and Normalization

5.1 Min–Max Normalization

For cohort-internal ranking, raw scores were optionally normalized using min–max scaling within each cohort:

$$\tilde{X} = \frac{X_{\text{raw}} - X_{\min}}{X_{\max} - X_{\min}},$$

where $X \in \{S, E\}$ and $X_{\min}$, $X_{\max}$ are the minimum and maximum values observed within the cohort.

5.2 Inverse AE Burden Normalization

To convert AE burden (where larger = worse) into a scale compatible with safety and efficacy (where larger = better), we computed the inverse AE burden:

$$AE^{-1} = \frac{1}{AE_{\text{score}} + \epsilon}, \qquad \epsilon = 10^{-9},$$

followed by cohort-level min–max normalization:

$$\widehat{AE^{-1}} = \frac{AE^{-1} - (AE^{-1})_{\min}}{(AE^{-1})_{\max} - (AE^{-1})_{\min}}.$$

5.3 Composite Score Computation

For screening, we combined normalized safety ($\tilde{S}$), efficacy ($\tilde{E}$), and inverse AE burden ($\widehat{AE^{-1}}$):

$$C = w_S \cdot \tilde{S} + w_E \cdot \tilde{E} + w_{AE} \cdot \widehat{AE^{-1}},$$

with default weights $w_S = 0.4$, $w_E = 0.4$, and $w_{AE} = 0.2$, constrained such that $w_S + w_E + w_{AE} = 1$.

When comparative efficacy (E) was unavailable, weights were renormalized over available components:

$$C = \frac{w_S}{w_S + w_{AE}} \cdot \tilde{S} + \frac{w_{AE}}{w_S + w_{AE}} \cdot \widehat{AE^{-1}} = 0.67 \cdot \tilde{S} + 0.33 \cdot \widehat{AE^{-1}}.$$

6 Avoid Lists: Definitions, Thresholds, and Stability

6.1 Primary Composite Avoid List

Drugs were ranked by composite score C, with lowest-scoring candidates forming the avoid list. We report three stringency levels: q05 (worst 5%), q10 (worst 10%, default), and q20 (worst 20%).

6.2 Secondary Threshold-Based Screening

As an independent safety check, we flagged drugs with extreme individual metrics (q20 thresholds): AE-extreme (top 20% of AE_{score}), safety-extreme (bottom 20% of S_{raw}), and efficacy-extreme (bottom 20% of E_{raw}). We tracked the number of triggered flags per drug ($n_{\text{extremes}} \in \{1, 2, 3\}$).

6.3 Counterfactual-Delta Avoid List

We identified drugs performing worse than the original intervention under substitution. Candidates were flagged as delta-worse when $\Delta S < 0$, $\Delta E < 0$, or $\Delta AE > 0$. The delta avoid list includes candidates in the worst decile for any metric, with multi-flag tracking ($n_{\text{delta}} \in \{1, 2, 3\}$).

6.4 Cross-List Concordance and Sensitivity Analysis

Candidates appearing across multiple lists (composite, threshold, and delta) represent higher-confidence deprioritization targets. We assessed sensitivity by repeating analyses across q05, q10, and q20 thresholds. Candidates stable across thresholds form a conservative "avoid" core; threshold-dependent candidates define a broader caution set.

7 Rule-Based Filtering Categories

7.1 Hard-Block Rules

Candidates were automatically excluded if they matched any of the following criteria:

- Oncology-specific agents (chemotherapeutics, targeted cancer therapies) without established non-oncology indications
- Acute-only interventions (e.g., thrombolytics, acute sedatives) unsuitable for chronic/outpatient Long COVID management
- Withdrawn or revoked drugs
- Agents with unacceptable safety profiles for the target population (e.g., known teratogens when reproductive-age populations are included)

7.2 Soft-Block Rules

Candidates were flagged for manual review (but not automatically excluded) if they:

- Require intensive monitoring (e.g., narrow therapeutic index, mandatory lab monitoring)
- Have significant drug–drug interaction potential relevant to common Long COVID comedications
- Are associated with known risks in Long COVID-relevant organ systems (cardiac, pulmonary, neurological) but may still offer net benefit
- Have limited safety data in post-viral or chronic fatigue populations

7.3 Keep-Hint Categories

Candidates were tagged for preserved visibility if they aligned with Long COVID-relevant therapeutic themes:

- **Cardiopulmonary/vascular**: Agents addressing cardiovascular symptoms, microvascular dysfunction, or endothelial health
- **Respiratory anti-inflammatory**: Inhaled or systemic anti-inflammatory agents for persistent respiratory symptoms
- **Neurocognitive**: Agents targeting cognitive dysfunction, neuroinflammation, or autonomic dysregulation
- **Metabolic/fibrotic**: Agents addressing metabolic dysregulation or fibrotic sequelae
- **Autonomic/POTS-related**: Agents for postural orthostatic tachycardia syndrome and related dysautonomia
- **Low-risk supportive**: Well-tolerated supportive agents (e.g., supplements, low-risk symptomatic treatments)

8 Pharmacology Enrichment Protocol

8.1 Data Sources

Enrichment data were collected programmatically from:

- **PubChem** [6, 5]: Cross-references, physicochemical properties, compound identifiers
- **ChEMBL** [2]: Potency/binding data (IC_{50}, EC_{50}, K_i, K_d), mechanism of action, target annotations
- **Open Targets** [9]: Disease associations, target tractability, clinical precedence
- **DailyMed** [13]: FDA label information, boxed warnings, contraindications

8.2 Physicochemical Properties Collected

- Molecular weight (MW)
- Lipophilicity (XLogP, ALogP)
- Topological polar surface area (TPSA)
- Hydrogen bond donors (HBD)
- Hydrogen bond acceptors (HBA)
- Rotatable bond count (flexibility)
- Heavy atom count

8.3 Lipinski Rule-of-Five Assessment

Developability was assessed using the Lipinski Rule-of-Five [7]. Violations were counted for:

- MW > 500 Da
- XLogP > 5
- HBD > 5
- HBA > 10

Compounds with ≤ 1 violation were considered to have favorable oral bioavailability profiles.

Part II: Supplementary Results

9 Key Terms and Definitions

For clarity, we define key terms used throughout the main text and supplementary materials:

- **Prediction**: PlaNet outputs for S, E, and AE probabilities in a given cohort context.
- **Counterfactual substitution**: Replacing the trial intervention while maintaining the cohort, protocol, and comparator fixed.
- **Evaluation**: The process of testing SPLIT as a framework, using counterfactual substitution to assess whether predicted scores distinguish drugs with different risk profiles.
- **Screening**: Applying predictions and scoring across a drug panel to generate ranks.
- **AE burden**: A weighted score combining individual predicted AE probabilities with severity and Long COVID relevance weights, summarizing the overall predicted toxicity load of a candidate.
- **Composite score**: A single ranking metric that combines normalized S, E, and inverse AE burden to summarize the overall risk-benefit profile of each candidate.
- **Deprioritization**: Labeling high risk or low benefit candidates as "avoid" based on ranks, thresholds, or deltas relative to the original trial drug.
- **Template cohort**: The primary clinical trial context (NCT04809974, cognitive phenotype) in which counterfactual substitutions are performed.
- **Sensitivity cohort**: A second clinical trial context (NCT04880161, respiratory phenotype) used to test whether deprioritization patterns generalize across phenotypes.

10 Causal Driver Gene Evidence

10.1 Gene Counts by Method and Evidence Category

Table 3 summarizes the distribution of causal driver genes across the three inference methods. CT contributed the largest set (1,641 genes), reflecting the scale of the protein–protein interaction network, while TWMR (50 genes) and DCE (45 genes) provided complementary, more focused evidence.

10.2 Convergent Driver Genes

Table 4 lists all 11 genes supported by two or more causal inference methods.

Table 3: **Summary of causal driver gene evidence across methods:** Mendelian Randomization (MR), Control Theory (CT), and Differential Causal Effects (DCE). Full gene list in Supplementary Section S1.

Evidence category	**Genes (n)**
Total causal driver genes	1,725
CT supported (any)	1,641
MR supported (any)	50
DCE supported (any)	45
Single method support	
CT only	1,631
MR only	45
DCE only	38
Support from multiple methods	
CT + DCE	6
CT + MR	4
DCE + MR	1
Convergent total	11

Table 4: **Long COVID causal driver genes supported by ≥2 causal inference methods.** Complete annotations in Supplementary Section S1.

Gene	**Supporting methods**
ERCC3	CT; MR
HIF1A	CT; DCE
IMMT	CT; MR
LEF1	CT; DCE
MED19	CT; MR
MYC	CT; DCE
NDUFA6	CT; MR
PNPO	DCE; MR
POMC	CT; DCE
STAT3	CT; DCE
TP53	CT; DCE

11 Drug–Gene Mapping Details

Table 5 presents the 10 drugs with the highest causal gene coverage. The dominance of oncology kinase inhibitors in this list motivated the explicit deprioritization step in the framework. Complete mapping tables are provided in Supplementary Section S2.

Table 5: **Top 10 drugs by number of mapped Long COVID causal gene targets.** Full mapping table in Supplementary Section S2.

Drug	**CG (n)**	**CG Methods**	**Drug–target sources**
Imatinib	210	CT; DCE	ChEMBL; DrugBank; DrugCentral; GtoPdb
Sirolimus	206	CT; DCE	ChEMBL; DrugCentral; GtoPdb
Palbociclib	201	CT	ChEMBL; DrugBank; DrugCentral; GtoPdb; TTD
Sunitinib	194	CT	ChEMBL; DrugBank; DrugCentral; GtoPdb
Sorafenib	191	CT; DCE	ChEMBL; DrugBank; DrugCentral; GtoPdb; TTD
Nilotinib	190	CT	ChEMBL; DrugBank; DrugCentral; GtoPdb
Bosutinib	189	CT; DCE	ChEMBL; DrugBank; DrugCentral; GtoPdb; TTD
Gefitinib	189	CT	ChEMBL; DrugBank; DrugCentral; GtoPdb; TTD
Erlotinib	186	CT	ChEMBL; DrugBank; DrugCentral; GtoPdb; TTD
Vandetanib	185	CT	ChEMBL; DrugBank; DrugCentral; GtoPdb; TTD

Abbreviations: n, Number; CG, Causal Gene Targets; CT, Control Theory; DCE, Differential Causal Effects.

12 PlaNet Counterfactual Predictions

Table 6 provides the full PlaNet predictions for the template cohort (NCT04809974), including the original trial drug (Niagen) and three counterfactual substitutions. These data underlie the multi-panel Figure 2 in the main text. Complete prediction outputs are provided in Supplementary Section S8.

Table 6: **PlaNet predictions for the NCT04809974 template cohort, original trial drug (Niagen), and counterfactual substitutions.** S and AE outputs are defined relative to a placebo baseline in the same population context, whereas the Δ values quantify differences between candidates within the cohort (candidate minus Niagen). S scores range from 0 to 1 (higher = safer); E represents $E = \Pr(\text{Drug} > \text{Placebo})$. Top AE is the highest probability event. Full AE profiles in Supplementary Section S3.

Drug	CG	S	ΔS	E	ΔE	Top AE (P)
Niagen (Original)	—	0.549	—	0.519	—	Hypertensive diseases (0.197)
Leronlimab	1	0.441	–0.108	0.521	+0.002	Hypertensive diseases (0.175)
Vortioxetine	11	0.332	–0.217	0.521	+0.002	Nausea (0.234)
Ritonavir	70	0.215	–0.334	0.527	+0.009	Diarrhea (0.192)

Abbreviations: CG, number of causal genes targeted; S, safety score; E, efficacy probability; AE, adverse event; P, probability.

13 Adverse Event Signatures by Drug

Table 7 presents the top 10 predicted AE for each counterfactual scenario in the NCT04809974 cohort, enabling distinction between drug-related toxicity and cohort background signals. Complete probability tables are provided in Supplementary Section S3, with mapped AE tables in Supplementary Section S10.

Table 7: **Top 10 predicted adverse events per scenario in the NCT04809974 cohort.** Probabilities represent PlaNet predictions. Full AE tables in Supplementary Section S3.

Rank	Niagen (Original)	P	Rank	Ritonavir	P
1	Hypertensive diseases	0.197	1	Diarrhoea	0.192
2	Diarrhoea	0.174	2	Hypertensive diseases	0.136
3	Confusion NOS	0.173	3	Acrodermatitis	0.125
4	Bronchopneumonia	0.157	4	Hypokalaemia	0.100
5	Hypokalaemia	0.139	5	Asthenia	0.099
6	Hyperglycaemia	0.122	6	Dyspnoea	0.095
7	Nausea with vomiting	0.118	7	Bronchopneumonia	0.084
8	Asthenia	0.115	8	Confusion NOS	0.083
9	Nausea	0.113	9	Dehydration	0.082
10	Dyspnoea	0.111	10	Headache	0.075
Rank	**Vortioxetine**	**P**	**Rank**	**Leronlimab**	**P**
1	Nausea	0.234	1	Hypertensive diseases	0.175
2	Confusion NOS	0.182	2	Asthenia	0.157
3	Diarrhoea	0.177	3	Diarrhoea	0.144
4	Nausea with vomiting	0.168	4	Hyperglycaemia	0.137
5	Insomnia	0.162	5	Dyspnoea	0.126
6	Constipation	0.155	6	Headache	0.124
7	Hypertensive diseases	0.148	7	Bronchopneumonia	0.118
8	Anorexia	0.142	8	Acrodermatitis	0.118
9	Asthenia	0.141	9	Hypokalaemia	0.111
10	Weight below normal	0.089	10	Confusion NOS	0.111

14 Avoid List Structure and Concordance

14.1 Avoid List Summary

Table 8 summarizes the structure and output of each deprioritization category for both cohorts.

Table 8: **Summary of deprioritization avoid lists for both cohorts.** Quantile thresholds (q05, q10, q20) indicate the percentage of worst ranked drugs captured. NCT04809974 deprioritization is driven by S/AE; NCT04880161 deprioritization is driven by E. Full avoid lists in Supplementary Section S14 (NCT04809974) and Supplementary Section S15 (NCT04880161).

List type	Threshold	C1	C2	Description
Primary composite (worst overall risk–benefit)				
Primary q05	Worst 5%	82	82	Core deprioritized set
Primary q10	Worst 10%	163	163	Default threshold
Primary q20	Worst 20%	325	325	Extended set
Secondary (extreme individual metrics)				
Threshold q20	Any extreme	855	803	Drugs with $\geq$1 extreme flag
AE extreme	Top 20% AE	326	326	High predicted toxicity burden
Safety extreme	Bottom 20% S	325	325	Low predicted S score
Efficacy extreme	Bottom 20% E	325	325	Low predicted E
Multiple extremes	$\geq$2 flags	121	168	Failed multiple thresholds
Counterfactual delta (worse than original trial drug)				
Delta q10	Any worse	457	412	Drugs worse than original
ΔAE worse	–	163	163	Higher AE burden
ΔS worse	–	163	163	Lower S
ΔE worse	–	164	163	Lower E
Concordance across lists (q10 primary)				
Primary $\cap$ Secondary	–	158 (96.9%)	162 (99.4%)	Overlap with threshold list
Primary $\cap$ Delta	–	113 (69.3%)	131 (80.4%)	Overlap with delta list
All three lists	–	113	131	Highest confidence targets

Abbreviations: C1, Cohort 1 (NCT04809974); C2, Cohort 2 (NCT04880161). **Note:** Original trial drugs are Niagen (NCT04809974) and Ampion (NCT04880161). Panel sizes: NCT04809974 = 1,625 drugs; NCT04880161 = 1,630 drugs.

14.2 Concordance Metrics

Table 9 provides detailed concordance metrics across avoid list categories.

Table 9: **Concordance across avoid lists.** The respiratory cohort shows higher concordance across all metrics, reflecting more consistent deprioritization signals when E provides strong discrimination.

Concordance metric	NCT04809974	NCT04880161	Difference
Primary list size	163 (10.0%)	163 (10.0%)	—
Secondary list size	855 (52.6%)	803 (49.4%)	–3.2%
Delta list size	457 (28.1%)	412 (25.4%)	–2.7%
Primary $\cap$ Secondary	158 (96.9%)	162 (99.4%)	+2.5%
Primary $\cap$ Delta	113 (69.3%)	131 (80.4%)	+11.1%
All three lists	113	131	+18 drugs

15 Dimensional Risk Profiles

Table 10 presents drugs with the most extreme unfavorable values on each individual risk dimension, revealing how each metric discriminates differently across the two cohorts.

Table 10: **Bottom 5 drugs by individual risk dimension for both cohorts.** Key finding: E shows minimal discrimination in the cognitive cohort (spread = 0.002) but strong discrimination in the respiratory cohort (spread = 0.015 among worst 10). AE burden is approximately 2× higher in the respiratory cohort. Full rankings in Supplementary Section S12 and Supplementary Section S13.

NCT04809974 (Cognitive)				NCT04880161 (Respiratory)			
Worst by AE Burden							
R	**Drug**	**AE**	**Class**	**R**	**Drug**	**AE**	**Class**
1	Lenvatinib	14.97	Multi-kinase	1	Cobimetinib	30.83	MEK inh.
2	Carfilzomib	14.85	Proteasome	2	Carfilzomib	30.11	Proteasome
3	Cobimetinib	14.72	MEK inh.	3	Lumacaftor	29.88	CFTR corr.
4	Regorafenib	14.24	Multi-kinase	4	Selumetinib	28.92	MEK inh.
5	Axitinib	13.72	VEGFR inh.	5	Lenvatinib	28.85	Multi-kinase
Range: 12.8–15.0				*Range: 25.2–30.8 (2× higher)*			
Worst by Safety							
R	**Drug**	**S**	**Class**	**R**	**Drug**	**S**	**Class**
1	Benzoyl peroxide	0.086	Topical	1	Benzoyl peroxide	0.217	Topical
2	Terconazole	0.094	Topical antifung.	2	Glucosamine	0.226	Nutraceutical
3	Cetylpyridinium	0.097	Topical antisep.	3	Cetylpyridinium	0.233	Topical antisep.
4	Donepezil	0.116	AChE inh.	4	Salsalate	0.243	NSAID
5	Scopolamine	0.116	Anticholinergic	5	Folic acid	0.259	Vitamin
Range: 0.09–0.12 (very low)				*Range: 0.22–0.27 (higher floor)*			
Worst by Efficacy							
R	**Drug**	**E**	**Class**	**R**	**Drug**	**E**	**Class**
1	Pivampicillin	0.514	Penicillin	1	Olaparib	0.009	PARP inh.
2	Upadacitinib	0.515	JAK inh.	2	Temozolomide	0.014	Alkylating
3	Chlorphentermine	0.515	Anorectic	3	Enzalutamide	0.015	Antiandrogen
4	Bithionol	0.515	Antiparasitic	4	Alectinib	0.017	ALK inh.
5	Estradiol cyp.	0.515	Estrogen	5	Trametinib	0.019	MEK inh.
Range: 0.514–0.516 (no discrim.)				*Range: 0.009–0.024 (strong discrim.)*			

Abbreviations: R, rank; inh., inhibitor; corr., corrector; antifung., antifungal; antisep., antiseptic; cyp., cypionate; discrim., discrimination.

16 Secondary Screening: Drugs Failing Multiple Thresholds

Table 11 presents the top 10 drugs failing multiple thresholds in both cohorts. Oncology kinase inhibitors dominated this subset (70% of flagged drugs), with the respiratory cohort showing 2–3× higher AE values and lower E.

Table 11: **Top 10 drugs failing multiple thresholds in both cohorts.** All listed drugs were flagged on both AE burden and E. Oncology kinase inhibitors dominate both lists, but failure modes differ quantitatively: the respiratory cohort shows 2–3× higher AE values and lower E. Full screening in Supplementary Section S14 and Supplementary Section S15.

NCT04809974 (Cognitive)					NCT04880161 (Respiratory)				
R	**Drug**	**AE**	**S**	**E**	**R**	**Drug**	**AE**	**S**	**E**
1	Cobimetinib	5.5	.51	.519	1	Carfilzomib	13.0	.37	.045
2	Regorafenib	5.2	.55	.517	2	Selumetinib	12.0	.35	.029
3	Infigratinib	4.6	.38	.519	3	Cobimetinib	11.8	.41	.039
4	Ruxolitinib	4.6	.53	.519	4	Pomalidomide	11.6	.37	.053
5	Tazemetostat	4.3	.36	.519	5	Lenvatinib	11.6	.40	.030
6	Ponatinib	4.2	.41	.517	6	Ixazomib	10.9	.37	.045
7	Tepotinib	4.1	.41	.518	7	Trametinib	10.4	.73	.019
8	Binimetinib	4.0	.39	.518	8	Momelotinib	10.3	.37	.034
9	Encorafenib	3.8	.45	.517	9	Ruxolitinib	9.8	.46	.058
10	Pexidartinib	3.7	.48	.519	10	Binimetinib	9.5	.41	.032
Multiple extremes total: 121 drugs (7.4%)					*Multiple extremes total: 168 drugs (10.3%)*				

17 Shared Deprioritized Drugs Across Cohorts

Table 12 lists the 18 drugs deprioritized in both the cognitive and respiratory cohorts. All shared drugs showed improved S in the respiratory cohort ($\Delta S = +0.06$ to $+0.16$) but reduced E ($\Delta E = -0.38$ to -0.44), indicating convergent deprioritization from different failure modes.

Table 12: **Drugs deprioritized in both cohorts (n=18).** All shared drugs show improved S but reduced E in the respiratory cohort, indicating convergent deprioritization from different failure modes: S driven in the cognitive cohort, E driven in the respiratory cohort.

	NCT04809974		NCT04880161		Δ (Resp–Cog)	
Drug	**S**	**E**	**S**	**E**	**ΔS**	**ΔE**
Secnidazole	0.124	0.516	0.285	0.122	+0.16	−0.39
Ethionamide	0.196	0.517	0.306	0.129	+0.11	−0.39
Tavaborole	0.168	0.519	0.309	0.119	+0.14	−0.40
Isoflurophate	0.240	0.517	0.347	0.109	+0.11	−0.41
Isradipine	0.220	0.517	0.340	0.124	+0.12	−0.39
Butoconazole	0.169	0.519	0.303	0.131	+0.13	−0.39
Lincomycin	0.234	0.518	0.342	0.123	+0.11	−0.40
Hydroxocobalamin	0.191	0.520	0.329	0.116	+0.14	−0.40
Uridine triacetate	0.243	0.518	0.318	0.134	+0.08	−0.38
Olmutinib	0.273	0.518	0.345	0.116	+0.07	−0.40
Canagliflozin	0.196	0.518	0.330	0.123	+0.13	−0.40
Ethosuximide	0.228	0.518	0.343	0.115	+0.12	−0.40
Linagliptin	0.231	0.518	0.357	0.109	+0.13	−0.41
Capmatinib	0.312	0.518	0.380	0.108	+0.07	−0.41
Phenazopyridine	0.257	0.517	0.350	0.117	+0.09	−0.40
Permethrin	0.242	0.519	0.298	0.076	+0.06	−0.44
Entrectinib	0.324	0.519	0.403	0.088	+0.08	−0.43
Copanlisib	0.341	0.519	0.416	0.085	+0.08	−0.43

18 Full Panel Score Distributions

Figure 1 compares S, E, and AE burden distributions across the complete screening panel for both cohorts, illustrating the different discriminatory behaviors that drive the cohort specific deprioritization mechanisms described in the main text.




Figure 1: **Complete panel score distributions across cohorts.** Left: S scores show similar distributions. Center: E scores show fundamentally different patterns: the cognitive cohort (blue) is tightly clustered around 0.52, while the respiratory cohort (coral) is spread across 0.01–0.64. Right: AE burden distributions with the respiratory cohort showing higher extreme values. This 37-fold difference in E spread explains why S drives deprioritization in the cognitive cohort while E drives deprioritization in the respiratory cohort.

References

[1] Matko Brbic, Michihiro Yasunaga, Pranav Agarwal, and Jure Leskovec. Planet: Predicting population response to drugs via clinical knowledge graph. *medRxiv*, 2024.

[2] Anna Gaulton et al. Chembl: a large-scale bioactivity database for drug discovery. *Nucleic Acids Research*, 40(D1):D1100–D1107, 2012.

[3] K. J. Gleason, F. Yang, and L. S. Chen. A robust two-sample transcriptome-wide Mendelian Randomization method integrating GWAS with multi-tissue eQTL summary statistics. *Genetic Epidemiology*, 45(4):353–371, Jun 2021.

[4] K. P. Jablonski, M. Pirkl, D. Ćevid, P. Bühlmann, and N. Beerenwinkel. Identifying cancer pathway dysregulations using differential causal effects. *Bioinformatics*, 38(6):1550–1559, 2022.

[5] Sunghwan Kim, Paul A. Thiessen, Evan E. Bolton, and Stephen H. Bryant. Pubchem pug-rest: a restful web service for programmatic access to pubchem data. *Nucleic Acids Research*, 46(W1):W563–W570, 2018.

[6] Sunghwan Kim, Paul A. Thiessen, Evan E. Bolton, Jie Chen, Gang Fu, Asta Gindulyte, Lianyi Han, Jane He, Siqian He, Benjamin A. Shoemaker, Jie Wang, Bo Yu, Jian Zhang, and Stephen H. Bryant. Pubchem: a public information system for analyzing bioactivities of small molecules. *Nucleic Acids Research*, 44(D1):D1202–D1213, 2016.

[7] Christopher A. Lipinski, Franco Lombardo, Beryl W. Dominy, and Paul J. Feeney. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Advanced Drug Delivery Reviews*, 23(1–3):3–25, 1997.

[8] National Cancer Institute. Common terminology criteria for adverse events (ctcae) v6.0 (meddra 28.0).

[9] David Ochoa et al. Open targets platform: supporting systematic drug–target identification and prioritisation. *Nucleic Acids Research*, 49(D1):D1302–D1310, 2021.

[10] Tuan Minh Pham, Sergei Rybakov, Hiroki Suzuki, Ka-Chun Yeung, Seungjin Hong, and Jie Sun. Optimal control nodes in disease-perturbed networks as targets for combination therapy. *Nature Communications*, 10:401, 2019.

[11] Sindy Pinero, Xiaomei Li, Lin Liu, Jiuyong Li, Sang Hong Lee, Marnie Winter, Thin Nguyen, Junpeng Zhang, and Thuc Duy Le. Integrative multi-omics framework for causal gene discovery in long covid. *medRxiv*, page 2025.02.09.25321751, 2025.

[12] Sindy Licette Pinero, Xiaomei Li, Lin Liu, Jiuyong Li, Sang Hong Lee, Marnie Winter, Thin Nguyen, Junpeng Zhang, and Thuc Duy Le. Taco: Tabpfn augmented causal outcomes for early detection of long covid. *medRxiv*, page 2025.10.02.25337138, 2025.

[13] U.S. National Library of Medicine. Dailymed: Drug label information. National Library of Medicine (NLM), n.d. Accessed programmatically via the DailyMed services API.

[14] A. Vinayagam, T. E. Gibson, H. J. Lee, B. Yilmazel, C. Roesel, Y. Hu, Y. Kwon, A. Sharma, Y. Y. Liu, N. Perrimon, and A. L. Barabási. Controllability analysis of the directed human protein interaction network identifies disease genes and drug targets. *Proceedings of the National Academy of Sciences of the United States of America*, 113(18):4976–4981, May 2016.

[15] Hua Xu, Shane P Stenner, Son Doan, Kevin B Johnson, Lemuel R Waitman, and Joshua C Denny. MedEx: a medication information extraction system for clinical narratives. *Journal of the American Medical Informatics Association*, 17(1):19–24, 2010.

[16] Chi Yuan, Patrick B Ryan, Casey Ta, Yixuan Guo, Ziran Li, Jill Hardin, Rupa Makadia, Peng Jin, Ning Shang, Tian Kang, and Chunhua Weng. Criteria2Query: a natural language interface to clinical databases for cohort definition. *Journal of the American Medical Informatics Association*, 26(4):294–305, 2019.