


Bounding the Infinite

A Resolution to Hilbert's 16th Problem

Uniform Bounds for Planar Polynomial Limit Cycles

Distilling the 2026 mathematical breakthrough, the 160-page 6-step proof, and the Lean-verified Quintic Liénard companion.




The System (Two Variables over Time)

$$\frac{dx}{dt} = P(x, y)$$


$$\frac{dy}{dt} = Q(x, y)$$

P and Q are real polynomials of degree d. At every point (x,y), the derivatives define a specific velocity vector.



What is a Limit Cycle? (Target: An isolated closed orbit)

The Non-Example: Center




A continuum of periodic orbits.

NOT ISOLATED

Moving slightly off one orbit simply lands you on another parallel orbit.

The Example: Van der Pol



An isolated periodic solution.

ISOLATED (LIMIT CYCLE)

A slight perturbation forces the trajectory to ultimately return to this specific Amber cycle.

The Poincaré Section: Reducing the Dimension



MECHANICS: Return Map $\Pi(r)$:
Measures the position after
one full circuit from
starting point r .

Displacement Function:
 $\Pi(r) - r$

Displacement = 0

Translates to: Closed Orbits

Isolated Zeros

Translates to: Limit Cycles (Our Target)

The Century-Old Mystery: Hilbert's 16th Problem

1900: Paris International Congress of Mathematicians.


What is the maximum number and relative position of limit cycles for polynomial vector fields of degree n ?

$H(n)$

Does a finite upper bound exist depending only on the degree n ?
(Note: $H(1) = 0$ as linear fields have no limit cycles. For $n > 1$, the answer has remained unknown).

Individual Finiteness


Does a single polynomial vector field have finitely many limit cycles?



- **1923:** Henri Dulac (Proof gap discovered in 1981).
- **1991 / 1992:** Yulii Ilyashenko & Jean Écalle (Independently proved: YES. Individual fields have finitely many cycles).

Uniform Boundedness

Is there a global cap $B(d)$ for every field of degree d ?



The Gap

Even if every individual field is finite, the maximum number across all fields of degree d could hypothetically be infinite.


Status: Open until 2026.

“For each integer $d \geq 1$, there is a finite nonnegative integer $B(d)$ such every real planar polynomial vector field of degree at most d has at most $B(d)$ limit cycles in the whole plane.”

RULE 1: This theorem proves a bound $B(d)$ EXISTS.

RULE 2: It gives NO VALUE for $B(d)$ or $H(n)$ —not even for $H(2)$.

Proof Step 1: Spatial Rescaling & Controlled Rotation



Rescaling

$$z = a + Ru$$

Bring all finite collections of limit cycles strictly inside one fixed bounded square.




Rotation

$$V_\mu = V + \mu(-Q, P)$$

A signed rotational perturbation guarantees the system produces at least as many hyperbolic (simple/isolated) cycles as any prescribed finite collection.


CELL DIVISION:

A limit cycle crosses each transverse boundary arc at most once. Therefore, every periodic orbit, apart from at most $T(d)$ exceptional orbits containing tangential boundary arcs, is represented by a closed word of bounded length.



THE "WORD":

A limit cycle crosses each transverse boundary arc at most once. Therefore, every periodic orbit, apart from at most $T(d)$ exceptional orbits containing tangential boundary arcs, is represented by a closed word of bounded length.



THE LINK: Each "letter" in the word represents exactly one of four scalar transfers.

EXAMPLE (SADDLE TRANSFER):


System: $x' = x, y' = -\lambda y$

Mapping: $(r, 1) \rightarrow (1, r^\lambda)$

Flight Time: $\log(1/r)$

TAKEAWAY: Intractable, dense geometry is reduced to precisely defined scalar equations.

Proof Step 4: Matching Equations & Separation



CLOSING THE LOOP:

For each valid "word", matching equations are generated to link the scalar transfers back together.

GEOMETRIC MEANING:

Exact local matching of these segments implies a closed periodic orbit.

THE SEPARATION PRINCIPLE:

Within each regular chart of these equations, representations of different hyperbolic limit cycles lie in different connected components of the fiber.

Proof Step 5: The Projection Count (The Fulcrum)



- **THE PRINCIPLE:** Counting the connected components in the matching system uniformly bounds the number of limit cycles.
- **THE CATCH:** This projection count only works if “**absolute isolated-zero finiteness**” holds for the closure of the generated systems. We must prove these isolated zeros are finite.

(The "monomial annuli" used here are pieces of a rewritten scalar equation, NOT physical rings in the real plane).

(The "monomial annuli" used here are pieces of a rewritten scalar equation NOT physical rings in the real plane)

(The "monomial annuli" used here are pieces of a rewritten scalar equation, NOT physical rings in the real plane).

Assume (for contradiction) an infinite sequence of isolated zeros exists.

Utilize exact “asymptotic packets” (finite saturated flags) evaluated on nested complex domains.

The calculus of these trees forces a separation, breaking the initial assumption. Finiteness is achieved.

The Computable Companion (Formal-Verified)

The Math

SYSTEM: Quintic Liénard

$$x' = y - F(x)$$

$$y' = -x$$

(where $\deg F \leq 5$)

THEOREM:


Yields at most 2 limit cycles.
The exact maximum is 2.

VERIFICATION:

Fully Lean-checked.

The Concrete Example

FUNCTION: $F(x) = \varepsilon \left(4x - \frac{20}{3}x^3 + \frac{8}{5}x^5 \right)$



Reality Check: What Is (and Isn't) Proved

WHAT WE HAVE

- A proof that a uniform finite bound $B(d)$ EXISTS for all degrees.
- A 160-page novel analytic proof utilizing asymptotic packets and matching equations.
- A Lean-checked proof that quintic Liénard systems have at most 2 limit cycles, and that 2 occur.

WHAT WE DON'T HAVE

- **NO VALUES:** NO exact value for $B(d)$ or $H(n)$ is provided (not even $H(2)$).
- **NO $H(5)$:** The Lean-verified '2' is not $H(5)$; it only applies to the restricted Liénard class.
- **NO FORMALIZATION:** The 160-page main proof is currently unformalized and unreviewed.
- **NO TRADITIONAL METHODS:** The main proof does not rely on Abelian integrals, Melnikov functions, computer searches, or Poincaré-Bendixson.