

## Alex Psomas: Lecture 15.

Bayes' Rule, Mutual Independence, Collisions and Collecting

# Alex Psomas: Lecture 15.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

1. Conditional Probability
2. Independence
3. Bayes' Rule
4. Balls and Bins
5. Coupons

# Conditional Probability: Review

# Conditional Probability: Review

Recall:

$$\blacktriangleright \Pr[A|B] = \frac{\Pr[A \cap B]}{\Pr[B]}.$$

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A],$

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A],$



# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A],$

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A]$ ,  
i.e., if  $Pr[A \cap B] = Pr[A]Pr[B].$

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A]$ ,  
i.e., if  $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note:  $B \subset A \Rightarrow A$  and  $B$  are

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A]$ ,  
i.e., if  $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note:  $B \subset A \Rightarrow A$  and  $B$  are positively correlated.

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A]$ ,  
i.e., if  $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note:  $B \subset A \Rightarrow A$  and  $B$  are positively correlated.  
( $Pr[A|B] = 1 > Pr[A]$ )

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A]$ ,  
i.e., if  $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note:  $B \subset A \Rightarrow A$  and  $B$  are positively correlated.  
( $Pr[A|B] = 1 > Pr[A]$ )
- ▶ Note:  $A \cap B = \emptyset \Rightarrow A$  and  $B$  are

# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A]$ ,  
i.e., if  $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note:  $B \subset A \Rightarrow A$  and  $B$  are positively correlated.  
( $Pr[A|B] = 1 > Pr[A]$ )
- ▶ Note:  $A \cap B = \emptyset \Rightarrow A$  and  $B$  are negatively correlated.



# Conditional Probability: Review

Recall:

- ▶  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}.$
- ▶ Hence,  $Pr[A \cap B] = Pr[B]Pr[A|B] = Pr[A]Pr[B|A].$
- ▶  $A$  and  $B$  are *positively correlated* if  $Pr[A|B] > Pr[A]$ ,  
i.e., if  $Pr[A \cap B] > Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *negatively correlated* if  $Pr[A|B] < Pr[A]$ ,  
i.e., if  $Pr[A \cap B] < Pr[A]Pr[B].$
- ▶  $A$  and  $B$  are *independent* if  $Pr[A|B] = Pr[A]$ ,  
i.e., if  $Pr[A \cap B] = Pr[A]Pr[B].$
- ▶ Note:  $B \subset A \Rightarrow A$  and  $B$  are positively correlated.  
( $Pr[A|B] = 1 > Pr[A]$ )
- ▶ Note:  $A \cap B = \emptyset \Rightarrow A$  and  $B$  are negatively correlated.  
( $Pr[A|B] = 0 < Pr[A]$ )

# Monty Hall

# Monty Hall

3 closed doors.

# Monty Hall

3 closed doors. Behind one of the doors there is a prize (car).

# Monty Hall

3 closed doors. Behind one of the doors there is a prize (car).  
The others have goats.

# Monty Hall

3 closed doors. Behind one of the doors there is a prize (car).  
The others have goats.

You pick a door. Say door number 1

# Monty Hall

3 closed doors. Behind one of the doors there is a prize (car).  
The others have goats.

You pick a door. Say door number 1

I open door 2 or door 3. One of the two that I **know** doesn't  
have the prize.

# Monty Hall

3 closed doors. Behind one of the doors there is a prize (car).  
The others have goats.

You pick a door. Say door number 1

I open door 2 or door 3. One of the two that I **know** doesn't  
have the prize. Say it was door 2



# Monty Hall

3 closed doors. Behind one of the doors there is a prize (car).  
The others have goats.

You pick a door. Say door number 1

I open door 2 or door 3. One of the two that I **know** doesn't  
have the prize. Say it was door 2

I ask: Would you like to change your door to number 3?

# Monty Hall

3 closed doors. Behind one of the doors there is a prize (car).  
The others have goats.

You pick a door. Say door number 1

I open door 2 or door 3. One of the two that I **know** doesn't  
have the prize. Say it was door 2

I ask: **Would you like to change your door to number 3?**

Question: What should you do in order to maximize the  
probability of winning?

# Monty Hall

# Monty Hall

Change!!!!

# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

How does that make any sense????

# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

How does that make any sense????

Say the original door where the prize is random.

# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

How does that make any sense????

Say the original door where the prize is random. So each door has probability  $\frac{1}{3}$ .



# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

How does that make any sense????

Say the original door where the prize is random. So each door has probability  $\frac{1}{3}$ .

You pick door 1.

# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

How does that make any sense????

Say the original door where the prize is random. So each door has probability  $\frac{1}{3}$ .

You pick door 1. What's the probability that it's in either 2 or 3?

# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

How does that make any sense????

Say the original door where the prize is random. So each door has probability  $\frac{1}{3}$ .

You pick door 1. What's the probability that it's in either 2 or 3?  
 $\frac{2}{3}$

# Monty Hall

Change!!!!

What is the probability that the prize is in door 3?  $\frac{2}{3}$ !

How does that make any sense????

Say the original door where the prize is random. So each door has probability  $\frac{1}{3}$ .

You pick door 1. What's the probability that it's in either 2 or 3?  
 $\frac{2}{3}$

The door I opened wasn't random! I knew it didn't have a prize!!

# Monty Hall

Change!!!!


What is the probability that the prize is in door 3?  $\frac{2}{3}$ !


How does that make any sense????

Say the original door where the prize is random. So each door has probability  $\frac{1}{3}$ .

You pick door 1. What's the probability that it's in either 2 or 3?  
 $\frac{2}{3}$

The door I opened wasn't random! I knew it didn't have a prize!!  
Therefore, switching, is like getting to pick two doors at the beginning!





## Balls in bins

I throw 5 (indistinguishable) balls in two bins.



## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ .

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart.

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .

# Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.



# Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ .

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well...

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well... I guess probability is wrong...

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well... I guess probability is wrong...

Or.....

## Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well... I guess probability is wrong...

Or..... Could one of the approaches be wrong???

# Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well... I guess probability is wrong...

Or..... Could one of the approaches be wrong???

Approach 1 is **WRONG!**

# Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes:  $(5, 0)$ ,  $(4, 1)$ ,  $(3, 2)$ ,  $(2, 3)$ ,  $(1, 4)$ ,  $(0, 5)$ . Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes:  $(1, 1, 1, 1, 1)$ ,  $(1, 1, 1, 1, 2)$ ,  $\dots$   $(2, 2, 2, 2, 2)$ .  $(x, 1, x, x, x)$  means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well... I guess probability is wrong...

Or..... Could one of the approaches be wrong???

Approach 1 is **WRONG!** Why did we divide by  $|\Omega|$ ???



# Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes: (5,0), (4,1), (3,2), (2,3), (1,4), (0,5). Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes: (1,1,1,1,1), (1,1,1,1,2), ... (2,2,2,2,2). (x,1,x,x,x) means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well... I guess probability is wrong...

Or..... Could one of the approaches be wrong???

Approach 1 is **WRONG!** Why did we divide by  $|\Omega|$ ???

Why??????? Noooooooooooooooooo

# Balls in bins

I throw 5 (indistinguishable) balls in two bins. What is the probability that the first bin is empty?

1. Approach 1: There are 6 outcomes: (5,0), (4,1), (3,2), (2,3), (1,4), (0,5). Probability that the first bin is empty is  $\frac{1}{6}$
2. Approach 2: I pretend I can tell the balls apart. There are  $2^5$  outcomes: (1,1,1,1,1), (1,1,1,1,2), ... (2,2,2,2,2). (x,1,x,x,x) means that the second ball I threw landed in the first bin.  
Probability that the first bin is empty is  $\frac{1}{2^5}$ . The fact that I can tell them apart shouldn't change the probability.

Well... I guess probability is wrong...

Or..... Could one of the approaches be wrong???




Approach 1 is **WRONG!** Why did we divide by  $|\Omega|$ ???

Why??????? Nooooooooooooooooooooooooooooo

## Conditional Probability: Pictures




# Conditional Probability: Pictures

Illustrations: Pick a point uniformly in the unit square



# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



► Left:  $A$  and  $B$  are

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- Left:  $A$  and  $B$  are independent.

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



► Left:  $A$  and  $B$  are independent.  $Pr[B] =$

# Conditional Probability: Pictures

Illustrations: Pick a point uniformly in the unit square




- Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;



# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



► Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] =$

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] =$

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] =$

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$ .

# Conditional Probability: Pictures

Illustrations: Pick a point uniformly in the unit square




- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_2, b_1)$ .



# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_2, b_1)$ .
- ▶ Right:  $A$  and  $B$  are

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_2, b_1)$ .
- ▶ Right:  $A$  and  $B$  are negatively correlated.

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square



- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_2, b_1)$ .
- ▶ Right:  $A$  and  $B$  are negatively correlated.  
 $Pr[B|A] = b_1 < Pr[B|\bar{A}] = b_2$ .

# Conditional Probability: Pictures


Illustrations: Pick a point uniformly in the unit square




- ▶ Left:  $A$  and  $B$  are independent.  $Pr[B] = b$ ;  $Pr[B|A] = b$ .
- ▶ Middle:  $A$  and  $B$  are positively correlated.  
 $Pr[B|A] = b_1 > Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_2, b_1)$ .
- ▶ Right:  $A$  and  $B$  are negatively correlated.  
 $Pr[B|A] = b_1 < Pr[B|\bar{A}] = b_2$ . Note:  $Pr[B] \in (b_1, b_2)$ .

# Bayes and Biased Coin

# Bayes and Biased Coin




# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] =$$




# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5;$$


# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] =$$


# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] =$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5;$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] =$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6;$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] =$$



# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] =$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6$$

# Bayes and Biased Coin




Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then


$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6}$$

# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then


$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6} = \frac{Pr[A]Pr[B|A]}{Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]}$$

# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then


$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$

$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6} = \frac{Pr[A]Pr[B|A]}{Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]} \\ \approx 0.46$$

# Bayes and Biased Coin



Pick a point uniformly at random in the unit square. Then

$$Pr[A] = 0.5; Pr[\bar{A}] = 0.5$$

$$Pr[B|A] = 0.5; Pr[B|\bar{A}] = 0.6; Pr[A \cap B] = 0.5 \times 0.5$$

$$Pr[B] = 0.5 \times 0.5 + 0.5 \times 0.6 = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]$$


$$Pr[A|B] = \frac{0.5 \times 0.5}{0.5 \times 0.5 + 0.5 \times 0.6} = \frac{Pr[A]Pr[B|A]}{Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}]}$$

$\approx 0.46 = \text{fraction of } B \text{ that is inside } A$




# Bayes: General Case

# Bayes: General Case




# Bayes: General Case



Pick a point uniformly at random in the unit square. Then


# Bayes: General Case



Pick a point uniformly at random in the unit square. Then

$$Pr[A_m] = p_m, m = 1, \dots, M$$

# Bayes: General Case




Pick a point uniformly at random in the unit square. Then

$$Pr[A_m] = p_m, m = 1, \dots, M$$

$$Pr[B|A_m] = q_m, m = 1, \dots, M;$$

# Bayes: General Case




Pick a point uniformly at random in the unit square. Then

$$Pr[A_m] = p_m, m = 1, \dots, M$$

$$Pr[B|A_m] = q_m, m = 1, \dots, M; Pr[A_m \cap B] =$$

# Bayes: General Case




Pick a point uniformly at random in the unit square. Then

$$Pr[A_m] = p_m, m = 1, \dots, M$$

$$Pr[B|A_m] = q_m, m = 1, \dots, M; Pr[A_m \cap B] = p_m q_m$$

# Bayes: General Case



Pick a point uniformly at random in the unit square. Then


$$Pr[A_m] = p_m, m = 1, \dots, M$$

$$Pr[B|A_m] = q_m, m = 1, \dots, M; Pr[A_m \cap B] = p_m q_m$$

$$Pr[B] = p_1 q_1 + \dots + p_M q_M$$



# Bayes: General Case



Pick a point uniformly at random in the unit square. Then


$$Pr[A_m] = p_m, m = 1, \dots, M$$

$$Pr[B|A_m] = q_m, m = 1, \dots, M; Pr[A_m \cap B] = p_m q_m$$

$$Pr[B] = p_1 q_1 + \dots p_M q_M$$

$$Pr[A_m|B] = \frac{p_m q_m}{p_1 q_1 + \dots p_M q_M}$$

# Bayes: General Case



Pick a point uniformly at random in the unit square. Then


$$Pr[A_m] = p_m, m = 1, \dots, M$$

$$Pr[B|A_m] = q_m, m = 1, \dots, M; Pr[A_m \cap B] = p_m q_m$$


$$Pr[B] = p_1 q_1 + \dots + p_M q_M$$

$$Pr[A_m|B] = \frac{p_m q_m}{p_1 q_1 + \dots + p_M q_M} = \text{fraction of } B \text{ inside } A_m.$$

# Why do you have a fever?




# Why do you have a fever?



Using Bayes' rule, we find


# Why do you have a fever?



Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

# Why do you have a fever?




Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

# Why do you have a fever?




Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$

# Why do you have a fever?



Using Bayes' rule, we find

$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$

The values  $0.58, 5 \times 10^{-8}, 0.42$  are the **posterior probabilities**.





Why do you have a fever?

# Why do you have a fever?

Our “Bayes’ Square” picture:

# Why do you have a fever?



Our “Bayes’ Square” picture:



58% of Fever = Flu  
 $\approx 0\%$  of Fever = Ebola  
42% of Fever = Other

# Why do you have a fever?

Our “Bayes’ Square” picture:





58% of Fever = Flu  
 $\approx 0\%$  of Fever = Ebola  
42% of Fever = Other

Note that even though  $Pr[\text{Fever}|\text{Ebola}] = 1$ ,

# Why do you have a fever?

Our “Bayes’ Square” picture:



58% of Fever = Flu

$\approx 0\%$  of Fever = Ebola


42% of Fever = Other

Note that even though  $Pr[\text{Fever}|\text{Ebola}] = 1$ , one has

$$Pr[\text{Ebola}|\text{Fever}] \approx 0.$$

# Why do you have a fever?

Our “Bayes’ Square” picture:



58% of Fever = Flu  
 $\approx 0\%$  of Fever = Ebola  
42% of Fever = Other


Note that even though  $Pr[\text{Fever}|\text{Ebola}] = 1$ , one has

$$Pr[\text{Ebola}|\text{Fever}] \approx 0.$$

This example shows the importance of the prior probabilities.


# Bayes' Rule Operations

# Bayes' Rule Operations





# Bayes' Rule Operations



Bayes' Rule is the canonical example of how information changes our opinions.

# Independence

Recall :

$A$  and  $B$  are independent

# Independence

Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

# Independence

Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow \Pr[A \cap B] = \Pr[A] \Pr[B]$$

$$\Leftrightarrow \Pr[A|B] = \Pr[A].$$

# Independence


Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



# Independence


Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



$(A_2, B)$  are independent:

# Independence


Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



$(A_2, B)$  are independent:  $Pr[A_2|B] = 0.5 = Pr[A_2]$ .

# Independence


Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



$(A_2, B)$  are independent:  $Pr[A_2|B] = 0.5 = Pr[A_2]$ .

$(A_2, \bar{B})$  are independent:



# Independence


Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



$(A_2, B)$  are independent:  $Pr[A_2|B] = 0.5 = Pr[A_2]$ .

$(A_2, \bar{B})$  are independent:  $Pr[A_2|\bar{B}] = 0.5 = Pr[A_2]$ .

# Independence


Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



$(A_2, B)$  are independent:  $Pr[A_2|B] = 0.5 = Pr[A_2]$ .

$(A_2, \bar{B})$  are independent:  $Pr[A_2|\bar{B}] = 0.5 = Pr[A_2]$ .

$(A_1, B)$  are not independent:

# Independence


Recall :

$A$  and  $B$  are independent

$$\Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B]$$

$$\Leftrightarrow Pr[A|B] = Pr[A].$$

Consider the example below:



$(A_2, B)$  are independent:  $Pr[A_2|B] = 0.5 = Pr[A_2]$ .

$(A_2, \bar{B})$  are independent:  $Pr[A_2|\bar{B}] = 0.5 = Pr[A_2]$ .

$(A_1, B)$  are not independent:  $Pr[A_1|B] = \frac{0.1}{0.5} = 0.2 \neq Pr[A_1] = 0.25$ .

# Pairwise Independence


Flip two fair coins. Let

- ▶  $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶  $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶  $C = \text{'the two coins are different'} = \{TH, HT\}.$

# Pairwise Independence

Flip two fair coins. Let


- ▶  $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶  $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶  $C = \text{'the two coins are different'} = \{TH, HT\}.$



# Pairwise Independence

Flip two fair coins. Let

- ▶  $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶  $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶  $C = \text{'the two coins are different'} = \{TH, HT\}.$




$A, C$  are independent;

# Pairwise Independence

Flip two fair coins. Let

- ▶  $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶  $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶  $C = \text{'the two coins are different'} = \{TH, HT\}.$




$A, C$  are independent;  $B, C$  are independent;

# Pairwise Independence

Flip two fair coins. Let

- ▶  $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶  $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶  $C = \text{'the two coins are different'} = \{TH, HT\}.$




$A, C$  are independent;  $B, C$  are independent;  
 $A \cap B, C$  are **not** independent.



# Pairwise Independence

Flip two fair coins. Let

- ▶  $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶  $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶  $C = \text{'the two coins are different'} = \{TH, HT\}.$




$A, C$  are independent;  $B, C$  are independent;

$A \cap B, C$  are **not** independent. ( $Pr[A \cap B \cap C] = 0 \neq Pr[A \cap B]Pr[C].$ )

# Pairwise Independence

Flip two fair coins. Let

- ▶  $A = \text{'first coin is H'} = \{HT, HH\};$
- ▶  $B = \text{'second coin is H'} = \{TH, HH\};$
- ▶  $C = \text{'the two coins are different'} = \{TH, HT\}.$



$A, C$  are independent;  $B, C$  are independent;

$A \cap B, C$  are **not** independent. ( $Pr[A \cap B \cap C] = 0 \neq Pr[A \cap B]Pr[C].$ )

$A$  did not say anything about  $C$  and  $B$  did not say anything about  $C$ , but  $A \cap B$  said something about  $C$ !

## Example 2

Flip a fair coin 5 times.

## Example 2

Flip a fair coin 5 times. Let  $A_n = \text{'coin } n \text{ is H'}$ , for  $n = 1, \dots, 5$ .

## Example 2

Flip a fair coin 5 times. Let  $A_n$  = 'coin  $n$  is H', for  $n = 1, \dots, 5$ .

Then,

$A_m, A_n$  are independent for all  $m \neq n$ .

## Example 2

Flip a fair coin 5 times. Let  $A_n = \text{'coin } n \text{ is H'}$ , for  $n = 1, \dots, 5$ .

Then,

$A_m, A_n$  are independent for all  $m \neq n$ .

Also,

$A_1$  and  $A_3 \cap A_5$  are independent.

## Example 2

Flip a fair coin 5 times. Let  $A_n$  = 'coin  $n$  is H', for  $n = 1, \dots, 5$ .

Then,

$A_m, A_n$  are independent for all  $m \neq n$ .

Also,

$A_1$  and  $A_3 \cap A_5$  are independent.

Indeed,

$$Pr[A_1 \cap (A_3 \cap A_5)] = \frac{1}{8} = Pr[A_1]Pr[A_3 \cap A_5]$$

.

## Example 2

Flip a fair coin 5 times. Let  $A_n$  = 'coin  $n$  is H', for  $n = 1, \dots, 5$ .

Then,

$A_m, A_n$  are independent for all  $m \neq n$ .

Also,

$A_1$  and  $A_3 \cap A_5$  are independent.

Indeed,

$$Pr[A_1 \cap (A_3 \cap A_5)] = \frac{1}{8} = Pr[A_1]Pr[A_3 \cap A_5]$$

. Similarly,

$A_1 \cap A_2$  and  $A_3 \cap A_4 \cap A_5$  are independent.



## Example 2

Flip a fair coin 5 times. Let  $A_n$  = 'coin  $n$  is H', for  $n = 1, \dots, 5$ .

Then,

$A_m, A_n$  are independent for all  $m \neq n$ .

Also,

$A_1$  and  $A_3 \cap A_5$  are independent.

Indeed,

$$Pr[A_1 \cap (A_3 \cap A_5)] = \frac{1}{8} = Pr[A_1]Pr[A_3 \cap A_5]$$

. Similarly,

$A_1 \cap A_2$  and  $A_3 \cap A_4 \cap A_5$  are independent.

This leads to a definition ....

# Mutual Independence

**Definition** Mutual Independence

# Mutual Independence

**Definition** Mutual Independence

(a) The events  $A_1, \dots, A_5$  are mutually independent if

# Mutual Independence

## **Definition** Mutual Independence

(a) The events  $A_1, \dots, A_5$  are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

# Mutual Independence

## **Definition** Mutual Independence

(a) The events  $A_1, \dots, A_5$  are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

(b) More generally, the events  $\{A_j, j \in J\}$  are **mutually independent** if

# Mutual Independence

## **Definition** Mutual Independence

(a) The events  $A_1, \dots, A_5$  are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

(b) More generally, the events  $\{A_j, j \in J\}$  are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all finite } K \subseteq J.$$

# Mutual Independence

## Definition Mutual Independence

(a) The events  $A_1, \dots, A_5$  are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all } K \subseteq \{1, \dots, 5\}.$$

(b) More generally, the events  $\{A_j, j \in J\}$  are **mutually independent** if

$$Pr[\cap_{k \in K} A_k] = \prod_{k \in K} Pr[A_k], \text{ for all finite } K \subseteq J.$$

Example: Flip a fair coin forever. Let  $A_n =$  'coin  $n$  is H.' Then the events  $A_n$  are mutually independent.

# Mutual Independence

## Theorem



# Mutual Independence

## Theorem

(a) If the events  $\{A_j, j \in J\}$  are mutually independent and if  $K_1$  and  $K_2$  are disjoint finite subsets of  $J$ , then

# Mutual Independence

## Theorem

(a) If the events  $\{A_j, j \in J\}$  are mutually independent and if  $K_1$  and  $K_2$  are disjoint finite subsets of  $J$ , then

$\cap_{k \in K_1} A_k$  and  $\cap_{k \in K_2} A_k$  are independent.

# Mutual Independence

## Theorem

(a) If the events  $\{A_j, j \in J\}$  are mutually independent and if  $K_1$  and  $K_2$  are disjoint finite subsets of  $J$ , then

$\cap_{k \in K_1} A_k$  and  $\cap_{k \in K_2} A_k$  are independent.

(b) More generally, if the  $K_n$  are pairwise disjoint finite subsets of  $J$ , then the events

$\cap_{k \in K_n} A_k$  are mutually independent.

# Mutual Independence

## Theorem

(a) If the events  $\{A_j, j \in J\}$  are mutually independent and if  $K_1$  and  $K_2$  are disjoint finite subsets of  $J$ , then

$\cap_{k \in K_1} A_k$  and  $\cap_{k \in K_2} A_k$  are independent.

(b) More generally, if the  $K_n$  are pairwise disjoint finite subsets of  $J$ , then the events

$\cap_{k \in K_n} A_k$  are mutually independent.

(c) Also, the same is true if we replace some of the  $A_k$  by  $\bar{A}_k$ .

# Balls in bins

# Balls in bins

One throws  $m$  balls into  $n > m$  bins.


# Balls in bins

One throws  $m$  balls into  $n > m$  bins.



# Balls in bins


One throws  $m$  balls into  $n > m$  bins.





# Balls in bins

One throws  $m$  balls into  $n > m$  bins.



**Theorem:**

$Pr[\text{no collision}] \approx \exp\left\{-\frac{m^2}{2n}\right\}$ , for large enough  $n$ .

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$\Pr[A_1] =$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$\Pr[A_1] = 1$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] =$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$\Pr[A_1] = 1$$

$$\Pr[A_2|A_1] = 1 - \frac{1}{n}$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] =$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] = 1 - \frac{2}{n}$$



## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] = 1 - \frac{2}{n}$$

$$Pr[A_i|A_{i-1} \cap \cdots \cap A_1] =$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] = 1 - \frac{2}{n}$$

$$Pr[A_i|A_{i-1} \cap \cdots \cap A_1] = (1 - \frac{i-1}{n}).$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] = 1 - \frac{2}{n}$$

$$Pr[A_i|A_{i-1} \cap \cdots \cap A_1] = (1 - \frac{i-1}{n}).$$

$$\text{no collision} = A_1 \cap \cdots \cap A_m.$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] = 1 - \frac{2}{n}$$

$$Pr[A_i|A_{i-1} \cap \cdots \cap A_1] = (1 - \frac{i-1}{n}).$$

$$\text{no collision} = A_1 \cap \cdots \cap A_m.$$

Product rule:

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] = 1 - \frac{2}{n}$$

$$Pr[A_i|A_{i-1} \cap \dots \cap A_1] = (1 - \frac{i-1}{n}).$$

$$\text{no collision} = A_1 \cap \dots \cap A_m.$$

Product rule:

$$Pr[A_1 \cap \dots \cap A_m] = Pr[A_1]Pr[A_2|A_1] \dots Pr[A_m|A_1 \cap \dots \cap A_{m-1}]$$

## The Calculation.

$A_i$  = no collision when  $i$ th ball is placed in a bin.

$$Pr[A_1] = 1$$

$$Pr[A_2|A_1] = 1 - \frac{1}{n}$$

$$Pr[A_3|A_1, A_2] = 1 - \frac{2}{n}$$

$$Pr[A_i|A_{i-1} \cap \dots \cap A_1] = \left(1 - \frac{i-1}{n}\right).$$

$$\text{no collision} = A_1 \cap \dots \cap A_m.$$

Product rule:

$$Pr[A_1 \cap \dots \cap A_m] = Pr[A_1]Pr[A_2|A_1] \dots Pr[A_m|A_1 \cap \dots \cap A_{m-1}]$$

$$\Rightarrow Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \dots \left(1 - \frac{m-1}{n}\right).$$

$$\Rightarrow \Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \cdots \left(1 - \frac{m-1}{n}\right).$$

$$\Rightarrow Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \cdots \left(1 - \frac{m-1}{n}\right).$$

Hence,

$$\ln(Pr[\text{no collision}]) = \sum_{k=1}^{m-1} \ln\left(1 - \frac{k}{n}\right)$$



$$\Rightarrow \Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \cdots \left(1 - \frac{m-1}{n}\right).$$

Hence,

$$\ln(\Pr[\text{no collision}]) = \sum_{k=1}^{m-1} \ln\left(1 - \frac{k}{n}\right) \approx \sum_{k=1}^{m-1} \left(-\frac{k}{n}\right) (*)$$

$$\Rightarrow \Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \cdots \left(1 - \frac{m-1}{n}\right).$$

Hence,

$$\begin{aligned} \ln(\Pr[\text{no collision}]) &= \sum_{k=1}^{m-1} \ln\left(1 - \frac{k}{n}\right) \approx \sum_{k=1}^{m-1} \left(-\frac{k}{n}\right) \quad (*) \\ &= -\frac{1}{n} \frac{m(m-1)}{2} \quad (\dagger) \approx \end{aligned}$$

$$\Rightarrow \Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \cdots \left(1 - \frac{m-1}{n}\right).$$

Hence,

$$\begin{aligned} \ln(\Pr[\text{no collision}]) &= \sum_{k=1}^{m-1} \ln\left(1 - \frac{k}{n}\right) \approx \sum_{k=1}^{m-1} \left(-\frac{k}{n}\right) \quad (*) \\ &= -\frac{1}{n} \frac{m(m-1)}{2} \quad (\dagger) \approx -\frac{m^2}{2n} \end{aligned}$$

$$\Rightarrow \Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \cdots \left(1 - \frac{m-1}{n}\right).$$

Hence,

$$\begin{aligned} \ln(\Pr[\text{no collision}]) &= \sum_{k=1}^{m-1} \ln\left(1 - \frac{k}{n}\right) \approx \sum_{k=1}^{m-1} \left(-\frac{k}{n}\right) (*) \\ &= -\frac{1}{n} \frac{m(m-1)}{2} (\dagger) \approx -\frac{m^2}{2n} \end{aligned}$$

(\*) We used  $\ln(1 - \varepsilon) \approx -\varepsilon$  for  $|\varepsilon| \ll 1$ .

$$\Rightarrow \Pr[\text{no collision}] = \left(1 - \frac{1}{n}\right) \cdots \left(1 - \frac{m-1}{n}\right).$$


Hence,

$$\begin{aligned} \ln(\Pr[\text{no collision}]) &= \sum_{k=1}^{m-1} \ln\left(1 - \frac{k}{n}\right) \approx \sum_{k=1}^{m-1} \left(-\frac{k}{n}\right) (*) \\ &= -\frac{1}{n} \frac{m(m-1)}{2} (\dagger) \approx -\frac{m^2}{2n} \end{aligned}$$

(\*) We used  $\ln(1 - \varepsilon) \approx -\varepsilon$  for  $|\varepsilon| \ll 1$ .

(†)  $1 + 2 + \cdots + m - 1 = (m - 1)m/2$ .

# Approximation



$$\exp\{-x\} = 1 - x + \frac{1}{2!}x^2 + \cdots \approx 1 - x, \text{ for } |x| \ll 1.$$

Hence,  $-x \approx \ln(1-x)$  for  $|x| \ll 1$ .

# Balls in bins


**Theorem:**

$Pr[\text{no collision}] \approx \exp\{-\frac{m^2}{2n}\}$ , for large enough  $n$ .

# Balls in bins

## Theorem:

$Pr[\text{no collision}] \approx \exp\{-\frac{m^2}{2n}\}$ , for large enough  $n$ .





# Balls in bins

**Theorem:**

$Pr[\text{no collision}] \approx \exp\{-\frac{m^2}{2n}\}$ , for large enough  $n$ .

# Balls in bins

**Theorem:**

$Pr[\text{no collision}] \approx \exp\{-\frac{m^2}{2n}\}$ , for large enough  $n$ .

In particular,  $Pr[\text{no collision}] \approx 1/2$  for  $m^2/(2n) \approx \ln(2)$ , i.e.,

$$m \approx \sqrt{2\ln(2)n} \approx 1.2\sqrt{n}.$$

# Balls in bins

**Theorem:**

$Pr[\text{no collision}] \approx \exp\{-\frac{m^2}{2n}\}$ , for large enough  $n$ .

In particular,  $Pr[\text{no collision}] \approx 1/2$  for  $m^2/(2n) \approx \ln(2)$ , i.e.,

$$m \approx \sqrt{2\ln(2)n} \approx 1.2\sqrt{n}.$$

E.g.,  $1.2\sqrt{20} \approx 5.4$ .

# Balls in bins

## Theorem:

$Pr[\text{no collision}] \approx \exp\{-\frac{m^2}{2n}\}$ , for large enough  $n$ .

In particular,  $Pr[\text{no collision}] \approx 1/2$  for  $m^2/(2n) \approx \ln(2)$ , i.e.,

$$m \approx \sqrt{2\ln(2)n} \approx 1.2\sqrt{n}.$$

E.g.,  $1.2\sqrt{20} \approx 5.4$ .

Roughly,  $Pr[\text{collision}] \approx 1/2$  for  $m = \sqrt{n}$ .

# Balls in bins

## Theorem:

$Pr[\text{no collision}] \approx \exp\{-\frac{m^2}{2n}\}$ , for large enough  $n$ .

In particular,  $Pr[\text{no collision}] \approx 1/2$  for  $m^2/(2n) \approx \ln(2)$ , i.e.,

$$m \approx \sqrt{2\ln(2)n} \approx 1.2\sqrt{n}.$$

E.g.,  $1.2\sqrt{20} \approx 5.4$ .

Roughly,  $Pr[\text{collision}] \approx 1/2$  for  $m = \sqrt{n}$ . ( $e^{-0.5} \approx 0.6$ .)

# The birthday paradox

Today's your birthday, it's my birthday too..

Probability that  $m$  people all have different birthdays?

Today's your birthday, it's my birthday too..

Probability that  $m$  people all have different birthdays?

With  $n = 365$ , one finds



Today's your birthday, it's my birthday too..

Probability that  $m$  people all have different birthdays?

With  $n = 365$ , one finds

$Pr[\text{collision}] \approx 1/2$  if  $m \approx 1.2\sqrt{365} \approx 23$ .

Today's your birthday, it's my birthday too..

Probability that  $m$  people all have different birthdays?

With  $n = 365$ , one finds

$Pr[\text{collision}] \approx 1/2$  if  $m \approx 1.2\sqrt{365} \approx 23$ .

If  $m = 60$ , we find that

Today's your birthday, it's my birthday too..

Probability that  $m$  people all have different birthdays?

With  $n = 365$ , one finds

$Pr[\text{collision}] \approx 1/2$  if  $m \approx 1.2\sqrt{365} \approx 23$ .

If  $m = 60$ , we find that

$$Pr[\text{no collision}] \approx \exp\left\{-\frac{m^2}{2n}\right\} = \exp\left\{-\frac{60^2}{2 \times 365}\right\} \approx 0.007.$$

Today's your birthday, it's my birthday too..

Probability that  $m$  people all have different birthdays?

With  $n = 365$ , one finds

$Pr[\text{collision}] \approx 1/2$  if  $m \approx 1.2\sqrt{365} \approx 23$ .

If  $m = 60$ , we find that

$$Pr[\text{no collision}] \approx \exp\left\{-\frac{m^2}{2n}\right\} = \exp\left\{-\frac{60^2}{2 \times 365}\right\} \approx 0.007.$$

If  $m = 366$ , then  $Pr[\text{no collision}] =$

Today's your birthday, it's my birthday too..

Probability that  $m$  people all have different birthdays?

With  $n = 365$ , one finds

$Pr[\text{collision}] \approx 1/2$  if  $m \approx 1.2\sqrt{365} \approx 23$ .

If  $m = 60$ , we find that

$$Pr[\text{no collision}] \approx \exp\left\{-\frac{m^2}{2n}\right\} = \exp\left\{-\frac{60^2}{2 \times 365}\right\} \approx 0.007.$$

If  $m = 366$ , then  $Pr[\text{no collision}] = 0$ . (No approximation here!)

# The birthday paradox

$n$	$p(n)$
1	0.0%
5	2.7%
10	11.7%
20	41.1%
23	50.7%
30	70.6%
40	89.1%
50	97.0%
60	99.4%
70	99.9%
100	99.99997%
200	99.999999999999999999999999999998%
300	$(100 - (6 \times 10^{-80}))\%$
350	$(100 - (3 \times 10^{-129}))\%$
365	$(100 - (1.45 \times 10^{-155}))\%$
366	100%
367	100%

# Checksums!

# Checksums!

Consider a set of  $m$  files.



# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\}$

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

Hence,

$$Pr[\text{no collision}] \approx 1 - 10^{-3}$$



# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

Hence,

$$Pr[\text{no collision}] \approx 1 - 10^{-3} \Leftrightarrow m^2/(2n) \approx 10^{-3}$$

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

Hence,

$$\begin{aligned} Pr[\text{no collision}] \approx 1 - 10^{-3} &\Leftrightarrow m^2/(2n) \approx 10^{-3} \\ &\Leftrightarrow 2n \approx m^2 10^3 \end{aligned}$$

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

Hence,

$$\begin{aligned} Pr[\text{no collision}] \approx 1 - 10^{-3} &\Leftrightarrow m^2/(2n) \approx 10^{-3} \\ &\Leftrightarrow 2n \approx m^2 10^3 \Leftrightarrow 2^{b+1} \approx m^2 2^{10} \end{aligned}$$

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $\Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $\Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

Hence,

$$\begin{aligned} \Pr[\text{no collision}] \approx 1 - 10^{-3} &\Leftrightarrow m^2/(2n) \approx 10^{-3} \\ &\Leftrightarrow 2n \approx m^2 10^3 \Leftrightarrow 2^{b+1} \approx m^2 2^{10} \\ &\Leftrightarrow b+1 \approx 10 + 2 \log_2(m) \end{aligned}$$

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $\Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9 \ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $\Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

Hence,

$$\begin{aligned} \Pr[\text{no collision}] \approx 1 - 10^{-3} &\Leftrightarrow m^2/(2n) \approx 10^{-3} \\ &\Leftrightarrow 2n \approx m^2 10^3 \Leftrightarrow 2^{b+1} \approx m^2 2^{10} \\ &\Leftrightarrow b+1 \approx 10 + 2 \log_2(m) \approx 10 + 2.9 \ln(m). \end{aligned}$$

# Checksums!

Consider a set of  $m$  files.

Each file has a checksum of  $b$  bits.

How large should  $b$  be for  $Pr[\text{share a checksum}] \leq 10^{-3}$ ?

**Claim:**  $b \geq 2.9\ln(m) + 9$ .

**Proof:**

Let  $n = 2^b$  be the number of checksums.

We know  $Pr[\text{no collision}] \approx \exp\{-m^2/(2n)\} \approx 1 - m^2/(2n)$ .

Hence,

$$\begin{aligned} Pr[\text{no collision}] \approx 1 - 10^{-3} &\Leftrightarrow m^2/(2n) \approx 10^{-3} \\ &\Leftrightarrow 2n \approx m^2 10^3 \Leftrightarrow 2^{b+1} \approx m^2 2^{10} \\ &\Leftrightarrow b+1 \approx 10 + 2\log_2(m) \approx 10 + 2.9\ln(m). \end{aligned}$$

Note:  $\log_2(x) = \log_2(e)\ln(x) \approx 1.44\ln(x)$ .

# Coupon Collector Problem.

There are  $n$  different baseball cards.

(Brian Wilson, Jackie Robinson, Roger Hornsby, ...)

## Coupon Collector Problem.

There are  $n$  different baseball cards.

(Brian Wilson, Jackie Robinson, Roger Hornsby, ...)

One random baseball card in each cereal box.



# Coupon Collector Problem.

There are  $n$  different baseball cards.

(Brian Wilson, Jackie Robinson, Roger Hornsby, ...)

One random baseball card in each cereal box.



# Coupon Collector Problem.

There are  $n$  different baseball cards.

(Brian Wilson, Jackie Robinson, Roger Hornsby, ...)

One random baseball card in each cereal box.



**Theorem:**

# Coupon Collector Problem.

There are  $n$  different baseball cards.

(Brian Wilson, Jackie Robinson, Roger Hornsby, ...)

One random baseball card in each cereal box.



**Theorem:** If you buy  $m$  boxes,

# Coupon Collector Problem.

There are  $n$  different baseball cards.

(Brian Wilson, Jackie Robinson, Roger Hornsby, ...)

One random baseball card in each cereal box.



**Theorem:** If you buy  $m$  boxes,

(a)  $Pr[\text{miss one specific item}] \approx e^{-\frac{m}{n}}$

# Coupon Collector Problem.

There are  $n$  different baseball cards.

(Brian Wilson, Jackie Robinson, Roger Hornsby, ...)

One random baseball card in each cereal box.



**Theorem:** If you buy  $m$  boxes,

(a)  $Pr[\text{miss one specific item}] \approx e^{-\frac{m}{n}}$

(b)  $Pr[\text{miss any one of the items}] \leq ne^{-\frac{m}{n}}.$

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$



## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ...

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ... for  $m$  times. Hence,

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ... for  $m$  times. Hence,

$$Pr[A_m] = (1 - \frac{1}{n}) \times \cdots \times (1 - \frac{1}{n})$$

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ... for  $m$  times. Hence,

$$\begin{aligned}Pr[A_m] &= (1 - \frac{1}{n}) \times \cdots \times (1 - \frac{1}{n}) \\&= (1 - \frac{1}{n})^m\end{aligned}$$

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ... for  $m$  times. Hence,

$$\begin{aligned}Pr[A_m] &= (1 - \frac{1}{n}) \times \cdots \times (1 - \frac{1}{n}) \\&= (1 - \frac{1}{n})^m\end{aligned}$$

$$\ln(Pr[A_m]) = m \ln(1 - \frac{1}{n}) \approx$$

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ... for  $m$  times. Hence,

$$\begin{aligned}Pr[A_m] &= (1 - \frac{1}{n}) \times \cdots \times (1 - \frac{1}{n}) \\&= (1 - \frac{1}{n})^m\end{aligned}$$

$$\ln(Pr[A_m]) = m \ln(1 - \frac{1}{n}) \approx m \times (-\frac{1}{n})$$

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ... for  $m$  times. Hence,

$$\begin{aligned}Pr[A_m] &= (1 - \frac{1}{n}) \times \cdots \times (1 - \frac{1}{n}) \\&= (1 - \frac{1}{n})^m\end{aligned}$$

$$\ln(Pr[A_m]) = m \ln(1 - \frac{1}{n}) \approx m \times (-\frac{1}{n})$$

$$Pr[A_m] \approx \exp\{-\frac{m}{n}\}.$$

## Coupon Collector Problem: Analysis.

Event  $A_m$  = 'fail to get Brian Wilson in  $m$  cereal boxes'

Fail the first time:  $(1 - \frac{1}{n})$

Fail the second time:  $(1 - \frac{1}{n})$

And so on ... for  $m$  times. Hence,

$$\begin{aligned}Pr[A_m] &= (1 - \frac{1}{n}) \times \cdots \times (1 - \frac{1}{n}) \\&= (1 - \frac{1}{n})^m\end{aligned}$$

$$\ln(Pr[A_m]) = m \ln(1 - \frac{1}{n}) \approx m \times (-\frac{1}{n})$$

$$Pr[A_m] \approx \exp\{-\frac{m}{n}\}.$$

For  $p_m = \frac{1}{2}$ , we need around  $n \ln 2 \approx 0.69n$  boxes.



## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

Events:  $E_k =$  'fail to get player  $k$ ' , for  $k = 1, \dots, n$

## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

Events:  $E_k =$  'fail to get player  $k$ ' , for  $k = 1, \dots, n$

## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

Events:  $E_k =$  'fail to get player  $k$ ' , for  $k = 1, \dots, n$

Probability of failing to get at least one of these  $n$  players:

$$p := Pr[E_1 \cup E_2 \cdots \cup E_n]$$

## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

Events:  $E_k =$  'fail to get player  $k$ ' , for  $k = 1, \dots, n$

Probability of failing to get at least one of these  $n$  players:

$$p := Pr[E_1 \cup E_2 \cdots \cup E_n]$$

How does one estimate  $p$ ?

## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

Events:  $E_k = \text{'fail to get player k'}$ , for  $k = 1, \dots, n$

Probability of failing to get at least one of these  $n$  players:

$$p := \Pr[E_1 \cup E_2 \cdots \cup E_n]$$

How does one estimate  $p$ ? **Union Bound:**

$$p = \Pr[E_1 \cup E_2 \cdots \cup E_n] \leq \Pr[E_1] + \Pr[E_2] \cdots \Pr[E_n].$$

## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

Events:  $E_k = \text{'fail to get player } k\text{'}$ , for  $k = 1, \dots, n$

Probability of failing to get at least one of these  $n$  players:

$$p := \Pr[E_1 \cup E_2 \cdots \cup E_n]$$

How does one estimate  $p$ ? **Union Bound:**

$$p = \Pr[E_1 \cup E_2 \cdots \cup E_n] \leq \Pr[E_1] + \Pr[E_2] \cdots \Pr[E_n].$$

$$\Pr[E_k] \approx e^{-\frac{m}{n}}, k = 1, \dots, n.$$

## Collect all cards?

Experiment: Choose  $m$  cards at random with replacement.

Events:  $E_k = \text{'fail to get player k'}$ , for  $k = 1, \dots, n$

Probability of failing to get at least one of these  $n$  players:

$$p := \Pr[E_1 \cup E_2 \cdots \cup E_n]$$

How does one estimate  $p$ ? **Union Bound:**

$$p = \Pr[E_1 \cup E_2 \cdots \cup E_n] \leq \Pr[E_1] + \Pr[E_2] \cdots \Pr[E_n].$$

$$\Pr[E_k] \approx e^{-\frac{m}{n}}, k = 1, \dots, n.$$

Plug in and get

$$p \leq ne^{-\frac{m}{n}}.$$



## Collect all cards?

Thus,

$$Pr[\text{missing at least one card}] \leq ne^{-\frac{m}{n}}.$$

## Collect all cards?

Thus,

$$Pr[\text{missing at least one card}] \leq ne^{-\frac{m}{n}}.$$

Hence,

$$Pr[\text{missing at least one card}] \leq p \text{ when } m \geq n \ln\left(\frac{n}{p}\right).$$

## Collect all cards?

Thus,

$$\Pr[\text{missing at least one card}] \leq ne^{-\frac{m}{n}}.$$

Hence,

$$\Pr[\text{missing at least one card}] \leq p \text{ when } m \geq n \ln\left(\frac{n}{p}\right).$$

To get  $p = 1/2$ , set  $m = n \ln(2n)$ .

## Collect all cards?

Thus,

$$\Pr[\text{missing at least one card}] \leq ne^{-\frac{m}{n}}.$$

Hence,

$$\Pr[\text{missing at least one card}] \leq p \text{ when } m \geq n \ln\left(\frac{n}{p}\right).$$

To get  $p = 1/2$ , set  $m = n \ln(2n)$ .

E.g.,  $n = 10^2 \Rightarrow m = 530$ ;

## Collect all cards?

Thus,

$$\Pr[\text{missing at least one card}] \leq ne^{-\frac{m}{n}}.$$

Hence,

$$\Pr[\text{missing at least one card}] \leq p \text{ when } m \geq n \ln\left(\frac{n}{p}\right).$$

To get  $p = 1/2$ , set  $m = n \ln(2n)$ .

E.g.,  $n = 10^2 \Rightarrow m = 530$ ;  $n = 10^3 \Rightarrow m = 7600$ .

# Summary.

Bayes' Rule, Mutual Independence, Collisions and Collecting

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- Bayes' Rule:  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$ .

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \cdots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \cdots \cap A_{n-1}]$ .



# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \cdots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \cdots \cap A_{n-1}]$ .
- ▶ **Balls in bins:**  $m$  balls into  $n > m$  bins.

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \cdots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \cdots \cap A_{n-1}]$ .
- ▶ **Balls in bins:**  $m$  balls into  $n > m$  bins.

$$Pr[\text{no collisions}] \approx \exp\left\{-\frac{m^2}{2n}\right\}$$

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \dots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \dots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \dots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]$ .
- ▶ **Balls in bins:**  $m$  balls into  $n > m$  bins.

$$Pr[\text{no collisions}] \approx \exp\left\{-\frac{m^2}{2n}\right\}$$

- ▶ **Coupon Collection:**  $n$  items. Buy  $m$  cereal boxes.

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \dots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \dots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \dots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]$ .
- ▶ **Balls in bins:**  $m$  balls into  $n > m$  bins.

$$Pr[\text{no collisions}] \approx \exp\left\{-\frac{m^2}{2n}\right\}$$

- ▶ **Coupon Collection:**  $n$  items. Buy  $m$  cereal boxes.  
 $Pr[\text{miss one specific item}] \approx e^{-\frac{m}{n}};$

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \dots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \dots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \dots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]$ .
- ▶ **Balls in bins:**  $m$  balls into  $n > m$  bins.

$$Pr[\text{no collisions}] \approx \exp\left\{-\frac{m^2}{2n}\right\}$$

- ▶ **Coupon Collection:**  $n$  items. Buy  $m$  cereal boxes.

$$Pr[\text{miss one specific item}] \approx e^{-\frac{m}{n}}; \quad Pr[\text{miss any one of the items}] \leq n e^{-\frac{m}{n}}.$$

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \dots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \dots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \dots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]$ .
- ▶ **Balls in bins:**  $m$  balls into  $n > m$  bins.

$$Pr[\text{no collisions}] \approx \exp\left\{-\frac{m^2}{2n}\right\}$$

- ▶ **Coupon Collection:**  $n$  items. Buy  $m$  cereal boxes.

$$Pr[\text{miss one specific item}] \approx e^{-\frac{m}{n}}; \quad Pr[\text{miss any one of the items}] \leq n e^{-\frac{m}{n}}.$$

Key Mathematical Fact:

# Summary.

## Bayes' Rule, Mutual Independence, Collisions and Collecting

Main results:

- ▶ **Bayes' Rule:**  $Pr[A_m|B] = p_m q_m / (p_1 q_1 + \cdots + p_M q_M)$ .
- ▶ **Product Rule:**  
 $Pr[A_1 \cap \cdots \cap A_n] = Pr[A_1] Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \cdots \cap A_{n-1}]$ .
- ▶ **Balls in bins:**  $m$  balls into  $n > m$  bins.

$$Pr[\text{no collisions}] \approx \exp\left\{-\frac{m^2}{2n}\right\}$$

- ▶ **Coupon Collection:**  $n$  items. Buy  $m$  cereal boxes.

$$Pr[\text{miss one specific item}] \approx e^{-\frac{m}{n}}; \quad Pr[\text{miss any one of the items}] \leq n e^{-\frac{m}{n}}.$$

Key Mathematical Fact:  $\ln(1 - \varepsilon) \approx -\varepsilon$ .