

Alex Psomas: Lecture 19.


1. Distributions
2. Tail bounds

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} Pr[X \geq i].$$

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} \Pr[X \geq i].$$



Probability mass at i , counted i times.

... Same as $\sum_{i=1}^{\infty} i \times \Pr[X = i]$.

A side step: Expected Value of Integer RV

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} Pr[X \geq i].$$

A side step: Expected Value of Integer RV

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} Pr[X \geq i].$$

Proof: One has

$$E[X] = \sum_{i=1}^{\infty} i \times Pr[X = i]$$

A side step: Expected Value of Integer RV

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} Pr[X \geq i].$$

Proof: One has

$$\begin{aligned} E[X] &= \sum_{i=1}^{\infty} i \times Pr[X = i] \\ &= \sum_{i=1}^{\infty} i (Pr[X \geq i] - Pr[X \geq i+1]) \end{aligned}$$

A side step: Expected Value of Integer RV

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} \Pr[X \geq i].$$

Proof: One has

$$\begin{aligned} E[X] &= \sum_{i=1}^{\infty} i \times \Pr[X = i] \\ &= \sum_{i=1}^{\infty} i (\Pr[X \geq i] - \Pr[X \geq i+1]) \\ &= \sum_{i=1}^{\infty} (i \times \Pr[X \geq i] - i \times \Pr[X \geq i+1]) \end{aligned}$$

A side step: Expected Value of Integer RV

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} Pr[X \geq i].$$

Proof: One has

$$\begin{aligned} E[X] &= \sum_{i=1}^{\infty} i \times Pr[X = i] \\ &= \sum_{i=1}^{\infty} i (Pr[X \geq i] - Pr[X \geq i+1]) \\ &= \sum_{i=1}^{\infty} (i \times Pr[X \geq i] - i \times Pr[X \geq i+1]) \\ &= \sum_{i=1}^{\infty} i \times Pr[X \geq i] - \sum_{i=1}^{\infty} i \times Pr[X \geq i+1] \end{aligned}$$

A side step: Expected Value of Integer RV

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} Pr[X \geq i].$$

Proof: One has

$$\begin{aligned} E[X] &= \sum_{i=1}^{\infty} i \times Pr[X = i] \\ &= \sum_{i=1}^{\infty} i (Pr[X \geq i] - Pr[X \geq i+1]) \\ &= \sum_{i=1}^{\infty} (i \times Pr[X \geq i] - i \times Pr[X \geq i+1]) \\ &= \sum_{i=1}^{\infty} i \times Pr[X \geq i] - \sum_{i=1}^{\infty} i \times Pr[X \geq i+1] \\ &= \sum_{i=1}^{\infty} i \times Pr[X \geq i] - \sum_{i=1}^{\infty} (i-1) \times Pr[X \geq i] \end{aligned}$$

A side step: Expected Value of Integer RV

Theorem: For a r.v. X that takes values in $\{0, 1, 2, \dots\}$, one has

$$E[X] = \sum_{i=1}^{\infty} Pr[X \geq i].$$

Proof: One has

$$\begin{aligned} E[X] &= \sum_{i=1}^{\infty} i \times Pr[X = i] \\ &= \sum_{i=1}^{\infty} i (Pr[X \geq i] - Pr[X \geq i+1]) \\ &= \sum_{i=1}^{\infty} (i \times Pr[X \geq i] - i \times Pr[X \geq i+1]) \\ &= \sum_{i=1}^{\infty} i \times Pr[X \geq i] - \sum_{i=1}^{\infty} i \times Pr[X \geq i+1] \\ &= \sum_{i=1}^{\infty} i \times Pr[X \geq i] - \sum_{i=1}^{\infty} (i-1) \times Pr[X \geq i] = \sum_{i=1}^{\infty} Pr[X \geq i]. \end{aligned}$$



Geometric Distribution: Memoryless

Let X be $Geom(p)$. **Theorem**

$$Pr[X > n + m | X > n] = Pr[X > m], m, n \geq 0.$$

Geometric Distribution: Memoryless

I flip a coin (probability of H is p) until I get H .

Geometric Distribution: Memoryless

I flip a coin (probability of H is p) until I get H .

What's the probability that I flip it exactly 100 times?

Geometric Distribution: Memoryless

I flip a coin (probability of H is p) until I get H .

What's the probability that I flip it exactly 100 times? $(1 - p)^{99}p$

Geometric Distribution: Memoryless

I flip a coin (probability of H is p) until I get H .

What's the probability that I flip it exactly 100 times? $(1 - p)^{99}p$

What's the probability that I flip it exactly 100 times if (given that) the first 20 were T ?

Geometric Distribution: Memoryless

I flip a coin (probability of H is p) until I get H .

What's the probability that I flip it exactly 100 times? $(1 - p)^{99}p$

What's the probability that I flip it exactly 100 times if (given that) the first 20 were T ?

Same as flipping it exactly 80 times!

Geometric Distribution: Memoryless

I flip a coin (probability of H is p) until I get H .

What's the probability that I flip it exactly 100 times? $(1 - p)^{99}p$

What's the probability that I flip it exactly 100 times if (given that) the first 20 were T ?

Same as flipping it exactly 80 times!

$(1 - p)^{79}p$.

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$.

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = (2 - p)/p^2$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = (2 - p)/p^2 \quad (\text{tricks})$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = (2 - p)/p^2 \quad (\text{tricks})$$

$$\text{var}[X] = E[X^2] - E[X]^2$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1} p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = (2 - p)/p^2 \quad (\text{tricks})$$

$$\text{var}[X] = E[X^2] - E[X]^2 = \frac{2-p}{p^2} - \frac{1}{p^2}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1}p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = (2 - p)/p^2 \quad (\text{tricks})$$

$$\text{var}[X] = E[X^2] - E[X]^2 = \frac{2-p}{p^2} - \frac{1}{p^2} = \frac{1-p}{p^2}.$$

$$\sigma(X) = \frac{\sqrt{1-p}}{p}$$

Variance of geometric distribution.

X is a geometrically distributed RV with parameter p .

Thus, $\Pr[X = n] = (1 - p)^{n-1} p$ for $n \geq 1$. Recall $E[X] = 1/p$.

$$E[X^2] = (2 - p)/p^2 \quad (\text{tricks})$$

$$\text{var}[X] = E[X^2] - E[X]^2 = \frac{2-p}{p^2} - \frac{1}{p^2} = \frac{1-p}{p^2}.$$

$$\sigma(X) = \frac{\sqrt{1-p}}{p} \approx E[X] \text{ when } p \text{ is small(ish).}$$

Poisson

Experiment: flip a coin n times. The coin is such that

$$Pr[H] = \lambda/n.$$

Random Variable: X - number of heads.

Poisson

Experiment: flip a coin n times. The coin is such that

$$Pr[H] = \lambda/n.$$

Random Variable: X - number of heads. Thus, $X = B(n, \lambda/n)$.

Poisson

Experiment: flip a coin n times. The coin is such that

$$\Pr[H] = \lambda/n.$$

Random Variable: X - number of heads. Thus, $X = B(n, \lambda/n)$.

Poisson Distribution is distribution of X “for large n .”


Poisson

Experiment: flip a coin n times. The coin is such that

$$Pr[H] = \lambda/n.$$

Random Variable: X - number of heads. Thus, $X = B(n, \lambda/n)$.

Poisson Distribution is distribution of X “for large n .”



Poisson

Experiment: flip a coin n times. The coin is such that

$$\Pr[H] = \lambda/n.$$

Random Variable: X - number of heads. Thus, $X = B(n, \lambda/n)$.

Poisson Distribution is distribution of X “for large n .”

Poisson

Experiment: flip a coin n times. The coin is such that

$$\Pr[H] = \lambda/n.$$

Random Variable: X - number of heads. Thus, $X = B(n, \lambda/n)$.

Poisson Distribution is distribution of X “for large n .”

We expect $X \ll n$.

Poisson

Experiment: flip a coin n times. The coin is such that

$$\Pr[H] = \lambda/n.$$

Random Variable: X - number of heads. Thus, $X = B(n, \lambda/n)$.

Poisson Distribution is distribution of X “for large n .”

We expect $X \ll n$. For $m \ll n$ one has

Poisson

Experiment: flip a coin n times. The coin is such that

$$Pr[H] = \lambda/n.$$

Random Variable: X - number of heads. Thus, $X = B(n, \lambda/n)$.

Poisson Distribution is distribution of X “for large n .”

We expect $X \ll n$. For $m \ll n$ one has

$$Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}.$$

Poisson Distribution: Definition and Mean

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \textcolor{blue}{Pr}[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Poisson Distribution: Definition and Mean

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Fact: $E[X] =$

Poisson Distribution: Definition and Mean

Definition Poisson Distribution with parameter $\lambda > 0$

$$X = P(\lambda) \Leftrightarrow \Pr[X = m] = \frac{\lambda^m}{m!} e^{-\lambda}, m \geq 0.$$

Fact: $E[X] = \lambda$.

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .


What is the maximum number of customers you might see?

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?




Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?




Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.


$n = 10$ sub-intervals.

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

$n = 10$ sub-intervals.


Binomial distribution, if only one event/interval!

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

$n = 10$ sub-intervals.


Binomial distribution, if only one event/interval!

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

$n = 10$ sub-intervals.

Binomial distribution, if only one event/interval!


Maybe more...

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

$n = 10$ sub-intervals.

Binomial distribution, if only one event/interval!

Maybe more...


and more.

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

$n = 10$ sub-intervals.

Binomial distribution, if only one event/interval!

Maybe more...

and more.


As n goes to infinity...analyze ...

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

$n = 10$ sub-intervals.

Binomial distribution, if only one event/interval!

Maybe more...

and more.

As n goes to infinity...analyze ...

.... $Pr[X = i] = \binom{n}{i} p^i (1 - p)^{n-i}$.


derive simple expression.

Poisson and Queueing.

Poisson: Distribution of how many events in an interval?

Average: λ .

What is the maximum number of customers you might see?



Idea: Cut into intervals so that “sum of Bernoulli (indicators)”.

$n = 10$ sub-intervals.

Binomial distribution, if only one event/interval!

Maybe more...

and more.

As n goes to infinity...analyze ...

$$\dots Pr[X = i] = \binom{n}{i} p^i (1 - p)^{n-i}.$$

derive simple expression.

...And we get the Poisson distribution!

When to use Poisson

If an event can occur $0, 1, 2, \dots$ times in an interval,

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ
and events are independent

When to use Poisson

If an event can occur 0,1,2,... times in an interval,

and the average number of events per interval is λ

and events are independent

and the probability of an event in an interval is proportional to the interval's length,

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ
and events are independent
and the probability of an event in an interval is proportional to
the interval's length,
then it might be appropriate to use Poisson distribution.

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ
and events are independent
and the probability of an event in an interval is proportional to
the interval's length,
then it might be appropriate to use Poisson distribution.

$$Pr[k \text{ events in interval}] = \frac{\lambda^k}{k!} e^{-\lambda}$$

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ
and events are independent
and the probability of an event in an interval is proportional to
the interval's length,
then it might be appropriate to use Poisson distribution.

$$Pr[k \text{ events in interval}] = \frac{\lambda^k}{k!} e^{-\lambda}$$

Examples:

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ
and events are independent
and the probability of an event in an interval is proportional to
the interval's length,
then it might be appropriate to use Poisson distribution.

$$Pr[k \text{ events in interval}] = \frac{\lambda^k}{k!} e^{-\lambda}$$

Examples: photons arriving at a telescope,

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ
and events are independent
and the probability of an event in an interval is proportional to
the interval's length,
then it might be appropriate to use Poisson distribution.

$$Pr[k \text{ events in interval}] = \frac{\lambda^k}{k!} e^{-\lambda}$$

Examples: photons arriving at a telescope, telephone calls
arriving in a system,

When to use Poisson

If an event can occur 0,1,2,... times in an interval,
and the average number of events per interval is λ
and events are independent
and the probability of an event in an interval is proportional to
the interval's length,
then it might be appropriate to use Poisson distribution.

$$Pr[k \text{ events in interval}] = \frac{\lambda^k}{k!} e^{-\lambda}$$

Examples: photons arriving at a telescope, telephone calls arriving in a system, the number of mutations on a strand of DNA per unit length...

Simeon Poisson

The Poisson distribution is named after:

Simeon Poisson

The Poisson distribution is named after:



“Life is good for only two things: doing mathematics and teaching it.”

Review: Distributions

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$
 $Var[X] = \frac{n^2-1}{12};$
- ▶ $Geom(p) : Pr[X = n] =$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$
 $Var[X] = \frac{n^2-1}{12};$
- ▶ $Geom(p) : Pr[X = n] = (1 - p)^{n-1} p, n = 1, 2, \dots;$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$
 $Var[X] = \frac{n^2-1}{12};$
- ▶ $Geom(p) : Pr[X = n] = (1 - p)^{n-1} p, n = 1, 2, \dots;$
 $E[X] = \frac{1}{p};$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$
 $Var[X] = \frac{n^2-1}{12};$
- ▶ $Geom(p) : Pr[X = n] = (1 - p)^{n-1} p, n = 1, 2, \dots;$
 $E[X] = \frac{1}{p};$
 $Var[X] = \frac{1-p}{p^2};$
- ▶ $P(\lambda) :$

Review: Distributions

- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$
 $Var[X] = \frac{n^2-1}{12};$
- ▶ $Geom(p) : Pr[X = n] = (1 - p)^{n-1} p, n = 1, 2, \dots;$
 $E[X] = \frac{1}{p};$
 $Var[X] = \frac{1-p}{p^2};$
- ▶ $P(\lambda) : Pr[X = n] =$

Review: Distributions


- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$
 $Var[X] = \frac{n^2-1}{12};$
- ▶ $Geom(p) : Pr[X = n] = (1 - p)^{n-1} p, n = 1, 2, \dots;$
 $E[X] = \frac{1}{p};$
 $Var[X] = \frac{1-p}{p^2};$
- ▶ $P(\lambda) : Pr[X = n] = \frac{\lambda^n}{n!} e^{-\lambda}, n \geq 0;$

Review: Distributions


- ▶ $Bern(p) : Pr[X = 1] = p;$
 $E[X] = p;$
 $Var[X] = p(1 - p);$
- ▶ $B(n, p) : Pr[X = m] = \binom{n}{m} p^m (1 - p)^{n-m}, m = 0, \dots, n;$
 $E[X] = np;$
 $Var[X] = np(1 - p);$
- ▶ $U[1, \dots, n] : Pr[X = m] = \frac{1}{n}, m = 1, \dots, n;$
 $E[X] = \frac{n+1}{2};$
 $Var[X] = \frac{n^2-1}{12};$
- ▶ $Geom(p) : Pr[X = n] = (1 - p)^{n-1} p, n = 1, 2, \dots;$
 $E[X] = \frac{1}{p};$
 $Var[X] = \frac{1-p}{p^2};$
- ▶ $P(\lambda) : Pr[X = n] = \frac{\lambda^n}{n!} e^{-\lambda}, n \geq 0;$
 $E[X] = \lambda;$
 $Var[X] = \lambda.$

Inequalities: An Overview

Distribution




Markov



$$Pr[X > a]$$

Chebyshev



$$Pr[|X - \mu| > \epsilon]$$

Andrey Markov

**Andrey (Andrei) Andreyevich
Markov**



Born	14 June 1856 N.S. Ryazan, Russian Empire
Died	20 July 1922 (aged 66) Petrograd, Russian SFSR

Andrey Markov

**Andrey (Andrei) Andreyevich
Markov**



Born	14 June 1856 N.S. Ryazan, Russian Empire
Died	20 July 1922 (aged 66) Petrograd, Russian SFSR

[Andrey Markov](#) is best known for his work on stochastic processes. A primary subject of his research later became known as Markov chains and Markov processes.

Andrey Markov

**Andrey (Andrei) Andreyevich
Markov**



Born 14 June 1856 N.S.
Ryazan, Russian Empire

Died 20 July 1922 (aged 66)
Petrograd, Russian SFSR

[Andrey Markov](#) is best known for his work on stochastic processes. A primary subject of his research later became known as Markov chains and Markov processes.

[Pafnuty Chebyshev](#) was one of his teachers.

Andrey Markov

**Andrey (Andrei) Andreyevich
Markov**



Born 14 June 1856 N.S.
Ryazan, Russian Empire

Died 20 July 1922 (aged 66)
Petrograd, Russian SFSR

[Andrey Markov](#) is best known for his work on stochastic processes. A primary subject of his research later became known as Markov chains and Markov processes.

[Pafnuty Chebyshev](#) was one of his teachers.

Markov's inequality

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev.

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing.

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing. Then, for a non-negative random variable X

$$Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}, \text{ for all } a \text{ such that } f(a) > 0.$$

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing. Then, for a non-negative random variable X

$$Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}, \text{ for all } a \text{ such that } f(a) > 0.$$

Proof:

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing. Then, for a non-negative random variable X

$$Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}, \text{ for all } a \text{ such that } f(a) > 0.$$

Proof:

Observe that

$$1_{\{X \geq a\}} \leq \frac{f(X)}{f(a)}.$$

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing. Then, for a non-negative random variable X

$$\Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}, \text{ for all } a \text{ such that } f(a) > 0.$$

Proof:

Observe that

$$1_{\{X \geq a\}} \leq \frac{f(X)}{f(a)}.$$

Indeed, if $X < a$, the inequality reads $0 \leq f(X)/f(a)$, which holds since $f(\cdot) \geq 0$.

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing. Then, for a non-negative random variable X

$$\Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}, \text{ for all } a \text{ such that } f(a) > 0.$$

Proof:

Observe that

$$1_{\{X \geq a\}} \leq \frac{f(X)}{f(a)}.$$

Indeed, if $X < a$, the inequality reads $0 \leq f(X)/f(a)$, which holds since $f(\cdot) \geq 0$. Also, if $X \geq a$, it reads $1 \leq f(X)/f(a)$, which holds since $f(\cdot)$ is nondecreasing.

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing. Then, for a non-negative random variable X

$$Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}, \text{ for all } a \text{ such that } f(a) > 0.$$

Proof:

Observe that

$$1_{\{X \geq a\}} \leq \frac{f(X)}{f(a)}.$$

Indeed, if $X < a$, the inequality reads $0 \leq f(X)/f(a)$, which holds since $f(\cdot) \geq 0$. Also, if $X \geq a$, it reads $1 \leq f(X)/f(a)$, which holds since $f(\cdot)$ is nondecreasing. Expectation is monotone: if $X(\omega) \leq Y(\omega)$ for all ω , then $E[X] \leq E[Y]$.

Markov's inequality

The inequality is named after Andrey Markov, although it appeared earlier in the work of Pafnuty Chebyshev. It should be (and is sometimes) called Chebyshev's first inequality.

Theorem Markov's Inequality (the fancy version)

Assume $f : \mathfrak{X} \rightarrow [0, \infty)$ is nondecreasing. Then, for a non-negative random variable X

$$Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}, \text{ for all } a \text{ such that } f(a) > 0.$$

Proof:


Observe that

$$1_{\{X \geq a\}} \leq \frac{f(X)}{f(a)}.$$

Indeed, if $X < a$, the inequality reads $0 \leq f(X)/f(a)$, which holds since $f(\cdot) \geq 0$. Also, if $X \geq a$, it reads $1 \leq f(X)/f(a)$, which holds since $f(\cdot)$ is nondecreasing. Expectation is monotone: if $X(\omega) \leq Y(\omega)$ for all ω , then $E[X] \leq E[Y]$. Therefore,

$$E[1_{\{X \geq a\}}] \leq \frac{E[f(X)]}{f(a)}.$$

A picture



$$f(a)1\{X \geq a\} \leq f(x) \Rightarrow 1\{X \geq a\} \leq \frac{f(X)}{f(a)}$$

$$\Rightarrow Pr[X \geq a] \leq \frac{E[f(X)]}{f(a)}$$

Markov Inequality Note

A more common version of Markov is for $f(x) = x$:

Markov Inequality Note

A more common version of Markov is for $f(x) = x$:

Theorem For a non-negative random variable X , and any $a > 0$,

$$Pr[X \geq a] \leq \frac{E[X]}{a}.$$

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$.

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$. Recall that $E[X] =$

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$. Recall that $E[X] = \frac{1}{p}$ and $E[X^2] =$

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$. Recall that $E[X] = \frac{1}{p}$ and $E[X^2] = \frac{2-p}{p^2}$.

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$. Recall that $E[X] = \frac{1}{p}$ and $E[X^2] = \frac{2-p}{p^2}$.

Choosing $f(x) = x$, we
get

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$. Recall that $E[X] = \frac{1}{p}$ and $E[X^2] = \frac{2-p}{p^2}$.

Choosing $f(x) = x$, we
get

$$\Pr[X \geq a] \leq \frac{E[X]}{a} = \frac{1}{ap}.$$

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$. Recall that $E[X] = \frac{1}{p}$ and $E[X^2] = \frac{2-p}{p^2}$.

Choosing $f(x) = x$, we
get

$$\Pr[X \geq a] \leq \frac{E[X]}{a} = \frac{1}{ap}.$$

Choosing $f(x) = x^2$,
we get

Markov Inequality Example: Geom(p)

Let $X \sim \text{Geom}(p)$. Recall that $E[X] = \frac{1}{p}$ and $E[X^2] = \frac{2-p}{p^2}$.

Choosing $f(x) = x$, we
get

$$\Pr[X \geq a] \leq \frac{E[X]}{a} = \frac{1}{ap}.$$

Choosing $f(x) = x^2$,
we get

$$\Pr[X \geq a] \leq \frac{E[X^2]}{a^2} = \frac{2-p}{p^2 a^2}.$$

Markov Inequality Example: Geom(p)


Let $X \sim \text{Geom}(p)$. Recall that $E[X] = \frac{1}{p}$ and $E[X^2] = \frac{2-p}{p^2}$.

Choosing $f(x) = x$, we get

$$\Pr[X \geq a] \leq \frac{E[X]}{a} = \frac{1}{ap}.$$

Choosing $f(x) = x^2$, we get

$$\Pr[X \geq a] \leq \frac{E[X^2]}{a^2} = \frac{2-p}{p^2 a^2}.$$



Markov's inequality example

$$Pr[X \geq a] \leq \frac{E[X]}{a}.$$

Markov's inequality example

$$Pr[X \geq a] \leq \frac{E[X]}{a}.$$

What is a bound on the probability that a random X takes value \geq than twice its' expectation?

Markov's inequality example

$$Pr[X \geq a] \leq \frac{E[X]}{a}.$$

What is a bound on the probability that a random X takes value \geq than twice its' expectation?

$$\frac{1}{2}.$$

Markov's inequality example

$$Pr[X \geq a] \leq \frac{E[X]}{a}.$$

What is a bound on the probability that a random X takes value \geq than twice its' expectation?

$\frac{1}{2}$. It can't be that more than half of the people are twice above the average!

Markov's inequality example

$$Pr[X \geq a] \leq \frac{E[X]}{a}.$$

What is a bound on the probability that a random X takes value \geq than twice its' expectation?

$\frac{1}{2}$. It can't be that more than half of the people are twice above the average!

What is a bound on the probability that a random X takes value \geq than k times its' expectation?

Markov's inequality example

$$Pr[X \geq a] \leq \frac{E[X]}{a}.$$

What is a bound on the probability that a random X takes value \geq than twice its' expectation?

$\frac{1}{2}$. It can't be that more than half of the people are twice above the average!

What is a bound on the probability that a random X takes value \geq than k times its' expectation?

$\frac{1}{k}$.

Markov's inequality example

Flip a coin n times. Probability of H is p .

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$$X \sim B(n, p).$$

$$E[X] =$$

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$$X \sim B(n, p).$$

$$E[X] = np.$$

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. Say $n = 1000$ and $p = 0.5$. $E[X] = 500$.

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. Say $n = 1000$ and $p = 0.5$. $E[X] = 500$.

Markov says that $Pr[X \geq 600] \leq$

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. Say $n = 1000$ and $p = 0.5$. $E[X] = 500$.

Markov says that $Pr[X \geq 600] \leq \frac{1000*0.5}{600} = \frac{5}{6} \approx 0.83$

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. Say $n = 1000$ and $p = 0.5$. $E[X] = 500$.

Markov says that $Pr[X \geq 600] \leq \frac{1000*0.5}{600} = \frac{5}{6} \approx 0.83$

Actual probability: < 0.000001

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. Say $n = 1000$ and $p = 0.5$. $E[X] = 500$.

Markov says that $Pr[X \geq 600] \leq \frac{1000*0.5}{600} = \frac{5}{6} \approx 0.83$

Actual probability: < 0.000001

Notice:

Markov's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. Say $n = 1000$ and $p = 0.5$. $E[X] = 500$.

Markov says that $Pr[X \geq 600] \leq \frac{1000*0.5}{600} = \frac{5}{6} \approx 0.83$

Actual probability: < 0.000001

Notice: Same bound for 10 coins and $Pr[X \geq 6]$

Chebyshev's Inequality

This is Pafnuty's inequality:

Chebyshev's Inequality

This is Pafnuty's inequality:

Theorem:

$$Pr[|X - E[X]| \geq a] \leq \frac{\text{var}[X]}{a^2}, \text{ for all } a > 0.$$

Chebyshev's Inequality

This is Pafnuty's inequality:

Theorem:

$$Pr[|X - E[X]| \geq a] \leq \frac{\text{var}[X]}{a^2}, \text{ for all } a > 0.$$

Proof: Let $Y = |X - E[X]|$ and $f(y) = y^2$.

Chebyshev's Inequality

This is Pafnuty's inequality:

Theorem:

$$Pr[|X - E[X]| \geq a] \leq \frac{\text{var}[X]}{a^2}, \text{ for all } a > 0.$$

Proof: Let $Y = |X - E[X]|$ and $f(y) = y^2$. Then,

$$Pr[Y \geq a] \leq \frac{E[f(Y)]}{f(a)}$$

Chebyshev's Inequality

This is Pafnuty's inequality:

Theorem:

$$Pr[|X - E[X]| \geq a] \leq \frac{\text{var}[X]}{a^2}, \text{ for all } a > 0.$$

Proof: Let $Y = |X - E[X]|$ and $f(y) = y^2$. Then,

$$Pr[Y \geq a] \leq \frac{E[f(Y)]}{f(a)} = \frac{E[|X - E[X]|^2]}{a^2}$$

Chebyshev's Inequality

This is Pafnuty's inequality:

Theorem:

$$Pr[|X - E[X]| \geq a] \leq \frac{\text{var}[X]}{a^2}, \text{ for all } a > 0.$$

Proof: Let $Y = |X - E[X]|$ and $f(y) = y^2$. Then,

$$Pr[Y \geq a] \leq \frac{E[f(Y)]}{f(a)} = \frac{E[|X - E[X]|^2]}{a^2} = \frac{\text{var}[X]}{a^2}.$$



Chebyshev's Inequality

This is Pafnuty's inequality:

Theorem:

$$Pr[|X - E[X]| \geq a] \leq \frac{\text{var}[X]}{a^2}, \text{ for all } a > 0.$$

Proof: Let $Y = |X - E[X]|$ and $f(y) = y^2$. Then,

$$Pr[Y \geq a] \leq \frac{E[f(Y)]}{f(a)} = \frac{E[|X - E[X]|^2]}{a^2} = \frac{\text{var}[X]}{a^2}.$$



This result confirms that the variance measures the “deviations from the mean.”

Chebyshev and Poisson

Chebyshev and Poisson

Let $X = P(\lambda)$. Then, $E[X] = \lambda$ and $\text{var}[X] =$

Chebyshev and Poisson

Let $X = P(\lambda)$. Then, $E[X] = \lambda$ and $\text{var}[X] = \lambda$.

Chebyshev and Poisson


Let $X = P(\lambda)$. Then, $E[X] = \lambda$ and $\text{var}[X] = \lambda$. Thus,

$$\Pr[|X - \lambda| \geq n] \leq \frac{\text{var}[X]}{n^2} = \frac{\lambda}{n^2}.$$

Chebyshev and Poisson

Let $X = P(\lambda)$. Then, $E[X] = \lambda$ and $\text{var}[X] = \lambda$. Thus,

$$\Pr[|X - \lambda| \geq n] \leq \frac{\text{var}[X]}{n^2} = \frac{\lambda}{n^2}.$$



Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$$X \sim B(n, p).$$

$$E[X] = np.$$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$$X \sim B(n, p).$$

$$E[X] = np. \text{ Var}[X] = np(1 - p).$$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$$X \sim B(n, p).$$

$$E[X] = np. \quad \text{Var}[X] = np(1 - p).$$

$$\text{Say } n = 1000 \text{ and } p = 0.5. \quad E[X] = 500. \quad \text{Var}[X] = 250.$$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev says that $Pr[X \geq 600]$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev says that $Pr[X \geq 600] = Pr[X - 500 \geq 100]$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev says that $Pr[X \geq 600] = Pr[X - 500 \geq 100] \leq Pr[|X - 500| \geq 100]$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev says that $Pr[X \geq 600] = Pr[X - 500 \geq 100] \leq Pr[|X - 500| \geq 100] \leq \frac{250}{10000} = 0.025$

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev says that $Pr[X \geq 600] = Pr[X - 500 \geq 100] \leq Pr[|X - 500| \geq 100] \leq \frac{250}{10000} = 0.025$

Actual probability: < 0.000001

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev says that $Pr[X \geq 600] = Pr[X - 500 \geq 100] \leq Pr[|X - 500| \geq 100] \leq \frac{250}{10000} = 0.025$

Actual probability: < 0.000001

Notice:

Chebyshev's inequality example

Flip a coin n times. Probability of H is p . X counts the number of heads.

X follows the Binomial distribution with parameters n and p .

$X \sim B(n, p)$.

$E[X] = np$. $Var[X] = np(1 - p)$.

Say $n = 1000$ and $p = 0.5$. $E[X] = 500$. $Var[X] = 250$.

Markov says that $Pr[X \geq 600] \leq \frac{500}{600} = \frac{5}{6} \approx 0.83$

Chebyshev says that $Pr[X \geq 600] = Pr[X - 500 \geq 100] \leq Pr[|X - 500| \geq 100] \leq \frac{250}{10000} = 0.025$

Actual probability: < 0.000001

Notice: If we had 100 coins, the bound for $Pr[X \geq 60]$ would be different.

Chebyshev's inequality example continued

What if we had more coins?

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.

p is still 0.5

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] =$$

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] = \frac{1}{n}E[\sum X_i] =$$

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] = \frac{1}{n}E[\sum X_i] = \frac{1}{n}np = p = 0.5$$

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] = \frac{1}{n}E[\sum X_i] = \frac{1}{n}np = p = 0.5$$

$$\text{Var}[Y_n] =$$

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] = \frac{1}{n} E[\sum X_i] = \frac{1}{n} np = p = 0.5$$

$$\text{Var}[Y_n] = \frac{1}{n^2} \text{Var}[\sum X_i] =$$

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] = \frac{1}{n} E[\sum X_i] = \frac{1}{n} np = p = 0.5$$

$$\text{Var}[Y_n] = \frac{1}{n^2} \text{Var}[\sum X_i] = \frac{1}{n^2} np(1-p) = \frac{p(1-p)}{n} = \frac{1}{4n}$$

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] = \frac{1}{n} E[\sum X_i] = \frac{1}{n} np = p = 0.5$$

$$\text{Var}[Y_n] = \frac{1}{n^2} \text{Var}[\sum X_i] = \frac{1}{n^2} np(1-p) = \frac{p(1-p)}{n} = \frac{1}{4n}$$

Let's try to bound how likely it is that the fraction of H 's differs from 50%.

Chebyshev's inequality example continued

What if we had more coins?

Also, let's count the fraction of H instead of their number.
 p is still 0.5

Let X_i be the indicator random variable for the i -th coin.

Define

$$Y_n = \frac{X_1 + \cdots + X_n}{n}, \text{ for } n \geq 1.$$

$$E[Y_n] = \frac{1}{n} E[\sum X_i] = \frac{1}{n} np = p = 0.5$$

$$\text{Var}[Y_n] = \frac{1}{n^2} \text{Var}[\sum X_i] = \frac{1}{n^2} np(1-p) = \frac{p(1-p)}{n} = \frac{1}{4n}$$

Let's try to bound how likely it is that the fraction of H 's differs from 50%.

$$\Pr[|Y_n - 0.5| \geq \varepsilon]?$$

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon]$$

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

As $n \rightarrow \infty$, this probability goes to zero.

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

As $n \rightarrow \infty$, this probability goes to zero.

In fact, for any $\varepsilon > 0$,

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

As $n \rightarrow \infty$, this probability goes to zero.

In fact, for any $\varepsilon > 0$, as $n \rightarrow \infty$,

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$\Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: \Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

As $n \rightarrow \infty$, this probability goes to zero.

In fact, for any $\varepsilon > 0$, as $n \rightarrow \infty$, the probability that the fraction of H s is within $\varepsilon > 0$ of 50% approaches 1:

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

As $n \rightarrow \infty$, this probability goes to zero.

In fact, for any $\varepsilon > 0$, as $n \rightarrow \infty$, the probability that the fraction of H s is within $\varepsilon > 0$ of 50% approaches 1:

$$Pr[|Y_n - 0.5| \leq \varepsilon] \rightarrow 1.$$

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

As $n \rightarrow \infty$, this probability goes to zero.

In fact, for any $\varepsilon > 0$, as $n \rightarrow \infty$, the probability that the fraction of H s is within $\varepsilon > 0$ of 50% approaches 1:

$$Pr[|Y_n - 0.5| \leq \varepsilon] \rightarrow 1.$$

This is an example of the Law of Large Numbers.

Chebyshev's inequality example continued

$$E[Y_n] = 0.5, \text{Var}[Y_n] = \frac{1}{4n}.$$

$$Pr[|Y_n - 0.5| \geq \varepsilon] \leq \frac{\text{Var}[Y_n]}{\varepsilon^2} = \frac{1}{4n\varepsilon^2}$$

$$\text{For } \varepsilon = 0.01: Pr[|Y_n - 0.5| \geq 0.01] \leq \frac{2500}{n}$$

For $n = 250,000$ this is 1%.

As $n \rightarrow \infty$, this probability goes to zero.

In fact, for any $\varepsilon > 0$, as $n \rightarrow \infty$, the probability that the fraction of H s is within $\varepsilon > 0$ of 50% approaches 1:

$$Pr[|Y_n - 0.5| \leq \varepsilon] \rightarrow 1.$$

This is an example of the Law of Large Numbers.

We look at a general case next.

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ .

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

$$Pr[|Y_n - \mu| \geq \varepsilon] \leq \frac{\text{var}[Y_n]}{\varepsilon^2}$$

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

$$Pr[|Y_n - \mu| \geq \varepsilon] \leq \frac{\text{var}[Y_n]}{\varepsilon^2} = \frac{\text{var}[X_1 + \dots + X_n]}{n^2 \varepsilon^2}$$

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

$$\begin{aligned} Pr[|Y_n - \mu| \geq \varepsilon] &\leq \frac{\text{var}[Y_n]}{\varepsilon^2} = \frac{\text{var}[X_1 + \dots + X_n]}{n^2 \varepsilon^2} \\ &= \frac{n \text{var}[X_1]}{n^2 \varepsilon^2} \end{aligned}$$

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

$$\begin{aligned} Pr[|Y_n - \mu| \geq \varepsilon] &\leq \frac{\text{var}[Y_n]}{\varepsilon^2} = \frac{\text{var}[X_1 + \dots + X_n]}{n^2 \varepsilon^2} \\ &= \frac{n \text{var}[X_1]}{n^2 \varepsilon^2} = \frac{\text{var}[X_1]}{n \varepsilon^2} \end{aligned}$$

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

$$\begin{aligned} Pr[|Y_n - \mu| \geq \varepsilon] &\leq \frac{\text{var}[Y_n]}{\varepsilon^2} = \frac{\text{var}[X_1 + \dots + X_n]}{n^2 \varepsilon^2} \\ &= \frac{n \text{var}[X_1]}{n^2 \varepsilon^2} = \frac{\text{var}[X_1]}{n \varepsilon^2} \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned}$$

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

$$\begin{aligned} Pr[|Y_n - \mu| \geq \varepsilon] &\leq \frac{\text{var}[Y_n]}{\varepsilon^2} = \frac{\text{var}[X_1 + \dots + X_n]}{n^2 \varepsilon^2} \\ &= \frac{n \text{var}[X_1]}{n^2 \varepsilon^2} = \frac{\text{var}[X_1]}{n \varepsilon^2} \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned}$$

(I used that variance is finite for this proof.)

Weak Law of Large Numbers

Theorem Weak Law of Large Numbers

Let X_1, X_2, \dots be pairwise independent with the same distribution and mean μ . Then, for all $\varepsilon > 0$,

$$Pr\left[\left|\frac{X_1 + \dots + X_n}{n} - \mu\right| \geq \varepsilon\right] \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Proof:

Let $Y_n = \frac{X_1 + \dots + X_n}{n}$. Then

$$\begin{aligned} Pr[|Y_n - \mu| \geq \varepsilon] &\leq \frac{\text{var}[Y_n]}{\varepsilon^2} = \frac{\text{var}[X_1 + \dots + X_n]}{n^2 \varepsilon^2} \\ &= \frac{n \text{var}[X_1]}{n^2 \varepsilon^2} = \frac{\text{var}[X_1]}{n \varepsilon^2} \rightarrow 0, \text{ as } n \rightarrow \infty. \end{aligned}$$

(I used that variance is finite for this proof. More complicated proof without this assumption.)



Confidence intervals example

Say p in the previous example was unknown.

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

$$E[\hat{p}] =$$

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

$$E[\hat{p}] = E[\frac{1}{n} \sum_{i=1}^n X_i]$$

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

$$E[\hat{p}] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = p$$

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

$$E[\hat{p}] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = p$$

$$\text{Var}[\hat{p}] =$$

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

$$E[\hat{p}] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = p$$

$$\text{Var}[\hat{p}] = \text{Var}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \text{Var}\left[\sum_{i=1}^n X_i\right] = \frac{p(1-p)}{n}$$

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

$$E[\hat{p}] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = p$$

$$\text{Var}[\hat{p}] = \text{Var}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \text{Var}\left[\sum_{i=1}^n X_i\right] = \frac{p(1-p)}{n}$$

$$\Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{\text{Var}[\hat{p}]}{\varepsilon^2}$$

Confidence intervals example

Say p in the previous example was unknown.

If you flip n coins, your estimate for p is $\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i$.

You many coins do you have to flip to make sure that your estimation \hat{p} is within 0.01 of the true p , with probability at least 95%?

$$E[\hat{p}] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = p$$

$$\text{Var}[\hat{p}] = \text{Var}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n^2} \text{Var}\left[\sum_{i=1}^n X_i\right] = \frac{p(1-p)}{n}$$

$$\Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{\text{Var}[\hat{p}]}{\varepsilon^2} = \frac{p(1-p)}{n\varepsilon^2}$$

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

We want to make $Pr[|\hat{p} - p| \leq 0.01]$ at least 0.95.

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

We want to make $Pr[|\hat{p} - p| \leq 0.01]$ at least 0.95.

Same as $Pr[|\hat{p} - p| \geq 0.01]$ at most 0.05.

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

We want to make $Pr[|\hat{p} - p| \leq 0.01]$ at least 0.95.

Same as $Pr[|\hat{p} - p| \geq 0.01]$ at most 0.05.

It's sufficient to have $\frac{p(1-p)}{n\varepsilon^2} \leq 0.05$

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

We want to make $Pr[|\hat{p} - p| \leq 0.01]$ at least 0.95.

Same as $Pr[|\hat{p} - p| \geq 0.01]$ at most 0.05.

It's sufficient to have $\frac{p(1-p)}{n\varepsilon^2} \leq 0.05$ or $n \geq \frac{20p(1-p)}{\varepsilon^2}$.

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

We want to make $Pr[|\hat{p} - p| \leq 0.01]$ at least 0.95.

Same as $Pr[|\hat{p} - p| \geq 0.01]$ at most 0.05.

It's sufficient to have $\frac{p(1-p)}{n\varepsilon^2} \leq 0.05$ or $n \geq \frac{20p(1-p)}{\varepsilon^2}$.

$p(1-p)$ is maximized for $p = 0.5$.

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

We want to make $Pr[|\hat{p} - p| \leq 0.01]$ at least 0.95.

Same as $Pr[|\hat{p} - p| \geq 0.01]$ at most 0.05.

It's sufficient to have $\frac{p(1-p)}{n\varepsilon^2} \leq 0.05$ or $n \geq \frac{20p(1-p)}{\varepsilon^2}$.

$p(1-p)$ is maximized for $p = 0.5$. Therefore it's sufficient to have $n \geq \frac{5}{\varepsilon^2}$.

Confidence intervals example continued

Estimation \hat{p} is within 0.01 of the true p , with probability at least 95%.

$$Pr[|\hat{p} - p| \geq \varepsilon] \leq \frac{p(1-p)}{n\varepsilon^2}$$

We want to make $Pr[|\hat{p} - p| \leq 0.01]$ at least 0.95.

Same as $Pr[|\hat{p} - p| \geq 0.01]$ at most 0.05.

It's sufficient to have $\frac{p(1-p)}{n\varepsilon^2} \leq 0.05$ or $n \geq \frac{20p(1-p)}{\varepsilon^2}$.

$p(1-p)$ is maximized for $p = 0.5$. Therefore it's sufficient to have $n \geq \frac{5}{\varepsilon^2}$.

For $\varepsilon = 0.01$ we get that $n \geq 50000$ coins are sufficient.

Today's gig: ?

Today's gig: ?

Gigs so far:

1. How to tell random from human.
2. Monty Hall.
3. Birthday Paradox.
4. St. Petersburg paradox.
5. Simpson's paradox.
6. Two envelopes problem.

Today:

Today's gig: ?

Gigs so far:

1. How to tell random from human.
2. Monty Hall.
3. Birthday Paradox.
4. St. Petersburg paradox.
5. Simpson's paradox.
6. Two envelopes problem.

Today: A magic trick.

Summary

- ▶ Variance of Geometric.
- ▶ Markov's Inequality
- ▶ Chebyshev's Inequality.