

# A Random Walk through CS70, Pt. II: Probability

CS70 Summer 2016 - Lecture 8C

---

David Dinh

10 August 2016

UC Berkeley

# Today

Same as yesterday (and tomorrow). Review, applications, gigs, cool examples, research questions...

Probability today!

Map of outcomes in a probability space  $\Omega$  to values in  $[0, 1]$ :

$$\sum_{\omega \in \Omega} \Pr[\omega] = 1$$

Events: set of outcomes.  $\Pr[E] = \sum_{\omega \in E} \Pr[\omega]$ .

# Fundamentals

Map of outcomes in a probability space  $\Omega$  to values in  $[0, 1]$ :

$$\sum_{\omega \in \Omega} \Pr[\omega] = 1$$

Events: set of outcomes.  $\Pr[E] = \sum_{\omega \in E} \Pr[\omega]$ .

Inclusion-Exclusion:  $\Pr[A \cup B] = \Pr[A] + \Pr[B] - \Pr[A \cap B]$ .

Union bound:  $\Pr[A_1 \cup A_2 \cup \dots \cup A_n] \leq \Pr[A_1] + \Pr[A_2] + \dots \Pr[A_n]$ .

Total probability: if  $A_1, \dots, A_n$  partition the entire sample space (disjoint, covers all of it), then  $\Pr[B] = \Pr[B \cap A_1] + \dots + \Pr[B \cap A_n]$ .

# Conditional Probability

Definition:

$$\Pr[A|B] = \frac{\Pr[A \cap B]}{\Pr[B]} .$$

Live demo.

Definition:

$$\Pr[A|B] = \frac{\Pr[A \cap B]}{\Pr[B]} .$$

Live demo.

From definition:  $\Pr[A \cap B] = \Pr[A] \Pr[B|A]$ .

Or, generally:  $\Pr[A_1 \cap \dots \cap A_n] = \Pr[A_1] \Pr[A_2|A_1] \dots \Pr[A_n|A_1 \cap \dots \cap A_{n-1}]$ .

# Bayes' Theorem

$$\Pr[A|B] = \frac{\Pr[A] \Pr[B|A]}{\Pr[B]}$$

# Bayes' Theorem

$$\Pr[A|B] = \frac{\Pr[A] \Pr[B|A]}{\Pr[B]}$$

Or if I know for sure that exactly one of  $A_1, \dots, A_n$  hold, then:

$$\Pr[A_k|B] = \frac{\Pr[A_k] \Pr[B|A_k]}{\sum_k \Pr[A_k] \Pr[B|A_k]} \ .$$



# Bayes' Theorem

$$\Pr[A|B] = \frac{\Pr[A] \Pr[B|A]}{\Pr[B]}$$

Or if I know for sure that exactly one of  $A_1, \dots, A_n$  hold, then:

$$\Pr[A_k|B] = \frac{\Pr[A_k] \Pr[B|A_k]}{\sum_k \Pr[A_k] \Pr[B|A_k]} .$$

Useful theorem for inference (updating beliefs). Heavily used in AI.  
CS188.

## Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

## Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

Random variables  $X, Y$  are independent if the events  $Y = a$  and  $X = b$  are independent for all  $a, b$ .

## Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

Random variables  $X, Y$  are independent if the events  $Y = a$  and  $X = b$  are independent for all  $a, b$ . If  $X, Y$  independent, then  $f(X), g(Y)$  independent for all  $f, g$ .

## Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

Random variables  $X, Y$  are independent if the events  $Y = a$  and  $X = b$  are independent for all  $a, b$ . If  $X, Y$  independent, then  $f(X), g(Y)$  independent for all  $f, g$ .

Expectation:  $E[X] = \sum_t t \Pr[X = t]$

Tail sum: for nonnegative r.v.  $X$ :  $E[X] = \sum_{i=0}^{\infty} \Pr[X > i]$ .

## Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

Random variables  $X, Y$  are independent if the events  $Y = a$  and  $X = b$  are independent for all  $a, b$ . If  $X, Y$  independent, then  $f(X), g(Y)$  independent for all  $f, g$ .

Expectation:  $E[X] = \sum_t t \Pr[X = t]$

Tail sum: for nonnegative r.v.  $X$ :  $E[X] = \sum_{i=0}^{\infty} \Pr[X > i]$ .

Expectation of function:  $E[g(X)] = \sum_t g(t) \Pr[X = t]$

# Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

Random variables  $X, Y$  are independent if the events  $Y = a$  and  $X = b$  are independent for all  $a, b$ . If  $X, Y$  independent, then  $f(X), g(Y)$  independent for all  $f, g$ .

Expectation:  $E[X] = \sum_t t \Pr[X = t]$

Tail sum: for nonnegative r.v.  $X$ :  $E[X] = \sum_{i=0}^{\infty} \Pr[X > i]$ .

Expectation of function:  $E[g(X)] = \sum_t g(t) \Pr[X = t]$

Variance:  $\text{Var}[X] = E[(X - E[X])^2] = E[X^2] - E[X]^2$

Standard deviation: square root of variance.

# Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

Random variables  $X, Y$  are independent if the events  $Y = a$  and  $X = b$  are independent for all  $a, b$ . If  $X, Y$  independent, then  $f(X), g(Y)$  independent for all  $f, g$ .

Expectation:  $E[X] = \sum_t t \Pr[X = t]$

Tail sum: for nonnegative r.v.  $X$ :  $E[X] = \sum_{i=0}^{\infty} \Pr[X > i]$ .

Expectation of function:  $E[g(X)] = \sum_t g(t) \Pr[X = t]$

Variance:  $\text{Var}[X] = E[(X - E[X])^2] = E[X^2] - E[X]^2$

Standard deviation: square root of variance.

Linearity of expectation:  $E[aX + bY] = aE[X] + bE[Y]$



# Random Variables: Discrete

Random variable: function that assigns a real number  $X(\omega)$  to each outcome  $\omega$  in a probability space.

Random variables  $X, Y$  are independent if the events  $Y = a$  and  $X = b$  are independent for all  $a, b$ . If  $X, Y$  independent, then  $f(X), g(Y)$  independent for all  $f, g$ .

Expectation:  $E[X] = \sum_t t \Pr[X = t]$

Tail sum: for nonnegative r.v.  $X$ :  $E[X] = \sum_{i=0}^{\infty} \Pr[X > i]$ .

Expectation of function:  $E[g(X)] = \sum_t g(t) \Pr[X = t]$

Variance:  $\text{Var}[X] = E[(X - E[X])^2] = E[X^2] - E[X]^2$

Standard deviation: square root of variance.

Linearity of expectation:  $E[aX + bY] = aE[X] + bE[Y]$

For independent RV:  $E[XY] = E[X]E[Y]$ ,  $\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y]$

## Example: Random-SAT

Let's say I have some Boolean clause that looks like this ("3-CNF")

$$(a \vee b \vee \bar{c}) \wedge (\bar{b} \wedge d \wedge e) \wedge \dots$$

$n$  clauses (three boolean variables, some may be negated). What is expected number of clauses that I satisfy with a random assignment?

## Example: Random-SAT

Let's say I have some Boolean clause that looks like this ("3-CNF")

$$(a \vee b \vee \bar{c}) \wedge (\bar{b} \wedge d \wedge e) \wedge \dots$$

$n$  clauses (three boolean variables, some may be negated). What is expected number of clauses that I satisfy with a random assignment?  $7n/8$ .

## Example: Random-SAT

Let's say I have some Boolean clause that looks like this ("3-CNF")

$$(a \vee b \vee \bar{c}) \wedge (\bar{b} \wedge d \wedge e) \wedge \dots$$

$n$  clauses (three boolean variables, some may be negated). What is expected number of clauses that I satisfy with a random assignment?  $7n/8$ .

Doesn't matter if variables are repeated! Expectation is linear.

## Example: Random-SAT

Let's say I have some Boolean clause that looks like this ("3-CNF")

$$(a \vee b \vee \bar{c}) \wedge (\bar{b} \wedge d \wedge e) \wedge \dots$$

$n$  clauses (three boolean variables, some may be negated). What is expected number of clauses that I satisfy with a random assignment?  $7n/8$ .

Doesn't matter if variables are repeated! Expectation is linear.

Also proves (by probabilistic method) that there exists some assignment satisfying at least  $7/8$  of the clauses.

## Example: Random-SAT

Let's say I have some Boolean clause that looks like this ("3-CNF")

$$(a \vee b \vee \bar{c}) \wedge (\bar{b} \wedge d \wedge e) \wedge \dots$$

$n$  clauses (three boolean variables, some may be negated). What is expected number of clauses that I satisfy with a random assignment?  $7n/8$ .

Doesn't matter if variables are repeated! Expectation is linear.

Also proves (by probabilistic method) that there exists some assignment satisfying at least  $7/8$  of the clauses.

Turns out that we don't know any better constant-factor approximation for this.  $7/8$  is the best we can do! If we can efficiently do better (i.e.  $7/8 + \epsilon$  fraction of clauses satisfied, for constant  $\epsilon$ ) this would prove  $P = NP$  which would, among many other things, render public key cryptography impossible!

## Example: Random-SAT

Let's say I have some Boolean clause that looks like this ("3-CNF")

$$(a \vee b \vee \bar{c}) \wedge (\bar{b} \wedge d \wedge e) \wedge \dots$$

$n$  clauses (three boolean variables, some may be negated). What is expected number of clauses that I satisfy with a random assignment?  $7n/8$ .


Doesn't matter if variables are repeated! Expectation is linear.

Also proves (by probabilistic method) that there exists some assignment satisfying at least  $7/8$  of the clauses.

Turns out that we don't know any better constant-factor approximation for this.  $7/8$  is the best we can do! If we can efficiently do better (i.e.  $7/8 + \epsilon$  fraction of clauses satisfied, for constant  $\epsilon$ ) this would prove  $P = NP$  which would, among many other things, render public key cryptography impossible!

"Hardness of approximation". Ongoing topic of research.

# Random Variables: Continuous



Distributions represented with a pdf


$$f_X(t) = \lim_{\delta \rightarrow 0} \frac{\Pr[X \in [t, t + \delta]]}{\delta}$$

...or, equivalently, a cdf:

$$F_X(t) = \Pr[X \leq t] = \int_{-\infty}^t f_X(z) dz.$$



# Random Variables: Continuous



Distributions represented with a pdf

$$f_X(t) = \lim_{\delta \rightarrow 0} \frac{\Pr[X \in [t, t + \delta]]}{\delta}$$

...or, equivalently, a cdf:

$$F_X(t) = \Pr[X \leq t] = \int_{-\infty}^t f_X(z) dz.$$

$$\Pr[X \in [a, b]] = \int_a^b f_X(t) dt = F_X(b) - F_X(a)$$

# Expectation/Variance for Continuous

Sum  $\rightarrow$  Integral. Most properties carry over.

$$E[X] = \int_{-\infty}^{\infty} tf_X(t)dt$$

$$E[g(X)] = \int_{-\infty}^{\infty} g(t)f_X(t)dt$$

$$\text{Var}[X] = E[(X - E[X])^2] = E[X^2] - E[X]^2$$

Linearity of expectation:  $E[aX + bY] = aE[X] + bE[Y]$

For independent RV:  $E[XY] = E[X]E[Y]$ ,  $\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y]$

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

Keep the minimum number of all the visitors to your website (only requires space for one number!).

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

Keep the minimum number of all the visitors to your website (only requires space for one number!).

What's the number you get? Minimum of  $Uniform(0,1)$ .

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

Keep the minimum number of all the visitors to your website (only requires space for one number!).

What's the number you get? Minimum of  $Uniform(0,1)$ .Distribution?

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

Keep the minimum number of all the visitors to your website (only requires space for one number!).

What's the number you get? Minimum of  $Uniform(0,1)$ . Distribution?  
CDF:

$$\Pr(\min_i X_i \leq x) =$$



## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

Keep the minimum number of all the visitors to your website (only requires space for one number!).

What's the number you get? Minimum of  $Uniform(0,1)$ . Distribution?  
CDF:

$$\Pr(\min_i X_i \leq x) = 1 - \Pr(\text{all } x_i \text{ at least } x) =$$

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

Keep the minimum number of all the visitors to your website (only requires space for one number!).

What's the number you get? Minimum of  $Uniform(0,1)$ . Distribution?  
CDF:

$$\Pr(\min_i X_i \leq x) = 1 - \Pr(\text{all } x_i \text{ at least } x) = 1 - (1-x)^n$$

So PDF is  $f(x) = n(1-x)^{n-1}$ . Expectation:  $\int_0^1 xn(1-x)^{n-1}dx = 1/(n+1)$ .

## Application: Streaming Algorithm for Counting Uniques

Let's say that you're building a server that wants to count unique visitors. But you only have a very small amount of memory - enough to remember one number. How do you distinguish between a million unique visitors and a single IP address sending a million requests to your site?

Map each IP address to a single number between 0 and 1 uniformly.

Keep the minimum number of all the visitors to your website (only requires space for one number!).

What's the number you get? Minimum of  $Uniform(0,1)$ . Distribution?  
CDF:

$$\Pr(\min_i X_i \leq x) = 1 - \Pr(\text{all } x_i \text{ at least } x) = 1 - (1 - x)^n$$

So PDF is  $f(x) = n(1 - x)^{n-1}$ . Expectation:  $\int_0^1 xn(1 - x)^{n-1}dx = 1/(n + 1)$ .  
Just invert the minimum number to estimate number of unique visitors!

# Distributions

**Discrete:** Uniform, Bernoulli, geometric, binomial, Poisson

**Continuous:** Exponential, normal, uniform.

# Distributions

**Discrete:** Uniform, Bernoulli, geometric, binomial, Poisson

**Continuous:** Exponential, normal, uniform.

Make sure you know what they mean intuitively (although formula sheet will have the formulas for them).

# Distributions

**Discrete:** Uniform, Bernoulli, geometric, binomial, Poisson

**Continuous:** Exponential, normal, uniform.

Make sure you know what they mean intuitively (although formula sheet will have the formulas for them).

For instance: What's the distribution of the sum of two independent binomial random variables? What's the distribution of the minimum of two independent geometric random variables?

# Distributions

**Discrete:** Uniform, Bernoulli, geometric, binomial, Poisson

**Continuous:** Exponential, normal, uniform.

Make sure you know what they mean intuitively (although formula sheet will have the formulas for them).

For instance: What's the distribution of the sum of two independent binomial random variables? What's the distribution of the minimum of two independent geometric random variables? Prove these formally for practice!

# Tail Bounds

Markov: For  $X$  non-negative,  $a$  positive,

$$\Pr[X \geq a] \leq \frac{E[X]}{a}.$$



# Tail Bounds

Markov: For  $X$  non-negative,  $a$  positive,

$$\Pr[X \geq a] \leq \frac{E[X]}{a}.$$

Chebyshev: For all  $a$  positive,

$$\Pr[|X - E[X]| \geq a] \leq \frac{\text{Var}[X]}{a^2}.$$

# Tail Bounds

Markov: For  $X$  non-negative,  $a$  positive,

$$\Pr[X \geq a] \leq \frac{E[X]}{a}.$$

Chebyshev: For all  $a$  positive,

$$\Pr[|X - E[X]| \geq a] \leq \frac{\text{Var}[X]}{a^2}.$$

Chernoff: Family of exponential bounds for sum of mutually independent 0-1 random variables. Derive by noting that  $\Pr[X \geq a] = \Pr[e^{tX} \geq e^{ta}]$ , and then applying Markov to bound

$$\Pr[e^{tX} \geq e^{ta}] \leq \frac{E[e^{tX}]}{e^{ta}}$$

for a good value of  $t$ .

# Law of Large Numbers and CLT

If  $X_1, X_2, \dots$  are pairwise independent, and identically distributed with mean  $\mu$ :  $\Pr\left[\left|\frac{\sum_i X_i}{n} - \mu\right| \geq \epsilon\right] \rightarrow 0$  as  $n \rightarrow \infty$ .

# Law of Large Numbers and CLT

If  $X_1, X_2, \dots$  are pairwise independent, and identically distributed with mean  $\mu$ :  $\Pr\left[\left|\frac{\sum_i X_i}{n} - \mu\right| \geq \epsilon\right] \rightarrow 0$  as  $n \rightarrow \infty$ .

With many i.i.d. samples we converge not only to the mean, but also to a normal distribution with the same variance.

CLT: Suppose  $X_1, X_2, \dots$  are i.i.d. random variables with expectation  $\mu$  and variance  $\sigma^2$ . Let

$$S_n := \frac{(\sum_i X_i) - n\mu}{\sigma\sqrt{n}}$$

Then  $S_n$  tends towards  $\mathcal{N}(0,1)$  as  $n \rightarrow \infty$ .

# Law of Large Numbers and CLT

If  $X_1, X_2, \dots$  are pairwise independent, and identically distributed with mean  $\mu$ :  $\Pr\left[\left|\frac{\sum_i X_i}{n} - \mu\right| \geq \epsilon\right] \rightarrow 0$  as  $n \rightarrow \infty$ .

With many i.i.d. samples we converge not only to the mean, but also to a normal distribution with the same variance.

CLT: Suppose  $X_1, X_2, \dots$  are i.i.d. random variables with expectation  $\mu$  and variance  $\sigma^2$ . Let

$$S_n := \frac{(\sum_i X_i) - n\mu}{\sigma\sqrt{n}}$$

Then  $S_n$  tends towards  $\mathcal{N}(0,1)$  as  $n \rightarrow \infty$ .

Or:

$$\Pr[S_n \leq a] \rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx$$

# Law of Large Numbers and CLT

If  $X_1, X_2, \dots$  are pairwise independent, and identically distributed with mean  $\mu$ :  $\Pr\left[\left|\frac{\sum_i X_i}{n} - \mu\right| \geq \epsilon\right] \rightarrow 0$  as  $n \rightarrow \infty$ .

With many i.i.d. samples we converge not only to the mean, but also to a normal distribution with the same variance.

CLT: Suppose  $X_1, X_2, \dots$  are i.i.d. random variables with expectation  $\mu$  and variance  $\sigma^2$ . Let

$$S_n := \frac{(\sum_i X_i) - n\mu}{\sigma\sqrt{n}}$$

Then  $S_n$  tends towards  $\mathcal{N}(0,1)$  as  $n \rightarrow \infty$ .

Or:

$$\Pr[S_n \leq a] \rightarrow \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx$$

This is an approximation, not a bound.

Live Demo

Live Demo

Transition matrix  $P$ . Timesteps correspond to matrix multiplication:

$$\pi \rightarrow \pi P.$$



Live Demo

Transition matrix  $P$ . Timesteps correspond to matrix multiplication:

$$\pi \rightarrow \pi P.$$

**Hitting time:** How long does it take us to get to some state  $j$ ?

Strategy: let  $\beta(i)$  be the time it takes to get to  $j$  from  $i$ , for each state  $i$ .  $\beta(j) = 0$ . Set up system of linear equations and solve.

# State Classifications

State  $j$  is **accessible** from  $i$ : can get from  $i$  to  $j$  with nonzero probability. Equivalently: exists path from  $i$  to  $j$ .

# State Classifications

State  $j$  is **accessible** from  $i$ : can get from  $i$  to  $j$  with nonzero probability. Equivalently: exists path from  $i$  to  $j$ .

$i$  accessible from  $j$  and  $j$  accessible from  $i$ :  $i, j$  **communicate**.

# State Classifications

State  $j$  is **accessible** from  $i$ : can get from  $i$  to  $j$  with nonzero probability. Equivalently: exists path from  $i$  to  $j$ .

$i$  accessible from  $j$  and  $j$  accessible from  $i$ :  $i, j$  **communicate**.

If, given that we're at some state, we will see that state again sometime in the future with probability 1, state is **recurrent**. If there is a nonzero probability that we don't ever see state again, state is **transient**.

# State Classifications

State  $j$  is **accessible** from  $i$ : can get from  $i$  to  $j$  with nonzero probability. Equivalently: exists path from  $i$  to  $j$ .

$i$  accessible from  $j$  and  $j$  accessible from  $i$ :  $i, j$  **communicate**.

If, given that we're at some state, we will see that state again sometime in the future with probability 1, state is **recurrent**. If there is a nonzero probability that we don't ever see state again, state is **transient**.

Every finite chain has a recurrent state.

# State Classifications

State  $j$  is **accessible** from  $i$ : can get from  $i$  to  $j$  with nonzero probability. Equivalently: exists path from  $i$  to  $j$ .

$i$  accessible from  $j$  and  $j$  accessible from  $i$ :  $i, j$  **communicate**.

If, given that we're at some state, we will see that state again sometime in the future with probability 1, state is **recurrent**. If there is a nonzero probability that we don't ever see state again, state is **transient**.

Every finite chain has a recurrent state.

State is **periodic** if, given that we're currently at that state, the probability that we are at that state  $s$  steps later is zero unless  $s$  divides some integer  $\Delta > 1$ .

# State Classifications

State  $j$  is **accessible** from  $i$ : can get from  $i$  to  $j$  with nonzero probability. Equivalently: exists path from  $i$  to  $j$ .

$i$  accessible from  $j$  and  $j$  accessible from  $i$ :  $i, j$  **communicate**.

If, given that we're at some state, we will see that state again sometime in the future with probability 1, state is **recurrent**. If there is a nonzero probability that we don't ever see state again, state is **transient**.

Every finite chain has a recurrent state.

State is **periodic** if, given that we're currently at that state, the probability that we are at that state  $s$  steps later is zero unless  $s$  divides some integer  $\Delta > 1$ .

**Ergodic** state: aperiodic + recurrent.

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.



## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ .

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ . Then there is a nonzero probability that, starting at  $i$ , we will go to  $j$ , at which point we will never be able to see  $i$  again. So  $i$  is transient.

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ . Then there is a nonzero probability that, starting at  $i$ , we will go to  $j$ , at which point we will never be able to see  $i$  again. So  $i$  is transient.

On the other hand, suppose  $i$  is accessible from  $j$ .

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ . Then there is a nonzero probability that, starting at  $i$ , we will go to  $j$ , at which point we will never be able to see  $i$  again. So  $i$  is transient.

On the other hand, suppose  $i$  is accessible from  $j$ . Suppose for contradiction that  $i$  is recurrent.

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ . Then there is a nonzero probability that, starting at  $i$ , we will go to  $j$ , at which point we will never be able to see  $i$  again. So  $i$  is transient.

On the other hand, suppose  $i$  is accessible from  $j$ . Suppose for contradiction that  $i$  is recurrent. Then if we're at  $j$ , we *have* to hit  $i$  again (because  $i$  is recurrent, so we have to go back to  $i$  if we go from  $i$  to  $j$ ).

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ . Then there is a nonzero probability that, starting at  $i$ , we will go to  $j$ , at which point we will never be able to see  $i$  again. So  $i$  is transient.

On the other hand, suppose  $i$  is accessible from  $j$ . Suppose for contradiction that  $i$  is recurrent. Then if we're at  $j$ , we *have* to hit  $i$  again (because  $i$  is recurrent, so we have to go back to  $i$  if we go from  $i$  to  $j$ ). But when we're at  $i$ , we know that we're definitely going to hit  $j$  sometime (because there's a nonzero chance of going to  $j$  from  $i$ , and we'll be back at  $i$  infinitely many times due to it being recurrent).

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ . Then there is a nonzero probability that, starting at  $i$ , we will go to  $j$ , at which point we will never be able to see  $i$  again. So  $i$  is transient.

On the other hand, suppose  $i$  is accessible from  $j$ . Suppose for contradiction that  $i$  is recurrent. Then if we're at  $j$ , we *have* to hit  $i$  again (because  $i$  is recurrent, so we have to go back to  $i$  if we go from  $i$  to  $j$ ). But when we're at  $i$ , we know that we're definitely going to hit  $j$  sometime (because there's a nonzero chance of going to  $j$  from  $i$ , and we'll be back at  $i$  infinitely many times due to it being recurrent). So  $j$  is recurrent. Contradiction!

## Example: Markov Proof

Here's a theorem and a proof of the sort that we might ask you to do on the test.

**Theorem:** If a transient state  $j$  is accessible from state  $i$ , then state  $i$  is transient.

**Proof:** Suppose  $i$  is not accessible from  $j$ . Then there is a nonzero probability that, starting at  $i$ , we will go to  $j$ , at which point we will never be able to see  $i$  again. So  $i$  is transient.

On the other hand, suppose  $i$  is accessible from  $j$ . Suppose for contradiction that  $i$  is recurrent. Then if we're at  $j$ , we *have* to hit  $i$  again (because  $i$  is recurrent, so we have to go back to  $i$  if we go from  $i$  to  $j$ ). But when we're at  $i$ , we know that we're definitely going to hit  $j$  sometime (because there's a nonzero chance of going to  $j$  from  $i$ , and we'll be back at  $i$  infinitely many times due to it being recurrent). So  $j$  is recurrent. Contradiction! So  $i$  has to be transient.



**Irreducible** Markov chain: all states communicate with every other state. Equivalently: graph representation is strongly connected.

# Markov Chain Classifications

**Irreducible** Markov chain: all states communicate with every other state. Equivalently: graph representation is strongly connected.

**Periodic** Markov chain: any state is periodic.

# Markov Chain Classifications

**Irreducible** Markov chain: all states communicate with every other state. Equivalently: graph representation is strongly connected.

**Periodic** Markov chain: any state is periodic.

**Ergodic** Markov chain: every state is ergodic. Any finite, irreducible, aperiodic Markov chain is ergodic.

# Stationary Distributions

Distribution is unchanged by state. Intuitively: if I have a lot (approaching infinity) of people on the same MC: the number of people at each state is constant (even if the individual people may move around).

# Stationary Distributions

Distribution is unchanged by state. Intuitively: if I have a lot (approaching infinity) of people on the same MC: the number of people at each state is constant (even if the individual people may move around).

To find limiting distribution? Solve **balance equations**:  $\pi = \pi P$ .

Let  $r_{i,j}^t$  be the probability that we first (if  $i = j$ , we don't count the zeroth timestep) hit  $j$  exactly  $t$  timesteps after we start at  $i$ . Then 
$$h_{i,j} = \sum_{t \geq 1} tr_{i,j}^t.$$

# Stationary Distributions

Distribution is unchanged by state. Intuitively: if I have a lot (approaching infinity) of people on the same MC: the number of people at each state is constant (even if the individual people may move around).

To find limiting distribution? Solve **balance equations**:  $\pi = \pi P$ .

Let  $r_{i,j}^t$  be the probability that we first (if  $i = j$ , we don't count the zeroth timestep) hit  $j$  exactly  $t$  timesteps after we start at  $i$ . Then  $h_{i,j} = \sum_{t \geq 1} tr_{i,j}^t$ .

Suppose we are given a finite, irreducible, aperiodic Markov chain. Then:

- There is a unique stationary distribution  $\pi$ .
- For all  $j, i$ , the limit  $\lim_{t \rightarrow \infty} P_{j,i}^t$  exists and is independent of  $j$ .
- $\pi_i = \lim_{t \rightarrow \infty} P_{j,i}^t = 1/h_{i,i}$

Markov chain on an undirected graph. At a vertex, pick edge with uniform probability and walk down it.

Markov chain on an undirected graph. At a vertex, pick edge with uniform probability and walk down it.

For undirected graphs: aperiodic if and only if graph is not bipartite.



Markov chain on an undirected graph. At a vertex, pick edge with uniform probability and walk down it.

For undirected graphs: aperiodic if and only if graph is not bipartite.

Stationary distribution:  $\pi_v = d(v)/(2|E|)$ .

Markov chain on an undirected graph. At a vertex, pick edge with uniform probability and walk down it.

For undirected graphs: aperiodic if and only if graph is not bipartite.

Stationary distribution:  $\pi_v = d(v)/(2|E|)$ .

Cover time (expected time that it takes to hit all the vertices, starting from the worst vertex possible): bounded above by  $4|V||E|$ .

## Example/Gig: Parrondo's Paradox

Let's say I have two slot machines. Each one takes some amount of money and then spits out some amount of money.

## Example/Gig: Parrondo's Paradox

Let's say I have two slot machines. Each one takes some amount of money and then spits out some amount of money.

Suppose that the expected return of each machine is negative - I get less money than I put in... the house always wins, after all. If I play machine 1 for a while, I expect to end up broke. Same with machine 2.

## Example/Gig: Parrondo's Paradox

Let's say I have two slot machines. Each one takes some amount of money and then spits out some amount of money.

Suppose that the expected return of each machine is negative - I get less money than I put in... the house always wins, after all. If I play machine 1 for a while, I expect to end up broke. Same with machine 2.

So if I play machine 1 and machine 2 alternately, I should expect to end up broke too, right? Hmm...

## Parrondo's Paradox II

Let's say that the slot machines work as follows:

Machine 1: Put in some money. You gain a dollar w.p. 0.49 and lose a dollar w.p. 0.51. Pretty obvious that you lose money playing this game.

## Parrondo's Paradox II

Let's say that the slot machines work as follows:

Machine 1: Put in some money. You gain a dollar w.p. 0.49 and lose a dollar w.p. 0.51. Pretty obvious that you lose money playing this game.

Machine 2: You put in  $d$  dollars.

- Case A: If  $d$  is a multiple of 3 then you gain a dollar w.p. 0.09 and lose a dollar w.p. 0.91.
- Case B: Otherwise, you gain a dollar w.p. 0.74 and lose a dollar w.p. 0.26.

What's the probability of winning a round?

## Parrondo's Paradox II

Let's say that the slot machines work as follows:

Machine 1: Put in some money. You gain a dollar w.p. 0.49 and lose a dollar w.p. 0.51. Pretty obvious that you lose money playing this game.

Machine 2: You put in  $d$  dollars.

- Case A: If  $d$  is a multiple of 3 then you gain a dollar w.p. 0.09 and lose a dollar w.p. 0.91.
- Case B: Otherwise, you gain a dollar w.p. 0.74 and lose a dollar w.p. 0.26.

What's the probability of winning a round?  $1/3$  probability of case A happening, so it would be

$$\frac{1}{3}(0.09) + \frac{2}{3}(0.74) = \frac{157}{300} > \frac{1}{2}$$

right?



## Parrondo's Paradox II

Let's say that the slot machines work as follows:

Machine 1: Put in some money. You gain a dollar w.p. 0.49 and lose a dollar w.p. 0.51. Pretty obvious that you lose money playing this game.

Machine 2: You put in  $d$  dollars.

- Case A: If  $d$  is a multiple of 3 then you gain a dollar w.p. 0.09 and lose a dollar w.p. 0.91.
- Case B: Otherwise, you gain a dollar w.p. 0.74 and lose a dollar w.p. 0.26.

What's the probability of winning a round?  $1/3$  probability of case A happening, so it would be

$$\frac{1}{3}(0.09) + \frac{2}{3}(0.74) = \frac{157}{300} > \frac{1}{2}$$

right? Are you sure?

# Parrondo's Paradox II

Let's say that the slot machines work as follows:

Machine 1: Put in some money. You gain a dollar w.p. 0.49 and lose a dollar w.p. 0.51. Pretty obvious that you lose money playing this game.

Machine 2: You put in  $d$  dollars.

- Case A: If  $d$  is a multiple of 3 then you gain a dollar w.p. 0.09 and lose a dollar w.p. 0.91.
- Case B: Otherwise, you gain a dollar w.p. 0.74 and lose a dollar w.p. 0.26.

What's the probability of winning a round?  $1/3$  probability of case A happening, so it would be

$$\frac{1}{3}(0.09) + \frac{2}{3}(0.74) = \frac{157}{300} > \frac{1}{2}$$

right? Are you sure? No!

# Parrondo's Paradox II

Let's say that the slot machines work as follows:

Machine 1: Put in some money. You gain a dollar w.p. 0.49 and lose a dollar w.p. 0.51. Pretty obvious that you lose money playing this game.

Machine 2: You put in  $d$  dollars.

- Case A: If  $d$  is a multiple of 3 then you gain a dollar w.p. 0.09 and lose a dollar w.p. 0.91.
- Case B: Otherwise, you gain a dollar w.p. 0.74 and lose a dollar w.p. 0.26.


What's the probability of winning a round?  $1/3$  probability of case A happening, so it would be

$$\frac{1}{3}(0.09) + \frac{2}{3}(0.74) = \frac{157}{300} > \frac{1}{2}$$

right? Are you sure? **No!** Probability of case A happening is not  $1/3$ ! (be careful about nonuniform probability spaces. MT2 1.1/1.2!)

## Parrondo's Paradox III


So how often do we end up with case A? Here's the approach: one state for each value of  $d \pmod 3$ .



Aperiodic? Irreducible? Yep!

## Parrondo's Paradox III


So how often do we end up with case A? Here's the approach: one state for each value of  $d \pmod 3$ .



Aperiodic? Irreducible? Yep! Limiting distribution = stationary distribution!

## Parrondo's Paradox III


So how often do we end up with case A? Here's the approach: one state for each value of  $d \pmod 3$ .



Aperiodic? Irreducible? Yep! Limiting distribution = stationary distribution! Just solve for the stationary distribution with  $\pi = \pi P$ .

## Parrondo's Paradox III

So how often do we end up with case A? Here's the approach: one state for each value of  $d \pmod 3$ .




Aperiodic? Irreducible? Yep! Limiting distribution = stationary distribution! Just solve for the stationary distribution with  $\pi = \pi P$ .

Result:  $\pi = [0.382604, 0.154728, 0.462668]$ .

## Parrondo's Paradox III

So how often do we end up with case A? Here's the approach: one state for each value of  $d \pmod 3$ .



Aperiodic? Irreducible? Yep! Limiting distribution = stationary distribution! Just solve for the stationary distribution with  $\pi = \pi P$ .

Result:  $\pi = [0.382604, 0.154728, 0.462668]$ . Plug in:

$$0.3826(0.09) + (0.1547 + 0.4627)(0.74) = 0.4913 < \frac{1}{2}$$

So I lose money in the long run.



## Parrondo's Paradox III

So now, what if I decide to flip a fair coin to figure out which machine to play?

## Parrondo's Paradox III

So now, what if I decide to flip a fair coin to figure out which machine to play?

I have  $d$  dollars... if  $d$  is a multiple of 3, probability of winning is:

$$\frac{1}{2}(0.49) + \frac{1}{2}(0.09) = 0.29$$

If  $d$  isn't a multiple of 3, probability of winning is:

$$\frac{1}{2}(0.49) + \frac{1}{2}(0.74) = 0.615$$

# Parrondo's Paradox III


So now, what if I decide to flip a fair coin to figure out which machine to play?

I have  $d$  dollars... if  $d$  is a multiple of 3, probability of winning is:

$$\frac{1}{2}(0.49) + \frac{1}{2}(0.09) = 0.29$$

If  $d$  isn't a multiple of 3, probability of winning is:

$$\frac{1}{2}(0.49) + \frac{1}{2}(0.74) = 0.615$$



Stationary distribution:  $\pi = [0.344583, 0.254343, 0.401075]$ .

## Parrondo's Paradox IV

Stationary distribution:  $\pi = [0.344583, 0.254343, 0.401075]$ .

Probability of winning:

$$0.3446(0.29) + (0.2543 + 0.4011)(0.615) = 0.503011 > \frac{1}{2}$$

## Parrondo's Paradox IV

Stationary distribution:  $\pi = [0.344583, 0.254343, 0.401075]$ .

Probability of winning:

$$0.3446(0.29) + (0.2543 + 0.4011)(0.615) = 0.503011 > \frac{1}{2}$$

So we expect to... gain money??!?!?!?!?!?

## Parrondo's Paradox IV

Stationary distribution:  $\pi = [0.344583, 0.254343, 0.401075]$ .

Probability of winning:

$$0.3446(0.29) + (0.2543 + 0.4011)(0.615) = 0.503011 > \frac{1}{2}$$

So we expect to... gain money??!?!?!?!?!?

Did we just break linearity of expectation?

## Parrondo's Paradox IV

Stationary distribution:  $\pi = [0.344583, 0.254343, 0.401075]$ .

Probability of winning:

$$0.3446(0.29) + (0.2543 + 0.4011)(0.615) = 0.503011 > \frac{1}{2}$$

So we expect to... gain money??!?!?!?!?!?

Did we just break linearity of expectation? No! It doesn't make a whole lot of sense to talk about "expected winnings" for a state without taking into account the current state. Our distribution across states changes between the two games!



Questions?