Continuous Probability

CS70 Summer 2016 - Lecture 6A

David Dinh 25 July 2016

UC Berkeley

Logistics

Tutoring Sections - M/W 5-8PM in 540 Cory.

- Conceptual discussions of material
- No homework discussion (take that to OH/HW party, please)

Midterm is this Friday - 11:30-1:30, same rooms as last time.

- · Covers material from MT1 to this Wednesday...
- ...but we will expect you to know everything we've covered from the start of class.
- One **double**-sided sheet of notes allowed (our advice: reuse sheet from MT1 and add MT2 topics to the other side).
- Students with time conflicts and DSP students should have been contacted by us - if you are one and you haven't heard from us, get in touch ASAP.

Today

- What is continuous probability?
- · Expectation and variance in the continuous setting.
- · Some common distributions.

Continuous Probability

Sometimes you can't model things discretely.

Sometimes you can't model things discretely. Random real numbers.

Sometimes you can't model things discretely. Random real numbers. Points on a map.

Sometimes you can't model things discretely. Random real numbers. Points on a map. Time.

Sometimes you can't model things discretely. Random real numbers. Points on a map. Time.

Probability space is **continuous**.

Sometimes you can't model things discretely. Random real numbers. Points on a map. Time.

Probability space is continuous.

What is probability? Function mapping events to [0, 1].

Sometimes you can't model things discretely. Random real numbers. Points on a map. Time.

Probability space is continuous.

What is probability? Function mapping events to [0, 1].

What is an event in continuous probability?

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here?

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32?

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32? Sample space: all times between 14:00 and 14:10.

Z

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32?

Sample space: all times between 14:00 and 14:10.

Size of sample space?

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32?

Sample space: all times between 14:00 and 14:10.

Size of sample space? How many numbers are there between 0 and 10?

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32? Sample space: all times between 14:00 and 14:10.

Size of sample space? How many numbers are there between 0 and 10? infinite

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32?

Sample space: all times between 14:00 and 14:10.

Size of sample space? How many numbers are there between 0 and 10? infinite

Chance of getting one event in an infinite sized uniform sample space?

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32?

Sample space: all times between 14:00 and 14:10.

Size of sample space? How many numbers are there between 0 and 10? infinite

Chance of getting one event in an infinite sized uniform sample space? ${\color{blue}0}$

Class starts at 14:10. You take your seat at some "uniform" random time between 14:00 and 14:10.

What's an event here? Probability of coming in at exactly 14:03:47.32?

Sample space: all times between 14:00 and 14:10.

Size of sample space? How many numbers are there between 0 and 10? infinite

Chance of getting one event in an infinite sized uniform sample space? 0

Not so simple to define events in continuous probability!



Look at intervals instead of specific times.



Look at *intervals* instead of specific times.

Probability that you come in between 14:00 and 14:10?




Look at intervals instead of specific times.

Probability that you come in between 14:00 and 14:10? 1.




- Look at *intervals* instead of specific times.
- Probability that you come in between 14:00 and 14:10? 1.
- Probability that you come in between 14:00 and 14:05?




Look at intervals instead of specific times.

Probability that you come in between 14:00 and 14:10? 1.

Probability that you come in between 14:00 and 14:05? 1/2.



- Look at intervals instead of specific times.
- Probability that you come in between 14:00 and 14:10? 1.
- Probability that you come in between 14:00 and 14:05? 1/2.
- Probability that you come between 14:03 and 14:04?




Look at intervals instead of specific times.

Probability that you come in between 14:00 and 14:10? 1.

Probability that you come in between 14:00 and 14:05? 1/2.

Probability that you come between 14:03 and 14:04? 1/10.




Look at intervals instead of specific times.

Probability that you come in between 14:00 and 14:10? 1.

Probability that you come in between 14:00 and 14:05? 1/2.

Probability that you come between 14:03 and 14:04? 1/10.

Probability that you come in some time interval of 10/k minutes?



Look at intervals instead of specific times.

Probability that you come in between 14:00 and 14:10? 1.

Probability that you come in between 14:00 and 14:05? 1/2.

Probability that you come between 14:03 and 14:04? 1/10.

Probability that you come in some time interval of 10/k minutes? 1/k.


What happens when you take $k \to \infty$? Probability goes to 0.

What happens when you take $k \to \infty$? Probability goes to 0.


What happens when you take $k \to \infty$? Probability goes to 0.




What happens when you take $k \to \infty$? Probability goes to 0.



What happens when you take $k \to \infty$? Probability goes to 0.




What happens when you take $k \to \infty$? Probability goes to 0.



PDF (no, not the file format)

What happens when you take $k \to \infty$? Probability goes to 0.


What do we do so that this doesn't disappear? If we split our sample space into k pieces - multiply each one by k.



PDF (no, not the file format)

What happens when you take $k \to \infty$? Probability goes to 0.

What do we do so that this doesn't disappear? If we split our sample space into k pieces - multiply each one by k.



The resulting curve as $k \to \infty$ is the **probability density function** (PDF).

PDF $f_X(t)$ of a random variable X is defined so that the probability of X taking on a value in $[t, t + \delta]$ is $\delta f(t)$ for infinitesimally small δ .

PDF $f_X(t)$ of a random variable X is defined so that the probability of X taking on a value in $[t, t + \delta]$ is $\delta f(t)$ for infinitesimally small δ .

$$f_X(t) = \lim_{\delta \to 0} \frac{\Pr[X \in [t, t + \delta]]}{\delta}$$

PDF $f_X(t)$ of a random variable X is defined so that the probability of X taking on a value in $[t, t + \delta]$ is $\delta f(t)$ for infinitesimally small δ .

$$f_X(t) = \lim_{\delta \to 0} \frac{\Pr[X \in [t, t + \delta]]}{\delta}$$

Another way of looking at it:

$$\Pr[X \in [a, b]] = \int_a^b f_X(t) dt$$

PDF $f_X(t)$ of a random variable X is defined so that the probability of X taking on a value in $[t, t + \delta]$ is $\delta f(t)$ for infinitesimally small δ .

$$f_X(t) = \lim_{\delta \to 0} \frac{\Pr[X \in [t, t + \delta]]}{\delta}$$

Another way of looking at it:

$$\Pr[X \in [a, b]] = \int_a^b f_X(t)dt$$

f is nonnegative (negative probability doesn't make much sense).

PDF $f_X(t)$ of a random variable X is defined so that the probability of X taking on a value in $[t, t + \delta]$ is $\delta f(t)$ for infinitesimally small δ .

$$f_X(t) = \lim_{\delta \to 0} \frac{\Pr[X \in [t, t + \delta]]}{\delta}$$

Another way of looking at it:

$$\Pr[X \in [a, b]] = \int_a^b f_X(t)dt$$

f is nonnegative (negative probability doesn't make much sense).

Total probability is 1: $\int_{-\infty}^{\infty} f_X(t) dt = 1$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

$$F_X(t) = \int_{-\infty}^t F_X(z) dz$$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

$$F_X(t) = \int_{-\infty}^t F_X(z) dz$$

$$Pr[X \in (a, b]] =$$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Or, in terms of PDF...

$$F_X(t) = \int_{-\infty}^t F_X(z) dz$$

$$Pr[X \in (a, b]] = Pr[X \le b] - Pr[X \le a]$$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Or, in terms of PDF...

$$F_X(t) = \int_{-\infty}^t F_X(z) dz$$

$$\Pr[X \in (a, b]] = \Pr[X \le b] - \Pr[X \le a]$$

= $F_X(b) - F_X(a)$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

$$F_X(t) = \int_{-\infty}^t F_X(z) dz$$

$$\Pr[X \in (a, b]] = \Pr[X \le b] - \Pr[X \le a]$$

= $F_X(b) - F_X(a)$

$$F_X(t) \in [0,1]$$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Or, in terms of PDF...

$$F_X(t) = \int_{-\infty}^t F_X(z)dz$$

$$Pr[X \in (a, b]] = Pr[X \le b] - Pr[X \le a]$$

$$= F_X(b) - F_X(a)$$

$$F_X(t) \in [0, 1]$$

$$\lim_{t \to -\infty} F_X(t) =$$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Or, in terms of PDF...

$$F_X(t) = \int_{-\infty}^t F_X(z)dz$$

$$Pr[X \in (a, b]] = Pr[X \le b] - Pr[X \le a]$$

$$= F_X(b) - F_X(a)$$

$$F_X(t) \in [0, 1]$$

$$\lim_{t \to -\infty} F_X(t) = 0$$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Or, in terms of PDF...

$$F_X(t) = \int_{-\infty}^t F_X(z) dz$$

$$Pr[X \in (a, b]] = Pr[X \le b] - Pr[X \le a]$$
$$= F_X(b) - F_X(a)$$

$$F_X(t) \in [0,1]$$

$$\lim_{t\to-\infty}F_X(t)=0$$

$$\lim_{t\to\infty}F_X(t)=$$

Cumulative distribution function (CDF): $F_X(t) = Pr[X \le t]$.

Or, in terms of PDF...

$$F_X(t) = \int_{-\infty}^t F_X(z) dz$$


$$Pr[X \in (a, b]] = Pr[X \le b] - Pr[X \le a]$$
$$= F_X(b) - F_X(a)$$

$$F_X(t) \in [0,1]$$

$$\lim_{t\to-\infty}F_X(t)=0$$

$$\lim_{t\to\infty}F_X(t)=1$$

In Pictures



Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case?

Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case? Sum \rightarrow integral.

$$E[X] = \int_{-\infty}^{\infty} t f_X(t) dt$$

Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case? Sum \rightarrow integral.

$$E[X] = \int_{-\infty}^{\infty} t f_X(t) dt$$

Expectation of a function?

$$E[g(X)] = \int_{-\infty}^{\infty} g(t)f_X(t)dt$$

Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case? Sum \rightarrow integral.

$$E[X] = \int_{-\infty}^{\infty} t f_X(t) dt$$

Expectation of a function?

$$E[g(X)] = \int_{-\infty}^{\infty} g(t) f_X(t) dt$$

Linearity of expectation:

$$E[aX + bY] = aE[X] + bE[Y]$$

Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case? Sum \rightarrow integral.

$$E[X] = \int_{-\infty}^{\infty} t f_X(t) dt$$

Expectation of a function?

$$E[g(X)] = \int_{-\infty}^{\infty} g(t) f_X(t) dt$$

Linearity of expectation:

$$E[aX + bY] = aE[X] + bE[Y]$$

Proof: similar to discrete case.

Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case? Sum \rightarrow integral.

$$E[X] = \int_{-\infty}^{\infty} t f_X(t) dt$$

Expectation of a function?

$$E[g(X)] = \int_{-\infty}^{\infty} g(t) f_X(t) dt$$

Linearity of expectation:

$$E[aX + bY] = aE[X] + bE[Y]$$

Proof: similar to discrete case.

If X, Y, Z are mutually independent, then E[XYZ] = E[X]E[Y]E[Z].

Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case? Sum \rightarrow integral.

$$E[X] = \int_{-\infty}^{\infty} t f_X(t) dt$$

Expectation of a function?

$$E[g(X)] = \int_{-\infty}^{\infty} g(t) f_X(t) dt$$

Linearity of expectation:

$$E[aX + bY] = aE[X] + bE[Y]$$

Proof: similar to discrete case.

If X, Y, Z are mutually independent, then E[XYZ] = E[X]E[Y]E[Z].

Proof: also similar to discrete case.

Discrete case: $E[X] = \sum_{t=-\infty}^{\infty} (\Pr[X=t]t)$

Continuous case? Sum \rightarrow integral.

$$E[X] = \int_{-\infty}^{\infty} t f_X(t) dt$$

Expectation of a function?

$$E[g(X)] = \int_{-\infty}^{\infty} g(t) f_X(t) dt$$

Linearity of expectation:

$$E[aX + bY] = aE[X] + bE[Y]$$

Proof: similar to discrete case.

If X, Y, Z are mutually independent, then E[XYZ] = E[X]E[Y]E[Z].

Proof: also similar to discrete case.

Exercise: try proving these yourself.

Variance

Variance is defined exactly like it is for the discrete case.

$$Var(X) = E[(X - E[X])^{2}]$$

= $E[X^{2}] - E[X]^{2}$

Variance

Variance is defined exactly like it is for the discrete case.

$$Var(X) = E[(X - E[X])^{2}]$$

= $E[X^{2}] - E[X]^{2}$

The standard properties for variance hold in the continuous case as well.

$$Var(aX) = a^2 Var(X)$$

For independent r.v. X, Y:

$$Var(X + Y) = Var(X) + Var(Y)$$

.

Suppose an archer always hits a circular target with 1 meter radius, and the exact point that he hits is distributed uniformly across the target. What is the distribution the distance between his arrow and the center (call this r.v. X)?

Suppose an archer always hits a circular target with 1 meter radius, and the exact point that he hits is distributed uniformly across the target. What is the distribution the distance between his arrow and the center (call this r.v. X)?



Suppose an archer always hits a circular target with 1 meter radius, and the exact point that he hits is distributed uniformly across the target. What is the distribution the distance between his arrow and the center (call this r.v. X)?



Suppose an archer always hits a circular target with 1 meter radius, and the exact point that he hits is distributed uniformly across the target. What is the distribution the distance between his arrow and the center (call this r.v. X)?



$$Pr[X \le t] = \frac{\text{area of small circle}}{\text{area of dartboard}}$$

Suppose an archer always hits a circular target with 1 meter radius, and the exact point that he hits is distributed uniformly across the target. What is the distribution the distance between his arrow and the center (call this r.v. X)?



$$Pr[X \le t] = \frac{\text{area of small circle}}{\text{area of dartboard}}$$

= $\frac{\pi t^2}{\pi}$

Suppose an archer always hits a circular target with 1 meter radius, and the exact point that he hits is distributed uniformly across the target. What is the distribution the distance between his arrow and the center (call this r.v. X)?



$$Pr[X \le t] = \frac{\text{area of small circle}}{\text{area of dartboard}}$$

= $\frac{\pi t^2}{\pi} = t^2$.

CDF:

$$F_{Y}(t) = Pr[Y \le t] = \begin{cases} 0 & \text{for } t < 0 \\ t^{2} & \text{for } 0 \le t \le 1 \\ 1 & \text{for } t > 1 \end{cases}$$

PDF?

$$f_Y(t) = F_Y(t)' =$$

CDF:

$$F_{Y}(t) = Pr[Y \le t] = \begin{cases} 0 & \text{for } t < 0 \\ t^{2} & \text{for } 0 \le t \le 1 \\ 1 & \text{for } t > 1 \end{cases}$$

PDF?

$$f_Y(t) = F_Y(t)' = \begin{cases} 2t & \text{for } 0 \le t \le 1\\ 0 & \text{otherwise} \end{cases}$$

Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t+\delta$ for small δ ?



Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t+\delta$ for small δ ?



Area of circle: π

Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t+\delta$ for small δ ?



Area of circle: π

$$\pi((t+\delta)^2-t^2)$$

Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t + \delta$ for small δ ?



Area of circle: π

$$\pi((t+\delta)^2 - t^2) = \pi(t^2 + 2t\delta + \delta^2 - t^2)$$

Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t + \delta$ for small δ ?



Area of circle: π

$$\pi((t+\delta)^2 - t^2) = \pi(t^2 + 2t\delta + \delta^2 - t^2) = \pi(2t\delta + \delta^2)$$

Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t + \delta$ for small δ ?



Area of circle: π

$$\pi((t+\delta)^2 - t^2) = \pi(t^2 + 2t\delta + \delta^2 - t^2) = \pi(2t\delta + \delta^2) \approx \pi 2t\delta$$

Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t + \delta$ for small δ ?



Area of circle: π

Area of ring:

$$\pi((t+\delta)^2 - t^2) = \pi(t^2 + 2t\delta + \delta^2 - t^2) = \pi(2t\delta + \delta^2) \approx \pi 2t\delta$$

Probability of hitting the ring: $2t\delta$.

Another way of attacking the same problem: what's the probability of hitting some ring with inner radius t and outer radius $t + \delta$ for small δ ?



Area of circle: π

Area of ring:

$$\pi((t+\delta)^2 - t^2) = \pi(t^2 + 2t\delta + \delta^2 - t^2) = \pi(2t\delta + \delta^2) \approx \pi 2t\delta$$

Probability of hitting the ring: $2t\delta$.

PDF for $t \le 1$: 2t

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$
$$= Pr[X \in (\frac{y - a}{b}, \frac{y + \delta - a}{b})]$$

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y + \delta - a}{b})]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y - a}{b} + \frac{\delta}{b})]$$

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y + \delta - a}{b})]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y - a}{b} + \frac{\delta}{b})]$$

$$= f_X(\frac{y - a}{b})\frac{\delta}{b}.$$

Let $f_X(x)$ be the pdf of X and Y = a + bX where b > 0. Then

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y + \delta - a}{b})]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y - a}{b} + \frac{\delta}{b})]$$

$$= f_X(\frac{y - a}{b})\frac{\delta}{b}.$$

Left-hand side is

Let $f_X(x)$ be the pdf of X and Y = a + bX where b > 0. Then

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y + \delta - a}{b})]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y - a}{b} + \frac{\delta}{b})]$$

$$= f_X(\frac{y - a}{b})\frac{\delta}{b}.$$

Left-hand side is $f_Y(y)\delta$.

Let $f_X(x)$ be the pdf of X and Y = a + bX where b > 0. Then

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y + \delta - a}{b})]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y - a}{b} + \frac{\delta}{b})]$$

$$= f_X(\frac{y - a}{b})\frac{\delta}{b}.$$

Left-hand side is $f_Y(y)\delta$. Hence,

$$f_Y(y) = \frac{1}{b} f_X(\frac{y-a}{b}).$$

Let $f_X(x)$ be the pdf of X and Y = a + bX where b > 0. Then

$$Pr[Y \in (y, y + \delta)] = Pr[a + bX \in (y, y + \delta)]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y + \delta - a}{b})]$$

$$= Pr[X \in (\frac{y - a}{b}, \frac{y - a}{b} + \frac{\delta}{b})]$$

$$= f_X(\frac{y - a}{b})\frac{\delta}{b}.$$

Left-hand side is $f_Y(y)\delta$. Hence,

$$f_Y(y) = \frac{1}{b} f_X(\frac{y-a}{b}).$$

Continuous Distributions

Uniform Distribution: CDF and PDF

PDF is constant over some interval [a, b], zero outside the interval.

What's the value of the constant in the interval?

$$\int_{-\infty}^{\infty} k dt = \int_{a}^{b} k dt = b - a = 1$$

so PDF is 1/(b-a) in [a,b] and 0 otherwise.

Uniform Distribution: CDF and PDF

PDF is constant over some interval [a, b], zero outside the interval.

What's the value of the constant in the interval?

$$\int_{-\infty}^{\infty} k dt = \int_{a}^{b} k dt = b - a = 1$$

so PDF is 1/(b-a) in [a,b] and 0 otherwise.


CDF?

$$\int_{-\infty}^{t} 1/(b-a)dz$$

0 for t < a, (t - a)/(b - a) for a < t < b, and 1 for t > b.

Uniform Distribution: CDF and PDF, Graphically

$$f_X(t) = \begin{cases} 1/(b-a) & a < t < b \\ 0 & \text{otherwise} \end{cases} \qquad F_X(t) = \begin{cases} 0 & t < a \\ (t-a)/(b-a) & a < t < b \\ 1 & b < t \end{cases}$$



Expectation?

Expectation?

$$E[X] = \int_{a}^{b} \frac{t}{b-a} dt = \frac{1}{2} \frac{b^{2}-a^{2}}{b-a} = \frac{b+a}{2}$$

Expectation?

$$E[X] = \int_{a}^{b} \frac{t}{b-a} dt = \frac{1}{2} \frac{b^{2} - a^{2}}{b-a} = \frac{b+a}{2}$$

$$Var[X] = E[X^2] - E[X]^2$$

Expectation?

$$E[X] = \int_{a}^{b} \frac{t}{b-a} dt = \frac{1}{2} \frac{b^{2} - a^{2}}{b-a} = \frac{b+a}{2}$$

$$Var[X] = E[X^2] - E[X]^2$$
$$= \int_a^b \frac{t^2}{b-a} dt - \left(\frac{b+a}{2}\right)^2$$

Expectation?

$$E[X] = \int_{a}^{b} \frac{t}{b-a} dt = \frac{1}{2} \frac{b^{2} - a^{2}}{b-a} = \frac{b+a}{2}$$

$$Var[X] = E[X^{2}] - E[X]^{2}$$

$$= \int_{a}^{b} \frac{t^{2}}{b-a} dt - \left(\frac{b+a}{2}\right)^{2}$$

$$= \frac{t^{3}}{3(b-a)} \Big|_{a}^{b} - \left(\frac{b+a}{2}\right)^{2}$$

Expectation?

$$E[X] = \int_{a}^{b} \frac{t}{b-a} dt = \frac{1}{2} \frac{b^{2} - a^{2}}{b-a} = \frac{b+a}{2}$$

$$Var[X] = E[X^{2}] - E[X]^{2}$$

$$= \int_{a}^{b} \frac{t^{2}}{b-a} dt - \left(\frac{b+a}{2}\right)^{2}$$

$$= \frac{t^{3}}{3(b-a)} \Big|_{a}^{b} - \left(\frac{b+a}{2}\right)^{2}$$

$$= \frac{(a-b)^{2}}{12}$$

- Continuous-time analogue of the geometric distribution.
- How long until a server fails?

Continuous-time analogue of the geometric distribution.

How long until a server fails? How long does it take you to run into pokemon?

Continuous-time analogue of the geometric distribution.

How long until a server fails? How long does it take you to run into pokemon?

Can't "continuously flip a coin". What do we do?

Look at geometric distributions representing processes with higher and higher granularity.

Suppose a server fails with probability λ every day.

Probability that server fails on the same day as time t:

Suppose a server fails with probability λ every day.

Probability that server fails on the same day as time t:

$$(1-\lambda)^{\lceil t \rceil - 1}\lambda$$

More precision! What's the probability that it fails in a 12-hour period?

Suppose a server fails with probability λ every day.

Probability that server fails on the same day as time t:

$$(1-\lambda)^{\lceil t \rceil - 1}\lambda$$

More precision! What's the probability that it fails in a 12-hour period? $\lambda/2$ if we assume that there is no time that it's more likely to fail than another.

Suppose a server fails with probability λ every day.

Probability that server fails on the same day as time t:

$$(1-\lambda)^{\lceil t \rceil - 1}\lambda$$

More precision! What's the probability that it fails in a 12-hour period? $\lambda/2$ if we assume that there is no time that it's more likely to fail than another.

Generally: server fails with probability λ/n during any 1/n-day time period.

Suppose a server fails with probability λ every day.

Probability that server fails on the same day as time t:

$$(1-\lambda)^{\lceil t \rceil - 1}\lambda$$

More precision! What's the probability that it fails in a 12-hour period? $\lambda/2$ if we assume that there is no time that it's more likely to fail than another.

Generally: server fails with probability λ/n during any 1/n-day time period.

Probability that server fails on the same 1/n-day time period as t:

Exponential Distribution: Motivation II

Suppose a server fails with probability λ every day.

Probability that server fails on the same day as time t:

$$(1-\lambda)^{\lceil t \rceil - 1}\lambda$$

More precision! What's the probability that it fails in a 12-hour period? $\lambda/2$ if we assume that there is no time that it's more likely to fail than another.

Generally: server fails with probability λ/n during any 1/n-day time period.

Probability that server fails on the same 1/n-day time period as t:

$$\left(1-\frac{\lambda}{n}\right)^{\lceil tn\rceil-1}\frac{\lambda}{n}$$

Exponential Distribution: Motivation III

$$\left(1-\frac{\lambda}{n}\right)^{\lceil tn\rceil-1}\frac{\lambda}{n}$$

What happens when we try to take n to ∞ ? Probability goes to zero...

Exponential Distribution: Motivation III

$$\left(1-\frac{\lambda}{n}\right)^{|tn|-1}\frac{\lambda}{n}$$

What happens when we try to take n to ∞ ?

Probability goes to zero...but we can make a PDF out of this!

Remove the width of the interval (1/n) and take the limit as $n \to \infty$ to get:

$$\lim_{n \to \infty} \left(1 - \frac{\lambda}{n} \right)^{|tn|-1} \lambda = \lambda \lim_{n \to \infty} \left(1 - \frac{\lambda}{n} \right)^{tn-1}$$

Exponential Distribution: Motivation III

$$\left(1-\frac{\lambda}{n}\right)^{|tn|-1}\frac{\lambda}{n}$$

What happens when we try to take n to ∞ ?

Probability goes to zero...but we can make a PDF out of this!

Remove the width of the interval (1/n) and take the limit as $n \to \infty$ to get:

$$\lim_{n \to \infty} \left(1 - \frac{\lambda}{n} \right)^{\lceil tn \rceil - 1} \lambda = \lambda \lim_{n \to \infty} \left(1 - \frac{\lambda}{n} \right)^{tn - 1}$$
$$= \lambda e^{-\lambda t}$$

This is the PDF of the exponential distribution!

Exponential Distribution: PDF and CDF


The exponential distribution with parameter $\lambda > 0$ is defined by


Exponential Distribution: PDF and CDF

The exponential distribution with parameter $\lambda > 0$ is defined by

$$f_X(t) = \begin{cases} 0, & \text{if } t < 0 \\ \lambda e^{-\lambda t}, & \text{if } t \ge 0. \end{cases}$$

$$F_X(t) = \begin{cases} 0, & \text{if } t < 0 \\ 1 - e^{-\lambda t}, & \text{if } t \ge 0. \end{cases}$$





Note that $Pr[X > t] = e^{-\lambda t}$ for t > 0.

$$X = Expo(\lambda).$$

$$X = Expo(\lambda)$$
. Then, $f_X(x) = \lambda e^{-\lambda x} 1\{x \ge 0\}$.

$$X=Expo(\lambda)$$
. Then, $f_X(x)=\lambda e^{-\lambda x}1\{x\geq 0\}$. Thus,
$$E[X]=\int_0^\infty x\lambda e^{-\lambda x}dx$$

$$X=Expo(\lambda)$$
. Then, $f_X(x)=\lambda e^{-\lambda x}1\{x\geq 0\}$. Thus,
$$E[X]=\int_0^\infty x\lambda e^{-\lambda x}dx=-\int_0^\infty xde^{-\lambda x}.$$

$$X=Expo(\lambda)$$
. Then, $f_X(x)=\lambda e^{-\lambda x}1\{x\geq 0\}$. Thus,
$$E[X]=\int_0^\infty x\lambda e^{-\lambda x}dx=-\int_0^\infty xde^{-\lambda x}.$$

Integration by parts:

$$\int_0^\infty x de^{-\lambda x} = [xe^{-\lambda x}]_0^\infty - \int_0^\infty e^{-\lambda x} dx$$

$$X=Expo(\lambda)$$
. Then, $f_X(x)=\lambda e^{-\lambda x}1\{x\geq 0\}$. Thus,
$$E[X]=\int_0^\infty x\lambda e^{-\lambda x}dx=-\int_0^\infty xde^{-\lambda x}.$$

Integration by parts:

$$\int_0^\infty x de^{-\lambda x} = [xe^{-\lambda x}]_0^\infty - \int_0^\infty e^{-\lambda x} dx$$
$$= 0 - 0 + \frac{1}{\lambda} \int_0^\infty de^{-\lambda x} =$$

$$X=Expo(\lambda)$$
. Then, $f_X(x)=\lambda e^{-\lambda x}1\{x\geq 0\}$. Thus,
$$E[X]=\int_0^\infty x\lambda e^{-\lambda x}dx=-\int_0^\infty xde^{-\lambda x}.$$

Integration by parts:

$$\int_0^\infty x de^{-\lambda x} = [xe^{-\lambda x}]_0^\infty - \int_0^\infty e^{-\lambda x} dx$$
$$= 0 - 0 + \frac{1}{\lambda} \int_0^\infty de^{-\lambda x} = -\frac{1}{\lambda}.$$

$$X=Expo(\lambda)$$
. Then, $f_X(x)=\lambda e^{-\lambda x}1\{x\geq 0\}$. Thus,
$$E[X]=\int_0^\infty x\lambda e^{-\lambda x}dx=-\int_0^\infty xde^{-\lambda x}.$$

Integration by parts:

$$\int_0^\infty x de^{-\lambda x} = [xe^{-\lambda x}]_0^\infty - \int_0^\infty e^{-\lambda x} dx$$
$$= 0 - 0 + \frac{1}{\lambda} \int_0^\infty de^{-\lambda x} = -\frac{1}{\lambda}.$$

So: expectation is $E[X] = \frac{1}{\lambda}$.

$$X = Expo(\lambda)$$
. Then, $f_X(x) = \lambda e^{-\lambda x} 1\{x \ge 0\}$. Thus,

$$E[X] = \int_0^\infty x\lambda e^{-\lambda x} dx = -\int_0^\infty xde^{-\lambda x}.$$

Integration by parts:

$$\int_0^\infty x de^{-\lambda x} = [xe^{-\lambda x}]_0^\infty - \int_0^\infty e^{-\lambda x} dx$$
$$= 0 - 0 + \frac{1}{\lambda} \int_0^\infty de^{-\lambda x} = -\frac{1}{\lambda}.$$

So: expectation is $E[X] = \frac{1}{\lambda}$.

Variance:

$$X = Expo(\lambda)$$
. Then, $f_X(x) = \lambda e^{-\lambda x} 1\{x \ge 0\}$. Thus,

$$E[X] = \int_0^\infty x\lambda e^{-\lambda x} dx = -\int_0^\infty xde^{-\lambda x}.$$

Integration by parts:

$$\int_0^\infty x de^{-\lambda x} = [xe^{-\lambda x}]_0^\infty - \int_0^\infty e^{-\lambda x} dx$$
$$= 0 - 0 + \frac{1}{\lambda} \int_0^\infty de^{-\lambda x} = -\frac{1}{\lambda}.$$

So: expectation is $E[X] = \frac{1}{\lambda}$.

Variance: $1/\lambda^2$

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

Let
$$X = Expo(\lambda)$$
.

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

Let
$$X = Expo(\lambda)$$
. Then, for $s, t > 0$,

$$Pr[X > t + s \mid X > s] =$$

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

Let
$$X = Expo(\lambda)$$
. Then, for $s, t > 0$,

$$Pr[X > t + s \mid X > s] = \frac{Pr[X > t + s]}{Pr[X > s]}$$

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

$$Pr[X > t + s \mid X > s] = \frac{Pr[X > t + s]}{Pr[X > s]}$$
$$= \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} =$$

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

$$Pr[X > t + s \mid X > s] = \frac{Pr[X > t + s]}{Pr[X > s]}$$
$$= \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} = e^{-\lambda t}$$

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

$$Pr[X > t + s \mid X > s] = \frac{Pr[X > t + s]}{Pr[X > s]}$$
$$= \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} = e^{-\lambda t}$$
$$= Pr[X > t].$$

Similar to memorylessness for geometric distributions.

"If your server doesn't fail today, it's in the same state as it was before today."

$$Pr[X > t + s \mid X > s] = \frac{Pr[X > t + s]}{Pr[X > s]}$$
$$= \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} = e^{-\lambda t}$$
$$= Pr[X > t].$$

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$.

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$. Then
$$Pr[Y > t] =$$

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$. Then
$$Pr[Y > t] = Pr[aX > t] =$$

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$. Then
$$Pr[Y > t] = Pr[aX > t] = Pr[X > t/a]$$

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$. Then

$$Pr[Y > t]$$
 = $Pr[aX > t] = Pr[X > t/a]$
= $e^{-\lambda(t/a)} = e^{-(\lambda/a)t} =$

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$. Then

$$Pr[Y > t] = Pr[aX > t] = Pr[X > t/a]$$

$$= e^{-\lambda(t/a)} = e^{-(\lambda/a)t} = Pr[Z > t] \text{ for } Z = Expo(\lambda/a).$$

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$. Then

$$Pr[Y > t]$$
 = $Pr[aX > t] = Pr[X > t/a]$
 = $e^{-\lambda(t/a)} = e^{-(\lambda/a)t} = Pr[Z > t]$ for $Z = Expo(\lambda/a)$.

Thus, $a \times Expo(\lambda) = Expo(\lambda/a)$.

Let
$$X = Expo(\lambda)$$
 and $Y = aX$ for some $a > 0$. Then

$$Pr[Y > t]$$
 = $Pr[aX > t] = Pr[X > t/a]$
 = $e^{-\lambda(t/a)} = e^{-(\lambda/a)t} = Pr[Z > t]$ for $Z = Expo(\lambda/a)$.

Thus,
$$a \times Expo(\lambda) = Expo(\lambda/a)$$
. Also, $Expo(\lambda) = \frac{1}{\lambda} Expo(1)$.

Normal Distribution

Continuous counterpart to Binomial dist. (more on this later)

Normal Distribution

Continuous counterpart to Binomial dist. (more on this later)

Normal (or Gaussian) distribution with parameters μ , σ^2 , denoted $\mathcal{N}(\mu, \sigma^2)$:


$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$

Normal Distribution

Continuous counterpart to Binomial dist. (more on this later)

Normal (or Gaussian) distribution with parameters μ , σ^2 , denoted $\mathcal{N}(\mu, \sigma^2)$:

$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$




Normal Distribution

Continuous counterpart to Binomial dist. (more on this later)

Normal (or Gaussian) distribution with parameters μ , σ^2 , denoted $\mathcal{N}(\mu, \sigma^2)$:

$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$



Sometimes called a "bell curve". Above: $\mathcal{N}(0,1)$.

PDF:
$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$

PDF:
$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$

CDF: involves an integral with no nice closed form (often expressed in terms of "erf", the error function). Won't discuss it here.

PDF:
$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$

CDF: involves an integral with no nice closed form (often expressed in terms of "erf", the error function). Won't discuss it here.

Expectation: μ (notice that PDF is symmetric around μ).

PDF:
$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$

CDF: involves an integral with no nice closed form (often expressed in terms of "erf", the error function). Won't discuss it here.

Expectation: μ (notice that PDF is symmetric around μ).

Variance: σ^2 (fairly straightforward integration)

PDF:
$$f_X(t) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}$$

CDF: involves an integral with no nice closed form (often expressed in terms of "erf", the error function). Won't discuss it here.

Expectation: μ (notice that PDF is symmetric around μ).

Variance: σ^2 (fairly straightforward integration)

Scaling/Shifting: if $X \sim \mathcal{N}(0,1)$ and $Y = \mu + \sigma X$, then $Y \sim \mathcal{N}(\mu, \sigma^2)$.

Basically: if you take a lot of i.i.d random variables from any* distribution with zero mean and the same variance and sum them up, the sum will converge to a random Gaussian with the same mean and variance.

Basically: if you take a lot of i.i.d random variables from any* distribution with zero mean and the same variance and sum them up, the sum will converge to a random Gaussian with the same mean and variance.

Suppose $X_1, X_2, ...$ are i.i.d. random variables with expectation μ and variance σ^2 . Let

$$S_n := \frac{A_n - \mu}{\sigma / \sqrt{n}} = \frac{(\sum_i X_i) - n\mu}{\sigma / \sqrt{n}}$$

Then S_n tends towards $\mathcal{N}(0,1)$ as $n \to \infty$.

Basically: if you take a lot of i.i.d random variables from any* distribution with zero mean and the same variance and sum them up, the sum will converge to a random Gaussian with the same mean and variance.

Suppose $X_1, X_2, ...$ are i.i.d. random variables with expectation μ and variance σ^2 . Let

$$S_n := \frac{A_n - \mu}{\sigma/\sqrt{n}} = \frac{(\sum_i X_i) - n\mu}{\sigma/\sqrt{n}}$$

Then S_n tends towards $\mathcal{N}(0,1)$ as $n \to \infty$.

Or:

$$\Pr[S_n \le a] \to \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\alpha} e^{-x^2/2} dx$$

Basically: if you take a lot of i.i.d random variables from any* distribution with zero mean and the same variance and sum them up, the sum will converge to a random Gaussian with the same mean and variance.

Suppose $X_1, X_2, ...$ are i.i.d. random variables with expectation μ and variance σ^2 . Let

$$S_n := \frac{A_n - \mu}{\sigma / \sqrt{n}} = \frac{(\sum_i X_i) - n\mu}{\sigma / \sqrt{n}}$$

Then S_n tends towards $\mathcal{N}(0,1)$ as $n \to \infty$.

Or:

$$\Pr[S_n \le a] \to \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\alpha} e^{-x^2/2} dx$$

Proof: EE126

Basically: if you take a lot of i.i.d random variables from any* distribution with zero mean and the same variance and sum them up, the sum will converge to a random Gaussian with the same mean and variance.

Suppose $X_1, X_2, ...$ are i.i.d. random variables with expectation μ and variance σ^2 . Let

$$S_n := \frac{A_n - \mu}{\sigma/\sqrt{n}} = \frac{(\sum_i X_i) - n\mu}{\sigma/\sqrt{n}}$$

Then S_n tends towards $\mathcal{N}(0,1)$ as $n \to \infty$.

Or:

$$\Pr[S_n \le a] \to \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\alpha} e^{-x^2/2} dx$$

Proof: EE126

Sum of Bernoullis (binomial) tends towards normal!

Continuous probability: translation of discrete probability to a continuous sample space with an infinite number of events.

Continuous probability: translation of discrete probability to a continuous sample space with an infinite number of events.

Concepts of variance, expectation, etc. translate to continuous too.

Continuous probability: translation of discrete probability to a continuous sample space with an infinite number of events.

Concepts of variance, expectation, etc. translate to continuous too.

Geometric distribution \rightarrow exponential distribution.

Continuous probability: translation of discrete probability to a continuous sample space with an infinite number of events.

Concepts of variance, expectation, etc. translate to continuous too.

Geometric distribution \rightarrow exponential distribution.

Binomial distribution \rightarrow normal distribution.

Continuous probability: translation of discrete probability to a continuous sample space with an infinite number of events.

Concepts of variance, expectation, etc. translate to continuous too.

Geometric distribution \rightarrow exponential distribution.

Binomial distribution \rightarrow normal distribution.

Central limit theorem: everything converges to normal if we take enough samples

Today's Gig: Cauchy Distribution



Augustin-Louis Cauchy (1789-1857)



Augustin-Louis Cauchy (1789-1857)

Practically invented complex analysis. Made fundamental contributions to calculus and group theory.



Augustin-Louis Cauchy (1789-1857)

Practically invented complex analysis. Made fundamental contributions to calculus and group theory.

"More concepts and theorems have been named for Cauchy than for any other mathematician."



Augustin-Louis Cauchy (1789-1857)

Practically invented complex analysis. Made fundamental contributions to calculus and group theory.

"More concepts and theorems have been named for Cauchy than for any other mathematician."

Was also a baron



Augustin-Louis Cauchy (1789-1857)

Practically invented complex analysis. Made fundamental contributions to calculus and group theory.

"More concepts and theorems have been named for Cauchy than for any other mathematician."

Was also a baron because he tutored a duke...



Augustin-Louis Cauchy (1789-1857)

Practically invented complex analysis. Made fundamental contributions to calculus and group theory.

"More concepts and theorems have been named for Cauchy than for any other mathematician."

Was also a baron because he tutored a duke... who ended up hating math.

Actually first written about by Poisson in 1824.

Actually first written about by Poisson in 1824. Cauchy became associated with it in 1853!

Actually first written about by Poisson in 1824. Cauchy became associated with it in 1853!

Suppose I have a wall on the x-axis.

Actually first written about by Poisson in 1824. Cauchy became associated with it in 1853!

Suppose I have a wall on the x-axis. Stand at (0,1) and point a laser at a uniform random angle such that the laser hits the wall.

Actually first written about by Poisson in 1824. Cauchy became associated with it in 1853!

Suppose I have a wall on the x-axis. Stand at (0,1) and point a laser at a uniform random angle such that the laser hits the wall.

What is the distribution of the point on the wall?

Actually first written about by Poisson in 1824. Cauchy became associated with it in 1853!

Suppose I have a wall on the x-axis. Stand at (0,1) and point a laser at a uniform random angle such that the laser hits the wall.

What is the distribution of the point on the wall?

$$\tan \theta = t$$

$$\theta = \tan^{-1} t$$

$$d\theta = \frac{1}{1 + t^2} dt$$

$$\frac{d\theta}{\pi} = \frac{1}{1 + t^2} \frac{dt}{\pi}$$

PDF:

$$\frac{1}{\pi(1+t^2)}$$

$$\int_{-\infty}^{\infty} \frac{t}{\pi (1+t^2)} dt$$

PDF:

$$\frac{1}{\pi(1+t^2)}$$

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt$$

PDF:

$$\frac{1}{\pi(1+t^2)}$$

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt = 0$$

PDF:

$$\frac{1}{\pi(1+t^2)}$$

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt = 0$$
$$= \lim_{a \to \infty} \int_{-a}^{2a} \frac{t}{\pi(1+t^2)} dt$$

PDF:

$$\frac{1}{\pi(1+t^2)}$$

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt = 0$$
$$= \lim_{a \to \infty} \int_{-a}^{2a} \frac{t}{\pi(1+t^2)} dt \neq 0$$

PDF:

$$\frac{1}{\pi(1+t^2)}$$

Expectation?

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt = 0$$
$$= \lim_{a \to \infty} \int_{-a}^{2a} \frac{t}{\pi(1+t^2)} dt \neq 0$$

Expectation doesn't exist!

PDF:

$$\frac{1}{\pi(1+t^2)}$$

Expectation?

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt = 0$$
$$= \lim_{a \to \infty} \int_{-a}^{2a} \frac{t}{\pi(1+t^2)} dt \neq 0$$

Expectation doesn't exist!

If you try to estimate the expectation by sampling points and averaging, you'll get crazy results.

PDF:

$$\frac{1}{\pi(1+t^2)}$$

Expectation?

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt = 0$$
$$= \lim_{a \to \infty} \int_{-a}^{2a} \frac{t}{\pi(1+t^2)} dt \neq 0$$

Expectation doesn't exist!

If you try to estimate the expectation by sampling points and averaging, you'll get crazy results.

Variance doesn't exist either.

PDF:

$$\frac{1}{\pi(1+t^2)}$$

Expectation?

$$\int_{-\infty}^{\infty} \frac{t}{\pi(1+t^2)} dt = \lim_{a \to \infty} \int_{-a}^{a} \frac{t}{\pi(1+t^2)} dt = 0$$
$$= \lim_{a \to \infty} \int_{-a}^{2a} \frac{t}{\pi(1+t^2)} dt \neq 0$$

Expectation doesn't exist!

If you try to estimate the expectation by sampling points and averaging, you'll get crazy results.

Variance doesn't exist either.

Main takeaway: there are some really badly-behaved distributions out there.

