

Analysis of Factors Affecting Number of People in Household of Philippines

Siqi Wang, Yubin Lyu, Liting Wang, Han Xu, Jiahao Mao

Introduction

The Family Income and Expenditure Survey (FIES) is a significant source of data for understanding the wellbeing of households in Philippines. It provides valuable information on family income and expenditure, which can be used to investigate various research questions related to household characteristics.

In this analysis, we are interested in identifying which household-related factors influence the size of a household. Using Generalized Linear Model (GLM), we will explore the datasets obtained from the FIES survey for XII - SOCCSKSARGEN region in Philippines. The results of our analysis could help the government to make informed decisions related to household policies and other related matters.

Data Processing

```
# Load data
household <- read.csv("dataset4.csv")
# Factorize the categorical variables
household$Electricity <- as.factor(household$Electricity)
household$Household.Head.Sex <- as.factor(household$Household.Head.Sex)
household$Type.of.Household <- as.factor(household$Type.of.Household)

# Simplified column names
colnames(household)[1]<-"Income"           # original name "Total.Household.Income"
colnames(household)[3]<-"FoodExp"          # original name "Total.Food.Expenditure"
colnames(household)[4]<-"Householder_Sex"  # original name "Household.Head.Sex"
colnames(household)[5]<-"Householder_Age"  # original name "Household.Head.Age"
colnames(household)[6]<-"Household_Type"   # original name "Type.of.Household"
colnames(household)[7]<-"Number_Members"   # original name "Total.Number.of.Family.members"
colnames(household)[8]<-"Floorarea"        # original name "House.Floor.Area"
colnames(household)[10]<-"Number_bedrooms" # original name "Number.of.bedrooms"
household$Household_Type <- ifelse(household$Household_Type == "Two or More Nonrelated Persons/Members"
```

Data Summary

```
# Summary of Categorical Variables
household_cat <- household %>%
  dplyr::select("Electricity", "Householder_Sex", "Household_Type")
summary(household_cat)
```

```
Electricity Householder_Sex Household_Type
0: 363      Female: 362      Length:2122
1:1759      Male   :1760      Class :character
                                Mode  :character
```

```
# Summary of Numerical Variables
household_num <- household[, sapply(household, is.numeric)]
my_skim <- skim_with(base = sfl(n = length))
household_num %>%
  my_skim() %>%
  transmute(Variable=skim_variable, n=n,
            Mean = format(signif(numeric.mean, 3), scientific = TRUE, digits = 2),
            SD = format(signif(numeric.sd, 3), scientific = TRUE, digits = 2),
            Min= format(signif(numeric.p0, 3), scientific = TRUE, digits = 2),
            Median=format(signif(numeric.p50, 3), scientific = TRUE, digits = 2),
            Max=format(signif(numeric.p100, 3), scientific = TRUE, digits = 2),
            IQR = format(signif(numeric.p75-numeric.p50, 3), scientific = TRUE, digits = 2) ) %>%
  kable(caption = '\\\\label{tab:summarybyskim} Summary statistics of variables',
        booktabs = TRUE, linesep = "", digits = 2) %>%
  kable_styling(font_size = 10, latex_options = "hold_position")
```

Table 1: Summary statistics of variables

Variable	n	Mean	SD	Min	Median	Max	IQR
Income	2122	1.8e+05	2.3e+05	1.5e+04	1.2e+05	3.2e+06	7.4e+04
FoodExp	2122	7.2e+04	4.5e+04	7.8e+03	6.3e+04	7.3e+05	2.4e+04
Householder_Age	2122	4.9e+01	1.4e+01	9.0e+00	4.8e+01	9.9e+01	1.1e+01
Number_Members	2122	4.5e+00	2.2e+00	1.0e+00	4.0e+00	1.9e+01	2.0e+00
Floorarea	2122	3.6e+01	3.5e+01	5.0e+00	2.6e+01	4.5e+02	1.4e+01
House.Age	2122	1.6e+01	1.1e+01	0.0e+00	1.4e+01	7.5e+01	7.0e+00
Number_bedrooms	2122	1.8e+00	1.0e+00	0.0e+00	2.0e+00	7.0e+00	0.0e+00

Distribution Check

Test if the distribution of y follows the poisson distribution.

```
# If a variable follows poisson distribution variance will equal to mean.
print(var(household$Number_Members))
```

```
[1] 4.9
```

```
print(mean(household$Number_Members))
```


```
[1] 4.5
```

```
# As we can see there is no significant difference between mean and variance
# Plot the histogram of response variable and compare it with poisson distribution.
hist(household$Number_Members, freq = FALSE, xlab = "Data",
```

```

    main = "Histogram of Number of Family members")
# Overlay a Poisson probability mass function
x <- 0:max(household$Number_Members)
lines(x, dpois(x, lambda = 4), col = "red")

```



Based on the plot we can see that the distribution follows poisson dist when lambda = 4

Correlation Matrix and ggpairs

```

# Create the correlation matrix of variables
cor_matrix <- cor(household_num) %>%
  round(2)
cor_matrix


```

	Income	FoodExp	Householder_Age	Number_Members	Floorarea
Income	1.00	0.59	0.08	0.12	0.38
FoodExp	0.59	1.00	0.04	0.43	0.28
Householder_Age	0.08	0.04	1.00	-0.08	0.12
Number_Members	0.12	0.43	-0.08	1.00	0.04
Floorarea	0.38	0.28	0.12	0.04	1.00
House.Age	0.12	0.09	0.32	-0.07	0.15
Number_bedrooms	0.42	0.41	0.15	0.16	0.43


House.Age Number_bedrooms

Income	0.12	0.42
FoodExp	0.09	0.41
Householder_Age	0.32	0.15
Number_Members	-0.07	0.16
Floorarea	0.15	0.43
House.Age	1.00	0.27
Number_bedrooms	0.27	1.00

```
corrplot <- corrplot(cor_matrix, method = "color", addCoef.col = "gray", type = "upper",)
```




```
# Create the ggpairs of variables
ggpairs(household_num)
```




Boxplots of Variables


```
# Explanatory analysis on numeric variables
ggplot(data = household, mapping = aes(x = Householder_Sex, y = Number_Members)) +
  geom_boxplot(aes(fill = Householder_Sex)) +
  labs(x = "Householder_Sex", y = "Number_Members",
       title = "gender") +
  scale_x_discrete(labels = c("female", "male"))
```



```
ggplot(data = household, mapping = aes(x = Household_Type, y = Number_Members)) +
  geom_boxplot(aes(fill = Household_Type))+
  labs(x = "Household_Type", y = "Number_Members", title = "type") +
  scale_x_discrete(labels = c("Extended", "Single", "Two or More"))+
  theme(legend.position = "bottom")
```



```
ggplot(data = household, mapping = aes(x = Electricity, y = Number_Members)) +
  geom_boxplot(aes(fill = Electricity ))+
  labs(x = "Electricity", y = "Number_Members",
       title = "Electricity") +
  scale_x_discrete(labels = c("no", "yes"))
```




Histogram of Variables


```
# Explanatory analysis on categorical variables
household %>%
  tabyl(Number_Members, Householder_Sex) %>%
  adorn_percentages() %>%
  adorn_pct_formatting() %>%
  adorn_ns() # To show original counts
```

Number_Members		Female		Male
1	37.2%	(45)	62.8%	(76)
2	30.8%	(69)	69.2%	(155)
3	25.1%	(92)	74.9%	(274)
4	12.7%	(58)	87.3%	(397)
5	10.3%	(40)	89.7%	(349)
6	12.2%	(28)	87.8%	(202)
7	7.0%	(9)	93.0%	(120)
8	11.4%	(12)	88.6%	(93)
9	12.8%	(6)	87.2%	(41)
10	3.8%	(1)	96.2%	(25)
11	0.0%	(0)	100.0%	(13)
12	0.0%	(0)	100.0%	(7)
13	33.3%	(1)	66.7%	(2)
14	0.0%	(0)	100.0%	(3)
15	0.0%	(0)	100.0%	(1)
18	0.0%	(0)	100.0%	(1)
19	50.0%	(1)	50.0%	(1)


```
# barplot of Number_Members and Householder_Sex
ggplot(household, aes(x= Number_Members, y = ..prop.., group=Householder_Sex, fill=Householder_Sex)) +
  geom_bar(position="dodge", stat="count") +
  labs(y = "Proportion")
```



```
# barplot of Number_Members and Electricity
ggplot(household, aes(x= Number_Members, y = ..prop.., group=Electricity, fill=Electricity)) +
  geom_bar(position="dodge", stat="count") +
  labs(y = "Proportion")
```



```
# barplot of Number_Members and Household_Type
ggplot(household, aes(x= Number_Members, y = ..prop.., group=Household_Type, fill=Household_Type)) +
  geom_bar(position="dodge", stat="count") +
  labs(y = "Proportion") +
  theme(legend.position = "bottom")
```

Model Fitting

Since our response variable follows poisson distribution, we will use glm with poisson distribution to fit the model.

```
#model1 with all variables
m1 <- glm(formula = Number_Members ~ Income + FoodExp + Householder_Sex +
           Householder_Age + Household_Type + Floorarea +
           House.Age + Number_bedrooms + Electricity,
           family = poisson(link = "log"), data = household)

#model2 with log transformation for Income and FoodExp
m2 <- glm(formula = Number_Members ~ log(Income) + log(FoodExp) + Householder_Sex +
           Householder_Age + Household_Type + Floorarea +
           House.Age + Number_bedrooms + Electricity,
           family = poisson(link = "log"), data = household)

summary(m1)
```

Call:

```
glm(formula = Number_Members ~ Income + FoodExp + Householder_Sex +
    Householder_Age + Household_Type + Floorarea + House.Age +
    Number_bedrooms + Electricity, family = poisson(link = "log"),
    data = household)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-4.523	-0.615	-0.113	0.423	4.115

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	1.60e+00	6.09e-02	26.21	< 2e-16 ***
Income	-2.39e-07	5.63e-08	-4.23	2.3e-05 ***

FoodExp	2.93e-06	1.88e-07	15.59	< 2e-16	***
Householder_SexMale	2.63e-01	3.05e-02	8.62	< 2e-16	***
Householder_Age	-3.80e-03	8.10e-04	-4.68	2.8e-06	***
Household_Type2	-3.47e-01	2.29e-02	-15.13	< 2e-16	***
Household_Typetwo or more	-1.06e-01	1.81e-01	-0.59	0.55842	
Floorarea	-4.94e-04	3.40e-04	-1.45	0.14648	
House.Age	-3.71e-03	1.03e-03	-3.61	0.00031	***
Number_bedrooms	5.01e-02	1.23e-02	4.06	4.9e-05	***
Electricity1	-9.03e-02	2.85e-02	-3.17	0.00154	**

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 2217.8 on 2121 degrees of freedom
 Residual deviance: 1551.8 on 2111 degrees of freedom
 AIC: 8512

Number of Fisher Scoring iterations: 5

```
summary(m2)
```

Call:

```
glm(formula = Number_Members ~ log(Income) + log(FoodExp) + Householder_Sex +
    Householder_Age + Household_Type + Floorarea + House.Age +
    Number_bedrooms + Electricity, family = poisson(link = "log"),
    data = household)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.960	-0.557	-0.110	0.422	3.859

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-2.951300	0.248609	-11.87	< 2e-16 ***
log(Income)	-0.137026	0.021684	-6.32	2.6e-10 ***
log(FoodExp)	0.577842	0.029121	19.84	< 2e-16 ***
Householder_SexMale	0.203725	0.030685	6.64	3.2e-11 ***
Householder_Age	-0.002625	0.000823	-3.19	0.00142 **
Household_Type2	-0.288165	0.023185	-12.43	< 2e-16 ***
Household_Typetwo or more	-0.035410	0.180946	-0.20	0.84485
Floorarea	-0.000904	0.000341	-2.65	0.00804 **
House.Age	-0.003815	0.001032	-3.70	0.00022 ***
Number_bedrooms	0.024816	0.012572	1.97	0.04840 *
Electricity1	-0.159250	0.029844	-5.34	9.5e-08 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 2217.8 on 2121 degrees of freedom
 Residual deviance: 1299.4 on 2111 degrees of freedom
 AIC: 8260

Number of Fisher Scoring iterations: 4

Use BIC to do variable selection

BIC is implemented to found the best fitting model.

```
output <- bic.glm(Number_Members ~ log(Income) + log(FoodExp) + Householder_Sex +
                  Householder_Age + Household_Type + Floorarea +
                  House.Age + Number_bedrooms + Electricity,
                  glm.family = "poisson" , data = household)

summary(output)
```

Call:

```
bic.glm.formula(f = Number_Members ~ log(Income) + log(FoodExp) +      Householder_Sex + Householder_Age
```

5 models were selected

Best 5 models (cumulative posterior probability = 1):

	p!=0	EV	SD	model 1	model 2
Intercept	100	-3.025085	0.246656	-2.97e+00	-3.10e+00
log(Income).x	100.0	-0.136207	0.021240	-1.37e-01	-1.28e-01
log(FoodExp).x	100.0	0.582453	0.029086	5.80e-01	5.84e-01
Householder_Sex.x	100.0				
.Male		0.205620	0.031117	2.03e-01	2.01e-01
Householder_Age.x	79.8	-0.002132	0.001308	-2.66e-03	-2.53e-03
Household_Type2.x	100.0	-0.286849	0.023801	-2.90e-01	-2.89e-01
Household_Typetwo or more.x	0.0	0.000000	0.000000	.	.
Floorarea.x	21.2	-0.000153	0.000331	.	-7.04e-04
House.Age.x	96.5	-0.003704	0.001268	-3.69e-03	-3.53e-03
Number_bedrooms.x	0.0	0.000000	0.000000	.	.
Electricity.x	100.0				
.1		-0.156645	0.029881	-1.57e-01	-1.57e-01
nVar				7	8
BIC				-1.49e+04	-1.49e+04
post prob				0.610	0.152
	model 3	model 4	model 5		
Intercept	-3.12e+00	-3.25e+00	-2.93e+00		
log(Income).x	-1.42e-01	-1.32e-01	-1.43e-01		
log(FoodExp).x	5.87e-01	5.90e-01	5.82e-01		
Householder_Sex.x					
.Male	2.18e-01	2.15e-01	2.07e-01		
Householder_Age.x	.	.	-3.46e-03		
Household_Type2.x	-2.74e-01	-2.74e-01	-2.87e-01		
Household_Typetwo or more.x	.	.	.		
Floorarea.x	.	-7.72e-04	.		
House.Age.x	-4.58e-03	-4.35e-03	.		
Number_bedrooms.x	.	.	.		
Electricity.x					

	.1	-1.54e-01	-1.55e-01	-1.68e-01
nVar	6	7	6	
BIC	-1.49e+04	-1.49e+04	-1.49e+04	
post prob	0.143	0.059	0.035	

1 observations deleted due to missingness.

```
# Name the best model selected by BIC m3
m3 <- glm(Number_Members ~ log(Income) + log(FoodExp) + Householder_Sex +
          Householder_Age + Household_Type +
          House.Age + Electricity,
          family = "poisson" , data = household)
summary(m3)
```

Call:

```
glm(formula = Number_Members ~ log(Income) + log(FoodExp) + Householder_Sex +
     Householder_Age + Household_Type + House.Age + Electricity,
     family = "poisson", data = household)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.912	-0.569	-0.108	0.420	3.924

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-2.96574	0.23096	-12.84	< 2e-16 ***
log(Income)	-0.13692	0.02069	-6.62	3.7e-11 ***
log(FoodExp)	0.58016	0.02894	20.04	< 2e-16 ***
Householder_SexMale	0.20301	0.03065	6.62	3.5e-11 ***
Householder_Age	-0.00266	0.00082	-3.25	0.00115 **
Household_Type2	-0.29073	0.02316	-12.55	< 2e-16 ***
Household_Typetwo or more	-0.03024	0.18093	-0.17	0.86726
House.Age	-0.00370	0.00102	-3.63	0.00028 ***
Electricity1	-0.15666	0.02980	-5.26	1.5e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)

Null deviance: 2217.8 on 2121 degrees of freedom
 Residual deviance: 1308.2 on 2113 degrees of freedom
 AIC: 8264

Number of Fisher Scoring iterations: 4

Negative Binomial Distribution

The variance(4.9) of y is slightly larger than the mean(4.5) of y, therefore a Negative Binomial Distribution model is fitted to reduce the issue of overdispersion.

```
m4 <- glm.nb(formula = Number_Members ~ log(Income) + log(FoodExp) + Householder_Sex +
              Householder_Age + Household_Type + Floorarea +
              House.Age + Number_bedrooms + Electricity, data = household)
summary(m4)
```

Call:

```
glm.nb(formula = Number_Members ~ log(Income) + log(FoodExp) +
       Householder_Sex + Householder_Age + Household_Type + Floorarea +
       House.Age + Number_bedrooms + Electricity, data = household,
       init.theta = 109689.4119, link = log)
```

Deviance Residuals:

```
      Min       1Q   Median       3Q      Max
-2.960  -0.557  -0.110   0.422   3.859
```

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-2.951350	0.248617	-11.87	< 2e-16 ***
log(Income)	-0.137029	0.021684	-6.32	2.6e-10 ***
log(FoodExp)	0.577850	0.029122	19.84	< 2e-16 ***
Householder_SexMale	0.203724	0.030685	6.64	3.2e-11 ***
Householder_Age	-0.002625	0.000823	-3.19	0.00143 **
Household_Type2	-0.288164	0.023186	-12.43	< 2e-16 ***
Household_Typetwo or more	-0.035409	0.180951	-0.20	0.84486
Floorarea	-0.000904	0.000341	-2.65	0.00804 **
House.Age	-0.003815	0.001032	-3.70	0.00022 ***
Number_bedrooms	0.024815	0.012572	1.97	0.04841 *
Electricity1	-0.159252	0.029845	-5.34	9.5e-08 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for Negative Binomial(109689) family taken to be 1)

```
Null deviance: 2217.7 on 2121 degrees of freedom
Residual deviance: 1299.4 on 2111 degrees of freedom
AIC: 8262
```

Number of Fisher Scoring iterations: 1

```
      Theta: 109689
    Std. Err.: 356155
Warning while fitting theta: iteration limit reached
```

```
2 x log-likelihood: -8238
```


Deviance plots

Deviance plots is shown below.

```

resp <- resid(m1, type = "pearson")
resd <- resid(m1, type = "deviance")
r1<- ggplot(m1, aes(sample = resp)) + geom_point(stat = "qq", color = "#7fc97f") + ylab("Pearson residuals")
r2<- ggplot(m1, aes(sample = resd)) + geom_point(stat = "qq", color = "#7fc97f") + ylab("Deviance residuals")
r3<- ggplot(m1, aes(x = predict(m1, type="link"), y =resd))+ geom_point(col = "#7fc97f") +
  ylab("Deviance residuals") + xlab("Linear predictor")
grid.arrange(r1, r2, r3, nrow = 1)


```



```

resp2 <- resid(m2, type = "pearson")
resd2 <- resid(m2, type = "deviance")
r4<- ggplot(m2, aes(sample = resp2)) + geom_point(stat = "qq", color = "#7fc97f") + ylab("Pearson residuals")
r5<- ggplot(m2, aes(sample = resd2)) + geom_point(stat = "qq", color = "#7fc97f") + ylab("Deviance residuals")
r6<- ggplot(m2, aes(x = predict(m2, type="link"), y =resd2))+ geom_point(col = "#7fc97f") +
  ylab("Deviance residuals") + xlab("Linear predictor")
grid.arrange(r4, r5, r6, nrow = 1)

```



Model Evaluation

```
# Poisson model
c(m1$deviance, m1$aic)
```

```
[1] 1552 8512
```

```
# poisson model with log transformation
c(m2$deviance, m2$aic)
```

```
[1] 1299 8260
```

```
# BIC model
c(m3$deviance, m3$aic)
```

```
[1] 1308 8264
```

```
# Negative binomial model
c(m4$deviance, m4$aic)
```

```
[1] 1299 8262
```

Goodness-of-fit test

```
chisq <- with(m1, sum((household$Number_Members- fitted.values)^2/fitted.values))
df <- with(m1, df.residual)
p <- with(m1, pchisq(chisq, df, lower.tail = FALSE))
cat("Chi-square test statistic = ", chisq, "\n")
```

Chi-square test statistic = 1584

```
cat("df = ", df, "\n")
```

df = 2111

```
cat("p-value = ", p, "\n")
```

p-value = 1

```
# The coef() function obtains the coefficients of the model.
# The confint() function obtains the confidence interval of the model coefficients.
```

```
exp(cbind(OR = coef(m1), confint(m1)))
```

	OR	2.5 %	97.5 %
(Intercept)	4.94	4.38	5.57
Income	1.00	1.00	1.00
FoodExp	1.00	1.00	1.00
Householder_SexMale	1.30	1.23	1.38
Householder_Age	1.00	0.99	1.00
Household_Type2	0.71	0.68	0.74
Household_Typetwo or more	0.90	0.62	1.26
Floorarea	1.00	1.00	1.00
House.Age	1.00	0.99	1.00
Number_bedrooms	1.05	1.03	1.08
Electricity1	0.91	0.86	0.97

```
#In Poisson regression, OR (odds ratio) represents the probability ratio (probability ratio) of a set o
```

```
#The OR value is equal to the regression coefficient of the indexed variable. The coefficient of an exp
```