



Intubation Checklist

Team Preparation

- ☐ Gloves / Glasses / N95 / Face Shield
- ☐ Establish Roles (Lead Intubator and Assistant)
- ☐ Establish Goals (First pass success, under 30 seconds between oxygenation, max 2 attempts)
- ☐ Rescue plan (iGel, BVM with adjuncts)

Patient Preparation

- ☐ Optimize Hemodynamics (Normal Saline / Dopamine if applicable)
- ☐ Position Patient (Head elevated, sniffing position, axis aligned)
- ☐ Pre-Oxygenate (BVM, Consider Hi-Flow Nasal Canula)
- ☐ Patient connected to monitor (4-lead, NIBP, SPO2, ETCO2)
- ☐ IV Access

Equipment


- ☐ BVM with Adjuncts Attached to O2
- ☐ Laryngoscope & Blade Tested
- ☐ Suction - Tested & Under Patient's Right Shoulder (Consider Big Stick Attachment for SALAD Maneuver)
- ☐ ET Tube - 10cc Syringe, Lubrication, Verify Cuff
- ☐ Bougie Stylet - Out of Package, Lubricated
- ☐ Thomas Tube Tie - Out of Package and Under Patient's Neck
- ☐ Cervical Collar for SET Protocol
- ☐ Rescue Airways - Nearby and Readily Available
- ☐ Stopwatch to Ensure Attempt Does Not Exceed 30 Seconds

Troubleshooting A Bad View


- Elevate the patient's head and/or lower your head improve view
- Utilize cricoid pressure or BURP and have assistant hold pressure
- Avoid levering and lift at 45 degrees (aim between wall and ceiling)
- Utilize atlanto-occipital joint role (tilt head 30 degrees to the right)
- Have assistant pull on patient's lip with finger to open view

Reference Material

Ideal Positioning



BURP Maneuver



Cormac Lehane Grades I-IV

