Human Reproduction Questions and Answers Chapter-3

| Marks Questions Failure of testes to descend into scrotal sacs leads to sterility. Why? Ans. High temperature of abdomen kills the spermatogenic tissue of the testes, so no sperm are formed. | | |
|--|----|--|
| 2. Both vaccine and colostrum produce immunity. Name type of immunity produced by these. Ans. Vaccine Active immunity Colostrum Passive immunity. | | |
| 3. How many sperms will be produced from 10 primary spermatocytes and how many egg will be produced from 10 primary oocytes? Ans. 40 sperms, 10 eggs. | gs | |
| 4. The spermatogonial cell has 46 chromosomes in human male. Give the number of chromosomes in (a) Primary spermatocyte (b) Spermatid Ans. (a) 46 in Primary spermatocyte (b) 23 in spermatid. | | |
| 5. In ovary which structure transforms as corpus luteum and name the hormone secreted by corpus luteum? Ans. Follicular cells of empty Graafian follicle. Progesterone. | | |

| 6. "Each and every coitus does not results in fertilisation and pregnancy". Justify the statement. Ans. Ovum and sperm should reach simultaneously to the ampullary – isthmic junction. | | |
|--|--|--|
| 7. Why are male testes located outside the abdominal cavity? Ans. The male testes are located in the scrotum outside the abdominal cavity as the scrotum provides low temperature than the normal body temperature required for spermatogenesis. | | |
| 8. State the function of leydig cells. Ans. The leydig cells synthesise and secrete testicular hormones called androgens. | | |
| 9. Where do we find fimbriae?Ans. Fimbriae are finger like projections found in the edges of the infundibulum. | | |
| 10. What is semen?Ans. The seminal plasma along with the sperms constitutes semen. | | |
| 11. Define parturition. Ans. The vigorous contraction of the uterus that results into the delivery of the child at the end of pregnancy is called parturition. | | |
| | | |

12. Where does fertilization normally takes place in a human female. **Ans.** Ampulla (fallopian tube).

| 13. Name the substance present in the sperm acrosome & which help in sperms entry into egg. Ans. Acrosome contains enzymes e.g. hyaluronidase that helps in dissolving membrane of ovum. |
|--|
| 14. Name the layer of cells that forms the outer wall of blastocyst. Ans. Trophoblast. |
| 15. At what stage is the mammalian embryo implanted in uterus? Ans. Blastocyst stage. |
| 16. Despite the presence of So many sperms in the vicinity of an egg cell, only one sperm enters the ovum. Why? Ans. Because when a sperm comes in contact with ovum (zona pellucida) & induces changes in membrane to block entry of other sperms. |
| 17. How many polar bodies are given out in production of one egg during cogenesis? Ans. Two polar bodies |
| |

2 Mark Questions


Chapter 3 Human Reproduction

- **2 Marks Questions**
- 1. Give the function of
- (a) Corpus luteum
- (b) Endometrium

Ans. Corpus luteum: It secretes progesterone which prepares endometrium of uterus for implantation and normal development of foetus.

Endometrium : It undergoes cyclic changes during menstrual cycle and prepares itself for implantation of blastocyst.


2. In the given figure, give the name and functions of parts labeled A and B.



Ans. A = Trophoblast Gets attached to endometrium and draws nutritive material secreted by uterine endometrium gland.

B = Inner cell mass Differentiates as Embryo.

3. Given below is an incomplete flow chart showing influence of hormone on gametogenesis in male, observe the flow chart carefully and fill in the blank A, B, C and D.



Ans. A = Testosterone; B = Spermatogenesis C = Sertoli cells; D Spermiogenesis

4. Give reason for the following:

- (a) The first half of the menstrual cycle is called follicular phase as well as proliferative phase.
- (b) The second half of the menstrual cycle is called luteal phase as well as secretory phase. Ans. (a) During this phase, primary follicles transform into Graafian follicle under FSH stimulation. Graafian follicles secrete estrogens with stimulate enlargement of Endometrium of uterus.
- (b) During this phase, Corpus luteum is fully formed and secretes large quantity of Progestrone.

5. What is meant by L.H. Surge? Write the role of L.H.


Ans. There are three phases in your menstrual cycle – follicular phase, ovulatory phase and luteal phase. In terms of the luteinizing hormone (LH) surge, the ovulatory phase is most important. During the follicular phase the follicle develops at the beginning of the menstrual cycle. This cycle begins with the menstrual period, the shedding of the interine lining and the shedding cleanses the lining of the uterus in preparation for ovulation during the ovulatory phase.

6. Explain significance of the condition in which the testes remain suspended in scrotum outside the abdomen.

Ans. Human sperm cells cannot develop at body temperature. Spermatogenesis and maintenance of the seminiferous tubules requires a temperature slightly lower than that of the body. This is provided by the scrotum, which lies outside the abdominal cavity.

7. Describe the structure of a sperm with a diagram.

Ans. The human sperm is a microscopic structure with a head, middle piece and a tail. The head has the haploid nucleus and an anterior acrosome that contains the enzymes required for the fertilization of the egg. The middle piece has numerous mitochondria to produce the energy for the mobility of the tail of the sperm.



8. Enlist any two functions of a female placenta.

Ans. The structural and the functional unit between the developing embryo and the mother called placenta facilitates the supply of nutrients, oxygen to the embryo and also the removal of carbon dioxide and other excretory products produced by the embryo. It also acts as endocrine tissue and produces several hormones

9. What is the number of chromosomes in the following cells? Primary oocyte, secondary oocyte, ootid and follicle.

Ans. The number of chromosome in the cells is as follows:

Primary oocyte: 23 pairs. Secondary oocyte: 23. Ootid: 23. Follicle: 23 pairs.


10. What is corpus luteum. How dose it functions as endocrine gland?

Ans. After ovulation, the graffian follicle ruptures & forms corpus luteum. Corpus luteum functions as endocrine glands as they secrete progesterone & estrogen in large quantities.

11. Where are leydig cells located? What do they secrete?

Ans. Leydig cells or interstitial cells are located in between the sominiferous tubules. Leydig cells secrete male sex hormone TESTOSTERONE which promotes development of accessory glands & control male secondary sexual characters.

12. Draw well labeled diagram of T.S. of ovary? Ans.




13. Why testes of human males are considered extra abdominal? What is the significance of this condition?

Ans. Testis in human males are called extra- abdominal because testis are located outside the abdominal cavity in a pouch called scrotum which provides a temperature 2-3oC lower than body temperature necessary for spermatogenesis.

14. Draw a diagram of the T.S. of seminiferous tubule of testis of an adult human male & label any four parts in it.

Ans.



15. What is colustrum? What is its significance to new born baby?

Ans. The milk secreted from mammary glands just after birth for 2 or 3 days is called colustrum. It is rich in proteins & low in fats. It also contains antibody IgA which provides immunity to new born infant.

3 Mark Questions

Chapter 3 Human Reproduction

3 Marks Questions

1. Mention the name and role of hormones which are involved in regulation of gamete formation in human male.

Ans. GnRH: Stimulates adenophysis to secrete gonadotrophins.

GSH: Stimulates Sertoli cells to secrete factors while help inspermatogenesis.

ICSH: Stimulates interstitial cells to secrete testosterone.


2. Three of the steps of neuro endocrine mechanism in respect of parturition are mentioned below.

Write the missing steps in proper sequence.

- (a) Signals originate from fully developed foetus and placenta.
- (b) _____.
- (c) _____.
- (d) Oxytocin causes strong uterine contraction
- (e) Uterine contraction stimulates further secretion of oxytocin.
- (f) _____.

Ans. (b) Foetal ejection reflex

- (c) The reflex triggers release of oxytocin
- (f) Expulsion of the baby out through birth canal.
- 3. The events of the menstrual cycle are represented below. Answer the following questions.



- (i) State the levels of FSH, LH and Progesterone simply by mentioning high or low around 13th and 14th day and 21st to 23rd day.
- (ii) In which of the above mentioned phases does egg travel to fallopian tube?
- (iii) Why there is no mensuration after fertilisation?


Ans. (i) 13-14th day 21st -23rd day

FSH – High Low

LH – High Low

Progesterone – Low High

- (ii) End of follicular or proliferative phase.
- (iii) Menstruation does not occur during pregnancy upon fertilization due to high level of progestenone secreted by persisting corpus luteum and Placenta.
- 4. (a) Read the graph given below. Correlate the ovarian events that take place in the human female according to the level of the pituitary hormone during the following day.



- (i) 10th 14th days (ii) 14th -15th days
- (iii) 16th 23th days (iv) 25th 29th days
- (If the ovum is not fertilised)
- (b) What are the uterine events that follow beyond 29th day if the ovum is not fertilised?

Ans. (a) (i) Gonadotropins and FSH increases

- (ii) LH attains peak level but FSH decreases
- (iii) LH and FSH level decreases
- (iv) LH remains low and FSH increases.
- **(b)** After 29th day there is a mentrual flow involving discharge of blood and cast off endometrium lining.

5. T.S. of mammalian testis revealing seminiferous tubules show different types of cell.

- (i) Name the two types of cells of germinal epithelium.
- (ii) Name of cells scattered in connective tissue and lying between seminiferous tubules. Differentiate between them on the basis of their functions.

Ans. (i) Germinal epithelium have two types of cell. 1. Spermatogonium. 2. Sertoli cells (ii) Leydig cells or Interstitial cells.

Functions

Spermatogonium undergoes meiotic division leading to sperm formation.

Sertoli cell: Nourishes germ cells

Leydig cell: Synthesise and Secrete hormone androgen.

6. What are the various male accessory glands? Give their function.


Ans. The male accessory glands include paired seminal vesicles, a prostrate gland and paired bulbourethral glands.

These glands secrete seminal plasma rich in fructose, calcium and certain enzymes. Secretions of bulbourethral glands help in lubrication of the penis.

7. Explain the menstrual cycle with a diagram.

Ans. Menstrual cycle has three phases: menstrual, proliferative and secretory.

- a) Menstrual Phase: The phase lasts for 3-5 days in human females and during this period the endometrial lining of the uterus is cast off and is slowly passed out from vagina as a mixture of blood.
- b) Proliferative or Follicular Phase: It lasts for 11 days between 6th to 16th day of the cycle. During this phase one ovarian follicle is changed into Graafian follicle and the endometrial layer is rebuilt along with repair of the ruptured blood vessels. Estrogen increases. It ends with ovulation.
- c) Secretory Phase: It lasts for 12 days between 17-28 days. The Graafian follicle is converted to Corpus Luteum. The endometrium grows and thickens further. Progesterone increases. It end with the conversion of corpus luteum to corpus albic



8. Differentiate between spermatogenesis and oogenesis. Ans.

| Spermatogenesis | Oogenesis |
|---|--|
| 1. It occurs inside the testes. | 1. It occurs inside the ovary. |
| 2. All the stages are completed inside the testes. | 2. Majority occurs inside the ovary but last stages occur in the oviduct. |
| 3. Spermatogonia develop from the germinal epithelium lining in the seminiferous tubules. | 3. Oogonia develop from the germinal epithelium overlying the ovary. |
| 4. All spermatogonia give rise to spermatocytes. | 4. Only few oogonia give rise to oocytes. |
| 5. Primary spermatocytes divide by meiosis I to give rise to two secondary spematocytes | 5. Primary oocyte undergoes meiosis I to give rise to one secondary oocyte and a polar body. |
| 6. Secondary spermatocyte divides by meiosisII to give rise to two spermatids. | 6. Secondary oocyte divides by meiosisII to form the ovum and the second polar body. |

9. 'A fertilized egg is the blue print of future development'. Explain

Ans. The sperm carries the genetic information from the father in form of 23 chromosomes (including the male sex chromosome X or Y) while the egg bears the genetic information from the mother (including the female sex chromosome X). Thus during fertilization the fusion of the male and the female gametes produce new genetic combination which introduces variation in the progeny. The zygote or the fertilized egg contain the genetic information which accordingly controls the development of the embryo.