

PRODUCT TECHNICAL BULLETIN

DIXIE CHEMICAL
10601 Bay Area Blvd.
Pasadena, TX 77507
Tel: (281) 474-3271
Fax: (281) 291-3384
E-Mail: info@dixiechemical.com

3-CHLORO-1, 2-PROPANEDIOL CHLOROGLYCERINE (CG)

CASRN 96-24-2

Typical Properties

Property	Value	Test Method
Appearance	Clear to light yellow viscous liquid	D-104
Color, Pt-Co	50 Max	D-103
Specific Gravity at 25°C	1.310 – 1.325	D-102
Chloroglycerine, %	98.5 Min	D-131C
Water, %	1.0 Max	D-101

Applications

Chloroglycerine can be used as a starting material for the preparation of pharmaceutical intermediates. It is also used as a fungicide and has found efficacy as a chemosterilant for rodent control. CG is commonly used to prepare glycidol ethers, esters and amines.

Handling & Storage

Chloroglycerine should be stored in a well-ventilated area. Drums should be kept tightly sealed.

Health Hazards

Read and understand the relevant Safety Data Sheets (SDS) for all products before use. Both mono- and dichloroglycerines are harmful when taken internally and when absorbed through the skin. Extreme care

J

TECHNICAL BULLETIN

Chloroglycerine (CG)

should be taken to avoid any exposure to this chemical. In case of contact, follow the procedures outlined in the SDS. Generally, these procedures include immediately flushing the affected skin or eyes with copious amounts of water for at least 15 minutes. In case of contact to eyes, skin or by inhalation, get medical attention. Wash contaminated clothing before reuse.

Follow the recommendations in the SDS for personal protective equipment when handling chloroglycerine. Impenetrable gloves, chemically impervious apron over lab coat and full coverage clothing are recommended. Wear chemical goggles and a face shield in combination. If ventilation is insufficient, wear respiratory protection.

Availability

Chloroglycerine is available from Dixie Chemical in Pasadena, Texas. Contact your Dixie representative for details.