

## Useful formulas for “Stochastic Processes”

### Chapter 1

- $\mathbb{P}\left(\bigcup_1^n A_i\right) = \sum_{k=1}^n (-1)^{k+1} \sum_{\{i_1, \dots, i_k\}} \mathbb{P}(A_{i_1} \cap \dots \cap A_{i_k})$
- The coupon collection problem:  
 $\mathbb{E}(N) = \sum_{k=1}^n (-1)^{k+1} \sum_{\{i_1, \dots, i_k\}} \frac{1}{p_{i_1} + \dots + p_{i_k}}$
- $X \geq 0$ :  $\mathbb{E}[X] = \int_0^\infty \mathbb{P}(X > u) du$ ,  
 $\mathbb{E}[X] = \sum_{k=0}^\infty \mathbb{P}(X > k)$ .
- Order statistics  $(X_{(1)}, \dots, X_{(n)})$  has joint density  
 $n! f(x_1) \cdots f(x_n) \mathbf{1}_{\{x_1 < \dots < x_n\}}$
- $i$ th order statistic  $X_{(i)}$  has density  
 $i \binom{n}{i} f(x) F^{i-1}(x) \{1 - F(x)\}^{n-i}$ .
- $\mathbb{E}(X) = \mathbb{E}\{\mathbb{E}(X|Y)\}$ .
- $\text{var}(X) = \mathbb{E}\{\text{var}(X|Y)\} + \text{var}\{\mathbb{E}(X|Y)\}$ .
- Random sum  $S_N = X_1 + \dots + X_N$ :  
 $\mathbb{E}(S_N) = \mu \mathbb{E}(N)$ ,  $\text{var}(S_N) = \sigma^2 \mathbb{E}(N) + \mu^2 \text{var}(N)$ .

### Chapter 2

- Hitting time of  $i$ :  
 $\{\tau_i = n\} = \{X_n = i, X_{n-1} \neq i, \dots, X_1 \neq i\}$ .
- $f_{ii}^{(n)} = \mathbb{P}(\tau_i = n | X_0 = i)$ ,  
 $f_i = \sum_{n=1}^\infty f_{ii}^{(n)} = \mathbb{P}(\tau_i < \infty | X_0 = i)$ .
- $P_{ii}^n = \sum_{k=1}^n f_{ii}^{(k)} P_{ii}^{n-k}$ ,  $n \geq 1$ .
- $f_{jk}^{(n)} = \mathbb{P}\{X_n = k, X_{n-1} \neq k, \dots, X_1 \neq k | X_0 = j\}$ .
- $P_{jk}^n = \sum_{\ell=1}^n f_{jk}^{(\ell)} P_{kk}^{n-\ell}$ ,  $n \geq 1$ .
- $U_{jk}(s) = \sum_{n=0}^\infty P_{jk}^n s^n$ ,  $F_{jk}(s) = \sum_{n=1}^\infty f_{jk}^{(n)} s^n$ .
- $U_{jk}(s) - \mathbf{1}_{\{j=k\}} = F_{jk}(s) U_{kk}(s)$ .
- Gambler's probabilities of reaching a fortune  $N$ ,  
 $0 \leq i \leq N$  and  $r = q/p$ :  

$$p_i = \begin{cases} (1 - r^i)/(1 - r^N), & \text{if } p \neq 1/2 \\ i/N, & \text{if } p = 1/2 \end{cases}$$
Ruin probabilities:  $q_i = 1 - p_i$ .
- $s_{ij} = \mathbb{E}\left[\sum_{n=0}^\infty \mathbf{1}_{\{X_n=j\}} \middle| X_0 = i\right]$   
 $= \sum_{n=0}^\infty \mathbb{P}(X_n = j | X_0 = i) = \sum_{n=0}^\infty P_{ij}^n$ .
- $s_{ij} = \mathbf{1}_{\{i=j\}} + \sum_{k=1}^M P_{ik} s_{kj}$ , or in matrix form,  
 $S = I + P_T S$ . Equivalently,  $S = (I - P_T)^{-1}$ .
- $f_{ij} = \sum_{n=1}^\infty f_{ij}^{(n)} = \frac{s_{ij} - \mathbf{1}_{\{i=j\}}}{s_{jj}}$ ,  $i, j \in T$ .

### Chapter 3

- $\min\{X_1, \dots, X_n\} \sim \mathcal{E}(\lambda_1 + \dots + \lambda_n)$

- $\mathbb{P}\{X_1 < X_2\} = \frac{\lambda_1}{\lambda_1 + \lambda_2}$
- $\mathbb{E}\left[X_1 \mathbf{1}_{\{X_1 < X_2\}}\right] = \frac{\lambda_1}{(\lambda_1 + \lambda_2)^2}$
- $\mathbb{P}\{X_1 < X_2 < X_3\} = \frac{\lambda_1 \lambda_2}{(\lambda_2 + \lambda_3)(\lambda_1 + \lambda_2 + \lambda_3)}$
- $\mathbb{P}(R_n = i) = \frac{\lambda_i}{\lambda_1 + \dots + \lambda_n}$
- $\mathbb{P}(S_n^1 < S_m^2) = \sum_{k=n}^{n+m-1} \binom{n+m-1}{k} p^k (1-p)^{n+m-1-k}$

### Chapter 4

- Kolmogorov backward equation:  $P'(s) = GP(s)$ , where  
 $G_{ij} > 0$  for  $i \neq j$ , and  $G_{ii} = -\sum_{j \neq i} G_{ij}$ .
- $\lambda_i = -G_{ii}$  and  $J_{ij} = G_{ij}/\lambda_i$  for  $i \neq j$ , and  $J_{ii} = 0$ .
- $P(t) = e^{tG}$ , for  $t \geq 0$ .
- M/M/1 queue stationary distribution  $\pi(i) = (1 - \theta)\theta^i$ ,  
where  $\theta = \lambda/\mu < 1$ .
- Ergodic Theorem: with  $\{\pi_i\}$  the stationary distribution,  
 $\mathbb{P}\left\{\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T f(X(t)) dt = \sum_{i \in E} \pi_i f(i)\right\} = 1$ .

### Chapter 5

- $f(\mathbf{x}) = (2\pi)^{-\frac{p}{2}} |\Sigma|^{-\frac{1}{2}} \exp\left\{-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})' \Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu})\right\}$
- For  $\mathbf{X} = \begin{pmatrix} X_1 \\ X_2 \end{pmatrix} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$   
 $\boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}$ ,  $\Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix}$ ,  
 $\mathbf{X}_1 | \mathbf{X}_2 = \mathbf{x}_2 \sim \mathcal{N}(\boldsymbol{\mu}_{1|2}, \Sigma_{11|2})$ , where  
 $\boldsymbol{\mu}_{1|2} = \mathbb{E}(\mathbf{X}_1 | \mathbf{X}_2 = \mathbf{x}_2) = \boldsymbol{\mu}_1 + \Sigma_{12} \Sigma_{22}^{-1}(\mathbf{x}_2 - \boldsymbol{\mu}_2)$ ,  
 $\Sigma_{11|2} = \text{Var}(\mathbf{X}_1 | \mathbf{X}_2 = \mathbf{x}_2) = \Sigma_{11} - \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21}$ .
- For  $0 = t_0 < t_1 < \dots < t_n$ ,  
 $(B(t_1), \dots, B(t_n)) \sim \mathcal{N}(\mathbf{0}, \Sigma)$  where  $\Sigma_{ij} = t_i \wedge t_j$ .
- Let  $0 < t_1 \leq t_2$ .  
Forecasting:  $B(t_2) | B(t_1) = x_1 \sim \mathcal{N}(x_1, t_2 - t_1)$ .  
Backcasting:  $B(t_1) | B(t_2) = x_2 \sim \mathcal{N}\left(\frac{t_1}{t_2} x_2, \frac{t_1}{t_2}(t_2 - t_1)\right)$ .
- $T_a = \inf\{t : B(t) = a\}$
- Reflection principle:  $B_t^* = \begin{cases} B_t, & \text{if } t \leq T_a, \\ 2a - B_t, & \text{if } t > T_a. \end{cases}$
- Running maximum:  $M_t = \max_{0 \leq s \leq t} B(s)$ ,  $t \geq 0$ .
- For  $0 < b \leq a$ ,  $\mathbb{P}(M_t \geq a, B_t \leq b) = \mathbb{P}(B_t \geq 2a - b)$ .
- $M_t$  has the same distribution as  $|B_t|$ .
- $T_a$  has the same distribution as  $\frac{a^2}{B_1^2}$

### Some common distributions

	pdf/pmf	mean	variance
Geometric	$(1-p)^{n-1}p$	$\frac{1}{p}$	$\frac{1-p}{p^2}$
Poisson	$e^{-\lambda} \frac{\lambda^k}{k!}$	$\lambda$	$\lambda$
Exponential	$\theta e^{-\theta x}$	$\frac{1}{\theta}$	$\frac{1}{\theta^2}$
Gamma	$\frac{\theta^k x^{k-1}}{\Gamma(k)} e^{-\theta x}$	$\frac{k}{\theta}$	$\frac{k}{\theta^2}$