


71028 v2

Catalogue of Salmon Streams
and Spawning Escapements
of Sub-Districts 110 and 120
(Yukon-Arctic)



M.J. Hancock and D.E. Marshall

Department of Fisheries & Oceans
Enhancement Services Branch
1090 West Pender Street
Vancouver, B.C. V6E 2P1

October 1984

Canadian Data Report of
Fisheries and Aquatic Sciences
No. 474



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Data Report of Fisheries and Aquatic Sciences

Data reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of data reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was introduced with the publication of report number 161.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport statistique canadien des sciences halieutiques et aquatiques

Les rapports statistiques servent à classer et à archiver les compilations de données pour lesquelles il y a peu ou point d'analyse. Ces compilations auront d'ordinaire été préparées à l'appui d'autres publications ou rapports. Les sujets des rapports statistiques reflètent la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports statistiques ne sont pas destinés à une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans autorisation écrite préalable de l'établissement auteur. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 25 de cette série ont été publiés à titre de relevés statistiques, Services des pêches et de la mer. Les numéros 26 à 160 ont été publiés à titre de rapports statistiques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 161.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Fisheries and Oceans
Canadian Data Report of
Fisheries and Aquatic Sciences

No. 474
October 1984

CATALOGUE OF SALMON STREAMS AND SPAWNING ESCAPEMENTS OF

SUB - DISTRICTS 110 AND 120
(Yukon - Arctic)

by

M. J. Hancock and D. E. Marshall

Fisheries and Oceans
Enhancement Services Branch
1090 West Pender Street
Vancouver, B.C.
V6E 2P1

Minister of Supply and Services Canada 1984
Cat. NO. Fs 97-13/474 ISSN 0706-6465

Correct citation for this publication:

Hancock, M.J. and D.E.Marshall. 1984 Catalogue of Salmon Streams
and Spawning Escapements of Sub-Districts 110 and 120 (Yukon-Arctic)
Can. Data Rep. Fish and Aquat. Sci. 474 xvi + 233p.

ABSTRACT

Hancock, M.J. and D.E.Marshall, 1984. Catalogue of Salmon Streams and Spawning Escapements of Sub-Districts 110 and 120 (Yukon-Arctic) Can. Data Rep. Fish and Aquat. Sci. 474: XVI + 233p.

Catalogue containing each stream's location, spawning distribution, barriers and points of difficult ascent, escapement records and other general data pertaining to the stream. The catalogue also includes a topographical map of the stream's location and in some cases a sketch which further describes the surrounding area.

Keywords: Yukon-Arctic, Sub-Districts 110 and 120, Salmon Streams, Spawning Escapements.

RESUME

Hancock, M.J. and D.E.Marshall, 1984. Catalogue of Salmon Streams and Spawning Escapements of Sub-Districts 110 and 120 (Yukon-Arctic) Can. Data Rep. Fish and Aquat. Sci. 474: XVI + 233p.

Le présent répertoire donne l'emplacement de chaque cours d'eau, la répartition de fraie, les points de remonte difficile, les données sur les saumons de remonte et d'autres information générales concernant le cours d'eau. On y trouve aussi une carte topographique de l'emplacement du cours d'eau et, dans quelque cas, un croquis décrivant la zone environnante.

Mots-clés: Yukon-Arctique, sous-secteur 110 and 120, cours d'eau à saumons, remonte.

ADDITIONAL REFERENCES

For additional information on the fisheries resources of the Yukon area, please refer to the following bibliography:

Ennis, G.L., A. Cinader, S. McIndoe, and T. Munsen. 1982.
An annotated bibliography and information summary on
the fisheries resources of the Yukon River Basin in Canada.
Can. MS Rept. Fish Aquat. Sci. 1657: V + 278 p.

CONTENTS

	<u>Page</u>
Abstract/Resume	iii
Additional References	iv
Table of Contents	v, vi
Statistical Areas	vii, viii
Management Divisions	ix
Federal Fisheries Districts and Sub-Districts	x
Standards used on stream data page	xi, xii
Map References	xiii
Federal Fisheries District and Sub-District Addresses	xiv
Total Escapement Record for Sub-District 110 Yukon Arctic	xv
Index to Map of Salmon Spawning Streams of Sub-Districts 110 Yukon Arctic	xvi

STREAM DATA

(1) BIG CREEK	1
(2) BIG SALMON RIVER	5
(3) SOUTH BIG SALMON RIVER	9
(4) BLIND CREEK	13
(5) CHANDINDU RIVER	17
(6) DONJEK RIVER	21
(7) EARN RIVER	25
(8) FIFTEEN MILE RIVER	29
(9) FISHING BRANCH RIVER	33
(10) FORTIN CREEK	39
(11) FORTY MILE CREEK	43
(12) GLADYS RIVER	47
(13) GLENYON RIVER	51
(55) HAYES CREEK	55
(14) HOOLE RIVER	57
(15) HUSKY DOG CREEK	61
(16) JANET CREEK	65
(17) JENNINGS RIVER	69
(18) KALZAS RIVER	73
(19) KLONDIKE RIVER	77
(20) KLUANE LAKE	81
(21) KLUANE RIVER	85
(22) KOIDERN RIVER	89
(23) LAPIE RIVER	93
(24) LEWIS LAKE	97
(25) LITTLE KALZAS RIVER	101
(26) LITTLE SALMON RIVER	105, 107
(27) McNEIL RIVER	113
(28) McQUESTEN RIVER	117
(29) MAYO RIVER	121

	<u>Page</u>
(30) MICHIE CREEK	127
(31) MINER RIVER	131
(32) MORLEY RIVER	135
(33) NISUTLIN RIVER	139
(34) NORDENSKIOLD RIVER	143
(35) NORTHERN LAKE	147
(36) (Ollie River and Lake)	151
(37) PELLY LAKES	155
(38) PELLY RIVER	159
(39) PLEASANT CREEK	163
(40) RED RIVER	167
(41) RIDDEL RIVER (South McMillan River)	171
(42) ROSS RIVER	175
(43) RUSSEL CREEK	179
(44) SHELDON LAKE	183
(45) SMART RIVER	187
(46) STEWART RIVER	191
(47) SWEDE JOHNSON CREEK	195
(48) SWIFT RIVER	199
(49) TAKHINI RIVER	203, 205
(50) TATCHUN RIVER	209
(51) TESLIN RIVER	213
(52) (Upper Teslin River)	217
(53) WOLF RIVER	221
(54) YUKON RIVER	225, 227, 229

READ CAREFULLY

1. PIN UP IN WHEELHOUSE.
2. WHEN DELIVERING YOUR CATCH, GIVE TALLY MAN THE MAP NUMBER, OR NUMBERS SHOWING THE AREA IN WHICH YOUR FISH WERE CAUGHT.
3. ACCURATE CATCH REPORTS WILL HELP PRESERVE YOUR FISHERIES.
4. FOR COMPLETE DETAILS, CONSULT BRITISH COLUMBIA FISHERIES REGULATIONS.

— STATISTICAL AREAS ARE DIVIDED BY RED LINES

— SALMON FISHING WITH NETS OF ANY KIND IS NOT PERMITTED OUTSIDE OF — THAT IS SEAWARD OF — THE HEAVY BLACK LINE.


● FISHERIES SERVICES OFFICES

DEPARTMENT OF THE ENVIRONMENT FISHERIES SERVICE


STATISTICAL MAP

SHOWING AREAS OF CATCH FOR
BRITISH COLUMBIA WATERS
(SOUTHERN HALF)

0 10 20 30 40 50
SCALE IN MILES



INSERT MAP - FRASER RIVER



FOR ALL FISH CAUGHT
OFF THE COAST OF THE
STATE OF WASHINGTON
SOUTH OF CAPE FLATTERY
REPORT AS AREA "C"

SHOWING AREAS OF CATCH FOR
BRITISH COLUMBIA WATERS
(NORTHERN HALF)

 FISHERIES SERVICE OFFICES

NOTE AREAS 5,6,7,8 AND 9 REVISED JAN. 1974

This map illustrates the Graham Island area, a large island in the Hecate Strait. The island is oriented vertically, with its northern tip at the top. Key features include:

- Islands and Inlets:**
 - Frederick I.** and **Thian-Hd.** are located at the northern tip.
 - Shag Rk.** and **Wiah Pt.** are on the northern coast.
 - Naden Harbour** and **Masset** are on the western coast.
 - Masset Inlet** and **Port Clements** are on the eastern coast.
 - Chippa Is.** is located south of the northern tip.
 - Graham Island** is the central, largest island.
 - Skidegate Inlet** and **Sandspit** are on the eastern coast of Graham Island.
 - Gray Pt.** is on the eastern coast.
 - Curmewah Inet** is on the eastern coast.
 - Chads R.** is on the western coast.
 - Tasou Sound** and **Sunday Harbour** are on the western coast.
 - McLean-Fraser Pt.** is on the southern coast.
 - Cape St James** is at the southern tip.
- Surrounding Regions:**
 - 2 WEST** and **2 EAST** are labeled on the western and eastern sides of the island.
 - 4** and **5** are labeled in the upper right corner.
 - Butterworth Rk.** and **Rose Pt.** are on the northern coast.
 - Stephens Is.** is on the northern coast.
 - Bonilla Is.** is on the eastern coast.
 - Scudder Pt.** is on the eastern coast.
 - Garcin Rks.** and **Lyman (East) Pt.** are on the eastern coast.
 - Kitgoro Pt.** is on the western coast.
 - Moresby** is on the western coast.
 - Green Charl. Pt.** is on the western coast.
 - Kitgoro Pt.** is on the western coast.
 - Kitgoro Pt.** is on the western coast.
- Water Bodies:**
 - Hecate Strait** is the body of water to the east of the island.
 - Kitgoro Pt.** is on the western coast.
 - Kitgoro Pt.** is on the western coast.
 - Kitgoro Pt.** is on the western coast.

1. PIN UP IN WHEELHOUSE
2. WHEN DELIVERING YOUR CATCH, GIVE TALLY MAN THE MAP NUMBER, OR NUMBERS SHOWING THE AREA IN WHICH YOUR FISH WERE CAUGHT
3. ACCURATE CATCH REPORTS WILL HELP PRESERVE YOUR FISHERIES.

30

AREA NORTH

AREA SOUTH

CAPE SCOTT

VANCOUVER
ISLAND

12

FIFTH EDITION JAN 1974

**MANAGEMENT DIVISIONS
FISHERIES & OCEANS
PACIFIC REGION**

Yukon River
Alsek R.
Mackenzie River
Liard R.
Peace R.
Skeena River
Fraser River
Columbia River

WHITEHORSE
PRINCE RUPERT
PRINCE GEORGE
KAMLOOPS
VANCOUVER
VICTORIA

Fraser River, Northern B.C. & Yukon Division

North & Central Coast Division

South Coast Division

ix

**MANAGEMENT DIVISIONS
FISHERIES & OCEANS
PACIFIC REGION**

Yukon River
Alsek R.
Mackenzie River
Liard R.
Peace R.
Skeena River
Fraser River
Columbia River

WHITEHORSE
PRINCE RUPERT
PRINCE GEORGE
KAMLOOPS
VANCOUVER
VICTORIA

Fraser River, Northern B.C. & Yukon Division

North & Central Coast Division

South Coast Division

ix

**MANAGEMENT DIVISIONS
FISHERIES & OCEANS
PACIFIC REGION**

Fraser River,
Northern B.C.
& Yukon
Division

North & Central
Coast Division

South Coast
Division

Yukon River
Alsek R.
Mackenzie River
Liard R.
Stikine River
Skeena River
Peace R.
Fraser River
Columbia River

WHITEHORSE
PRINCE RUPERT
PRINCE GEORGE
KAMLOOPS
VANCOUVER
VICTORIA

ix

**MANAGEMENT DIVISIONS
FISHERIES & OCEANS
PACIFIC REGION**

Fraser River,
Northern B.C.
& Yukon
Division

North & Central
Coast Division

South Coast
Division

Yukon River
Alsek R.
Mackenzie River
Liard R.
Stikine River
Skeena River
Peace R.
Fraser River
Columbia River

WHITEHORSE
PRINCE RUPERT
PRINCE GEORGE
KAMLOOPS
VANCOUVER
VICTORIA

ix

**MANAGEMENT DIVISIONS
FISHERIES & OCEANS
PACIFIC REGION**

Fraser River,
Northern B.C.
& Yukon
Division


North & Central
Coast Division

South Coast
Division

Yukon River
Alsek R.
Mackenzie River
Liard R.
Stikine River
Skeena River
Peace R.
Fraser River
Columbia River

WHITEHORSE
PRINCE RUPERT
PRINCE GEORGE
KAMLOOPS
VANCOUVER
VICTORIA

ix



BRITISH COLUMBIA

FISHERIES & OCEANS - Pacific Region

DISTRICT/SUB-DISTRICT OFFICES

DISTRICT/SUB-DISTRICT	ADDRESS	PHONE	SEN-DISTRICT
DISTRICT 1 - Kamloops	202 - 1117 Broadway St., Kamloops, B.C. V2S 2G9	274-4322	246
DISTRICT 2 - New Westminster	Box 1100, 441 Healy Ave., New Westminster, B.C. V3M 2V5	832-8027	246
DISTRICT 3 - Nanaimo	2700 Quilley Road, Prince Rupert, B.C. V8P 2G5	544-1030	246
DISTRICT 4 - Port Alberni	Box 610, Clearwater, B.C. V6M 1A0	474-2431	246
DISTRICT 5 - Victoria	Box 105, Lillooet, B.C. V3V 1G9	256-5025	246
DISTRICT 6 - Vancouver	Box 436, Quilley, V2Z 3V7	990-4544	246
DISTRICT 7 - Whitehorse	Box 100, Whitehorse, B.C. Y1A 1A1	930-4544	246
DISTRICT 8 - Prince George	Box 100, Prince George, B.C. V2N 1A1	930-4544	246
DISTRICT 9 - Hazelton	Box 100, Hazelton, B.C. V2A 1A1	930-4544	246
DISTRICT 10 - Bulkley Valley	Box 100, Bulkley Valley, B.C. V2A 1A1	930-4544	246
DISTRICT 11 - Skeena	Box 100, Skeena, B.C. V2A 1A1	930-4544	246
DISTRICT 12 - Fraser	Box 100, Fraser, B.C. V2A 1A1	930-4544	246
DISTRICT 13 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 14 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 15 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 16 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 17 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 18 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 19 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 20 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 21 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 22 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 23 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 24 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 25 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 26 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 27 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 28 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 29 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 30 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 31 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 32 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 33 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 34 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 35 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 36 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 37 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 38 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 39 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 40 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 41 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 42 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 43 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 44 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 45 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 46 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 47 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 48 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 49 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 50 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 51 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 52 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 53 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 54 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 55 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 56 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 57 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 58 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 59 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 60 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 61 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 62 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 63 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 64 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 65 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 66 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 67 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 68 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 69 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 70 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 71 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 72 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 73 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 74 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 75 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 76 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 77 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 78 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 79 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 80 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 81 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 82 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 83 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 84 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 85 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 86 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 87 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 88 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 89 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 90 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 91 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 92 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 93 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 94 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 95 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 96 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 97 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 98 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 99 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246
DISTRICT 100 - Bulkley	Box 100, Bulkley, B.C. V2A 1A1	930-4544	246

HABITAT MANAGEMENT OFFICES

Vanouver	1095 West Pender Street, Vancouver, B.C. V2T 1A1	666-4544
New Westminster	311 - 545 Columbia Street, New Westminster, B.C. V3M 1A1	524-7144
Nanaimo	400 Front Street, Nanaimo, B.C. V9S 5G7	753-7248
Prince Rupert	202 - 477 2nd Ave. West, Prince Rupert, B.C. V8P 1G5	424-9393

LEGEND

- DISTRICT BOUNDARY
- SUBDISTRICT BOUNDARY
- DISTRICT OFFICE
- SUBDISTRICT OFFICE

Habitat Management Division
Department of Fisheries and
Oceans

STANDARDS USED ON STREAM DATA PAGE

Name of Stream: Name as given in Gazetteer of Canada: Yukon Territory, Fourth Edition, Dept. of Energy, Mines and Resources, Ottawa, 1981; local or non-gazetted names are added in lower case type.

Districts and Sub-Districts: As shown on map produced by Habitat Management Division, D.F.O., June 1983.

Map References: Most of the maps used in this publication are the 1:250,000 scale, this being the largest scale available for complete map coverage of the Yukon Territory in the National Topographical System. Some of the maps are smaller scale, 1:500,000 and 1:1,000,000, these are indicated on the map labels:

1: 250,000 4cm = 10km
1: 500,000 4cm = 20km
1: 1,000,000 4cm = 40km

Geographic Co-ordinate Reference: The geographic co-ordinates given for the streams indicate the position of the mouth, and the co-ordinates are to the nearest minute.

Length: The portion of the stream accessible to spawning salmon.

Drainage: Area in square kilometers of the entire drainage basin feeding the stream.

Discharge: Extremes of maximum and minimum daily discharge for period of record.

Discharge and drainage data has been taken from the "Historical Streamflow Summary" Yukon and Northwest Territories to 1982, Inland Waters Directorate, Water Resources Branch, Water Survey of Canada, Ottawa, 1983.

STANDARDS USED ON STREAM DATA PAGE

Barriers and Points of Difficult Ascent: Complete and partial barriers to salmon and their distance from the stream mouth. Species likely to be affected may be listed. Both natural and man-made obstructions are defined.

Spawning Distribution: Portion of the stream utilized by each species. Distribution is indicated by brief comments opposite the species.

General Remarks: Emphasizes features of stream and spawning populations. Also includes industrial activity, routes of accessibility, etc. The comments with dates are taken from the "Annual Reports of Salmon Streams and Spawning Grounds". In some cases, references to additional information not included in the General Remarks may be given.

Escapement Records: The escapement represents the mid point of the coded range of escapement for each species. For example: the letter "H" representing 5000-10000 fish would be entered as 7500. Where absolute numbers are provided by Fisheries Personnel, these numbers are entered. N/O means the stream was inspected but no fish were observed; UNK means there was evidence of fish present but no estimates were made; NO RECORDS means no escapement records for the applicable years could be found in the escapement files. N/I means the stream was not inspected.

Timing: Dates when salmon arrive in the stream, begin to spawn, reach peak spawning period and finish spawning.

E = early (1st to 10th of the month)
M = mid (11th to 20th of the month)
L = late (21st to end of the month)

MAP REFERENCES

Roads:

hard surface, all weather	more than 2 lanes
hard surface, all weather	2 lanes
loose surface, all weather	2 lanes wide or more
"	less than 2 lanes
Private Road, Trail	Private Road Trail

Railways:

normal gauge, multiple track	Station
normal gauge, single track	Stop
abandoned, or under construction	Siding
narrow gauge, single track	
Bridge, underpass or overpass	
Tunnel	

Boundary, International

" Province	
" County or District	
" Township or Parish	
" City or Town	
" Reservation, Indian, Military, etc	
Power Transmission Line	
Telephone or Telegraph, trunk route	
Horizontal Control Point	
Boundary Marker	
Bench Mark	BM
Spot Elevation, (in feet)	2975
Mine or Pit	5752

Road, Hard Surface, All Weather

" Loose Surface, All Weather	2 Lanes
" Loose Surface, Less than 2 lanes	All Weather Dry Weather
" Private (Logging, Mining etc)	
" Four Wheel Drive	
Trail	
Railway	
Main Telephone Line	
Main Electric Power Line	
Horizontal Control Station	
Contours (Interval 500 feet)	5000
Elevation in feet above mean sea-level	2584' 6312'
Intermittent Stream	
Swamp or Marsh	
Dam	
Spring	
Navigation Light	
Mine	
Glacier	
Customs Office	

House, Building

School	S
Church	+
" with conspicuous Tower or Spire	+
Post Office	P
Tower, Radio Mast, Lookout, etc	o
Cemetery	Cem
Quarry	
Sand or Gravel Pit	
Cliff	
Cutting	
Embankment	
Saw Mill	SM

Lighthouse

Wharf or Pier	
Foreshore Flats	
Swamp or Marsh	
Lake or Pond, intermittent	
Glacier or Snowfield	
Stream, intermittent	
Irrigation Canals, Ditches	
Inundated Land, seasonal	
Contours, elevation	500 400
" depression	500 400
" approximate	500 400
Forest, unclassified	

Surveyed timber license number

Lot number	TL 2841
Building	L 124 or S 66
School	
Non-perennial stream	
Marsh or Swamp	
Glacier	
Foreshore flats	
Contours, elevation	500
Contours, depression	
Forest	

City or large town

Town	
Village or settlement	
Streams:	
intermittent or dry	
indefinite	
Irrigation canal or ditch	
Rapids, falls	
Aerodrome	
Landing ground	
Post office	P
School	S
Church	+
Intermittent lake	
Marsh or swamp	
Sand, gravel or mud	
Wooded areas	
Seaplane base	
Seaplane anchorage	
Boundary monument	
Astronomical position	
Horizontal control point	

Streams

Highways	
Roads	
Trails	
Houses	
Railroad	
Falls	
Rapids	
Rip-Rap	
Bridges	
Dam	
Log Jams	
Log	
Power Line	
Coho	CO
Chum	CM
Pink	P
Chinook	CK
Sockeye	S

FISHERIES & OCEANS - Pacific Region

DISTRICT/SUB-DISTRICT OFFICES

<u>DISTRICT/SUB-DISTRICT</u>	<u>ADDRESS</u>	<u>TELEPHONE</u>	<u>SUB-DISTRICT NUMBER</u>
<u>DISTRICT #1 - Kamloops</u>	202 - 317 Seymour St., Kamloops, V2C 2E9	374-4322	
Salmon Arm	Box 1160, 461 Beatty Ave. NW, Salmon Arm V0E 2T0	832-8037	29K
Prince George	2392 Ospika Blvd., Prince George, V2N 3N5	564-7030	29I
Clearwater	Box 610, Clearwater, V0E 1N0	674-2633	29J
Lillooet	Box 315, Lillooet, V0K 1V0	256-4525	29F
Quesnel	Box 4340, Quesnel, V2J 3J3	992-2434	29H
Williams Lake	540 Borland St, Williams Lake, V2G 1R9	398-6544	29G
<u>DISTRICT #2 - New Westminster</u>	309 - 549 Columbia St., New West., V3L 1B3	524-7181	
Vancouver	309 - 549 Columbia St., New West., V3L 1B3	524-7306	28A
Surrey	309 - 549 Columbia St., New West., V3L 1B3	524-7171	29B
Coquitlam	309 - 549 Columbia St., New West., V3L 1B3	524-7169	29C
Steveston	1255 No. 1 Road, Richmond, V7E 1T7	274-7217	29A
Squamish	Box 85, Squamish, V0N 3G0	892-3230	28B
Mission	Box 3308, Mission, V2V 4J5	826-3664	29D
Chilliwack	Suite 5, 9375 Mary St., Chilliwack, V2P 4G9	792-6011	29E
<u>DISTRICT #3 - Nanaimo</u>	60 Front St., Nanaimo, V9R 5H7	754-3257	
Nanaimo/Ladysmith	60 Front St., Nanaimo, V9R 5H7	754-3257	17
Qualicum Beach	Box 1270, Qualicum Beach, V0R 2T0	752-9712	14S
Comox	Box 1328, Comox, V9N 3Z0	339-2031	14N
Duncan	Box 241, 191 Ingram St., Duncan, V0L 3X3	746-6221	18
Powell River	4488 Marine Avenue, Powell River, V8A 2K2	485-9621	15
Pender Harbour	Box 10, Madeira Park, V0N 2H0	883-2313	16
<u>DISTRICT #4 - Port Alberni</u>	Box 280, Federal Building, Port Alberni, V9Y 7M7	724-0195	
Port Alberni	Box 280, Federal Building, Port Alberni, V9Y 7M7	724-0195	23
Quatsino Sound	Box 10, Port Hardy, V0N 2P0	949-6422	27
Kyuquot	Box 549, Tahsis, V0P 1X0	934-6606	26
Tahsis	Box 549, Tahsis, V0P 1X0	934-6606	25
Tofino	Box 48, Tofino, V0R 2Z0	725-3468	24
<u>DISTRICT #5 - Campbell River</u>	215 - 950 Alder St., Campbell River, V0W 2P8	287-2102	
Campbell River	215 - 950 Alder St., Campbell River, V0W 2P8	287-2102	13
Seymour Inlet	Box 10, Port Hardy, V0N 2P0	949-6422	11
Alert Bay	Box 10, Alert Bay, V0N 1A0	974-5216	12
<u>DISTRICT #6 - Victoria</u>	116 - 816 Government St., Victoria, V8W 1W9	566-3252	
Victoria/Saanich	116 - 816 Government St., Victoria, V8W 1W9	566-3252	19
Sooke	Box 460, Sooke, V0Z 1N0	642-5322	20
<u>DISTRICT #7 - Kitimat</u>	315 - 450 Federal Building, Kitimat, V8C 1T6	632-4884	
Butedale	315 - 450 Federal Building, Kitimat, V8C 1T6	632-4884	6
Bella Bella	Box 38, Bella Bella, V0T 1B0	957-2363	7
Bella Coola	Box 130, Bella Coola, V0T 1C0	799-5345	8
Rivers Inlet	Dawson Landing P.O., Rivers Inlet, V0N 1M0		9
Smith Inlet	Dawson Landing P.O., Rivers Inlet, V0N 1M0		10
<u>DISTRICT #8 - Prince Rupert</u>	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	
Waterfront	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	
Skeena	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	4A
Greenville - Principe	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	5
Lower Nass	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	3A
Upper Nass	Box 29, Nass Camp, V0J 3J0	633-2408	3B
Hazelton	Box 327, Field Street, Hazelton, V0J 1Y0	842-6327	4C
Smithers	Box 578, Smithers, V0J 2N0	847-2312	4D
Terrace	4721-B Lazelle Ave., Terrace, V8G 1R6	635-2206	4B
<u>DISTRICT #9 - Queen Charlotte Is.</u>	Box 99, Queen Charlotte City, V0T 1S0	559-4413	
West Coast Q.C.I.	Box 99, Queen Charlotte City, V0T 1S0	559-4413	2W
Masset	Box 99, Masset, V0T 1M0	626-3316	1
Sandspit	Box 222, Sandspit, V0T 1T0	637-5340	2E
<u>DISTRICT #10 - Whitehorse</u>	122 Industrial Road, Whitehorse, Y.T., Y1A 2T9	667-2235	
Yukon South/Northern B.C.	122 Industrial Road, Whitehorse, Y.T., Y1A 2T9	667-2235	120
Yukon-Arctic	122 Industrial Road, Whitehorse, Y.T., Y1A 2T9	667-2235	110
Alsek-Taku	Box 5341, Haines Junction, Y.T., Y0B 1L0	634-2235	130


TOTAL ESCAPEMENT RECORD FOR SUB-DISTRICT 110 - YUKON ARCTIC

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		10,325		2,800		
67		11,725		3,525		
68		18,900		3,525		
69		4,306		7,525		
70		16,450				
71		5,975	25	79,250		
72		4,033		500		
73		415	25	18,500		
74		25,758	300	58,400		
75		2,084		357,200		
76		7,885		15,020		
77		46,093		32,300		
78		93,551		10,200		
79		62,975		215,100		
80		80,750	5,400	60,300		
81		60,143		106,550		
82		4,464		16,870		
83						
84						
85						
TOTALS		455,832	5,750	987,565		


TIMING

ARRIVE						
START						
PEAK						
END						

REMARKS _____



- (1) BIG CREEK
- (2) BIG SALMON RIVER
- (3) SOUTH BIG SALMON RIVER
- (4) BLIND CREEK
- (5) CHANDINDU RIVER
- (6) DONJEK RIVER
- (7) EARN RIVER
- (8) FIFTEEN MILE RIVER
- (9) FISHING BRANCH RIVER
- (10) FORTIN CREEK
- (11) FORTY MILE CREEK
- (12) GLADYS RIVER
- (13) GLENLYON RIVER
- (14) HAYES CREEK
- (15) HOOLE RIVER
- (16) HUSKY DOG CREEK
- (17) JANET CREEK
- (18) JENNINGS RIVER
- (19) KALZAS RIVER
- (20) KLONDIKE RIVER
- (21) KLUANE LAKE
- (22) KLUANE RIVER
- (23) KOIDERN RIVER
- (24) LAPTE RIVER
- (25) LEWIS LAKE
- (26) LITTLE KALZAS RIVER
- (27) LITTLE SALMON RIVER
- (28) McNEIL RIVER
- (29) McQUESTEN RIVER
- (30) MAYO RIVER
- (31) MICHIE CREEK
- (32) MINER RIVER
- (33) MORLEY RIVER
- (34) NISUTLIN RIVER
- (35) NORDENSKIOLD RIVER
- (36) NORTHERN LAKE
- (37) (Ollie River and Lake)
- (38) PELLY LAKES
- (39) PELLY RIVER
- (40) PLEASANT CREEK
- (41) RED RIVER
- (42) RIDDEL RIVER (South McMillan River)
- (43) ROSS RIVER
- (44) RUSSEL CREEK
- (45) SHELTON LAKE
- (46) SMART RIVER
- (47) STEWART RIVER
- (48) SWEDE JOHNSON CREEK
- (49) SWIFT RIVER
- (50) TAKHINI RIVER
- (51) TATCHUN RIVER
- (52) TESLIN RIVER
- (53) (Upper Teslin River)
- (54) WOLF RIVER
- (55) YUKON RIVER



NAME OF STREAM BIG CREEKCONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110LOCATION OF MOUTH Flows into Yukon RiverPOSITION 62° 37' 137° 00'LENGTH _____ km WIDTH _____ m DRAINAGE 1750 km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²DISCHARGE (m³/s) Near the mouth max 206 Jul 15 1976 min 0.020B Apl. 4 1978

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1980 Flown by fixed wing aircraft. Totally frozen over. Nov. 11
1981 31st September Lower reaches of river flown while flying the Yukon River
for chum salmon, could have been too early.

ESCAPEMENT RECORD FOR BIG CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82					UNK	
83					Not Inspected	
84						
85						

TIMING


ARRIVE

START

PEAK

END

REMARKS



NAME OF STREAM BIG SALMON RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 61° 53' 134° 55'
 LENGTH 145 km WIDTH m DRAINAGE 6760 km²
 COMPOSITION: BEDROCK BOULDER COARSE FINE
 SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s) Near Carmacks max 671 Jun. 23 1962 min 11.0B Apl. 2 1971

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Usually no obstructions to headwaters
approx. 145 km.
1972 Log jam 1.6km from lake.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- heaviest concentration near lake — and in pockets throughout.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS 1967 Improvement of Canel Road permits easy public access to this stream. Many fishing parties and hunters in area during past season.
1969 High brackish water precluded accurate count.
1973 Aerial survey showed only 15 Chinook. Probably a little late for this river. No reasonable estimate can be made for river this year.
1974 River flown by Alaskans in early September and they located only 40 chinook. River flown later by our personnel and 150 chinook were sighted. Also numerous redds. Weather unfavourable for aerial inspection.
1975 800 chinook counted from air. Probably conservative as count was made fairly late in year. (end of August)

continued.....

BIG SALMON RIVER

- 1977 Flown Aug 25, 305 live fish and 11 carcasses observed.
- 1978 Aerial surveys by both Americans and Canadians. It is believed that both surveys were done after peak spawning.
Americans Aug. 18 1150
Canadians Aug. 22 225
- 1979 American and Canadian Surveys done together.
Conducted Aug. 18. Weather warm — broken cloud. It appears that the survey was late — there were redds with no fish on them as well as large redds with small numbers of fish. 637 chinook — 37 carcasses.
- 1980 Flown by U.S. Fish and Game Branch. Judging from the number of carcasses and redds, it was felt that survey was flown near the peak of spawning.
* Count was 1568 on grounds.
- 1981 Flown by A.D.F.G. Salmon were just spawning and there was evidence of many redds.
- 1982 Flown by A.D.F.G. Large number of jacks observed.

Light predation by bears and eagles.

*Breakdown:

Big Salmon Lake to Scurvey Creek	470
Scurvey Cr. to Pleasant Lake Cr.	288
Pleasant Lake Cr. to Bat Creek .	559
Bat Creek to Sauch Creek.	119
Sauch Creek to Next North Cr.	132

ESCAPEMENT RECORD FOR BIG SALMON RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		2,500				
67		3,500				
68		3,500				
69		1,500				
70		3,500				
71		400				
72		750				
73		15				
74		150				
75		800				
76		86				
77		316				
78		1,500				
79		1,200				
80		3,000				
81		2,573				
82		1,081				
83						
84						
85						


TIMING

ARRIVE		M Jul-E Aug			
START		L Jul-E Aug			
PEAK		E Aug-M Aug			
END		L Aug-M Sept			

REMARKS 1973 Late inspection. Not an accurate count.
 1982 Large number of Jacks observed.

1971 Sex ratios M 60% F 40%

1972 Sex ratios M 50% F 50%



NAME OF STREAM SOUTH BIG SALMON RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Big Salmon River

POSITION 61° 36' 134° 26'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1981 Aug.12 and Aug.20. River flown by Alaska Fish and Game on
August 12th. Water was too turbid due to mining on Livingstone Creek for a count.
Flown by fixed wing on Aug. 20th, but no salmon were seen. A report from a prospect-
or in the area states that there were a considerable number of chinook in the system.
No numbers were given.


ESCAPEMENT RECORD FOR SOUTH BIG SALMON RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81		N/O				
82		UNK				
83						
84						
85						

TIMING

ARRIVE						
START						
PEAK						
END						

REMARKS 1981 Local resident reported considerable number of chinook in system.
Numbers unknown.



BLIND CREEK

NAME OF STREAM _____
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Pelly River

POSITION 62° 11' 133° 13'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	_____
0.25 - 0.50	_____
0.50 - 0.75	_____
0.75 - 1.00	_____
> 1.00	_____

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	_____
CHINOOK	_____
COHO	_____
CHUM	_____
PINK (ODD YEAR)	_____
PINK (EVEN YEAR)	_____
STEELHEAD	_____

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1975 Local reports of Chinook being present in this creek.

Several Chinook caught by Faro residents.

1981 Locals report that there are several people in this area who are taking over their catch limit.

1982 Sportsfishermen report that the run was very poor this year. Survey done by boat by Yukon River Basin study.


ESCAPEMENT RECORD FOR BLIND CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75		N/O				
76		N/I				
77		-				
78		-				
79		-				
80		-				
81		N/I				
82		6				
83						
84						
85						

TIMING

ARRIVE		Jul				
START		Jul				
PEAK		Aug				
END		Sept				

REMARKS 1982 Survey done by boat and by Yukon River Basin Study.



NAME OF STREAM CHANDINDU RIVERCONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110LOCATION OF MOUTH Flows into Yukon RiverPOSITION 64° 15' 139° 43'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

>1.00

WETTED AREA _____ m²SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION UNKNOWN

SPECIES

SECTION OF STREAM USED

SOCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1980 - Helicopter survey. River was frozen over with open patches of good spawning grounds.

ESCAPEMENT RECORD FOR CHANDINDU RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80				N/O		
81						
82						
83						
84						
85						

TIMING


ARRIVE

START

PEAK

END

REMARKS



0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

BARRIERS OR POINTS OF DIFFICULT ASCENT

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	- in Koidern River downstream of Koidern River Lodge.
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

Light predation. Normal seasonal fluctuations in water levels.

ESCAPEMENT RECORD FOR DONJEK RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79				40,000		
80				100		
81						
82						
83						
84						
85						

TIMING

ARRIVE				Oct		
START				Oct		
PEAK				Oct		
END				Nov		

REMARKS Inspected by Foothills Pipeline Personnel


Earn River

N

0

10

kilometres



NAME OF STREAM EARN RIVER

CONSERVATION DISTRICT 10 SUB-DISTRICT 110

LOCATION OF MOUTH Flows into Pelly River

POSITION 62° 44' 134° 42'

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT 1972 reported passable log jams throughout.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- scattered over 11km of stream up to lake.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS A small narrow river -- difficult to survey from the air.

Narrow canyon in lower reaches .8km from mouth.

1967 Game guides reported numerous chinook.

1969 Heavy rainfall and early snowstorms in August prevented access to this area
on one occasion, and high water precluded any sightings on the second investi-
gation.

1970 Effects of erosion evident in water due to runoff from a large burn.

1981 Water was rather organic brown making observation difficult. Mining camp
has been set up at lake.

Predators: Grizzly bears.


ESCAPEMENT RECORD FOR EARN RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		1,250				
67		1,500				
68		400				
69		N/I				
70		750				
71		75				
72		75				
73		N/I				
74		N/I				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/I				
81		100				
82		N/I				
83						
84						
85						

TIMING

ARRIVE		L Aug-E Aug				
START		L Jul-E Aug				
PEAK		E Aug-M Aug				
END		L Aug- Sept				

REMARKS



NAME OF STREAM FIFTEEN - MILE RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 64°17'139°49'

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

>1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1980 Nov 8 Flown by helicopter. River frozen over with open areas of good spawning grounds.


ESCAPEMENT RECORD FOR FIFTEEN MILE RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80		N/O		N/O		
81						
82						
83						
84						
85						

TIMING

ARRIVE						
START						
PEAK						
END						

REMARKS 1980 Flown by helicopter. No fish observed.



NAME OF STREAM FISHING BRANCH RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Porcupine River
 POSITION 66°27' 138°35'
 LENGTH 48 km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- Bear Cave Mountain area
COHO	- Above Bear Cave Mountain area
CHUM	- Heaviest around Bear Cave Mountain -- light on flats to Porcupine.
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1971 Spring fed river. Probably thermal as the upper portion of South branch is open all winter even though temperatures of -65 degrees are not uncommon. In January 1972 springs were observed coming from half way up the mountain opposite Bear Cave. The area is now under seismic exploration.
1973 Counting fence located in Bear Cave Mountain area manned throughout season counted 16,239 chum.
1974 Counting fence used throughout most of the chum run.
1975 Fence was put in after commencement of run and approx 40,000 had already passed. When camp was pulled Oct.10,75 chum still passing at 5-10,000 per day, so estimate is conservative.


continued.....

continuation

FISHING BRANCH RIVER

- 1976 Aerial surveys done twice by Alaskans - count of 13,500 chum - estimate of 15,000. Net set at Old Crow, December 2, 1976 produced 2 chum. Condition of fish was dark and full of spawn.
- 1977 Joint survey by Canadians and Americans.
- 1978 Aerial survey done Oct. 13 by Americans and Canadians. A ground survey was done by Americans Oct. 13-16 by floating river. 1 tag was recovered.
- 1980 Oct. 2 Zealand and Johnston flew this river for a count of 8907 live, 114 dead. Oct. 13 U.S. Fish and Wildlife flew system again counted 20,215 live, 104 dead. In 1979 a flight was made on Oct. 25 with a 60% dead count. Two other attempts were made to fly the Fishing Branch on the way to Old Crow, but bad weather prevented the survey.
- 1981 Survey was done by helicopter, but was not completed due to mechanical failure about 3.2km short of the confluence with the Porcupine River. Two other attempts were made at a later date, but bad weather prevented surveys.
- 1982 Helicopter survey -- conditions good. (October 12)

Predation: Grizzly bear, wolf, eagle, merganser. 1978 reported very heavy bear predation.



Sketch of Fishing Branch River
and Miner River 1971.


ESCAPEMENT RECORD FOR FISHING BRANCH

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71			25	75,000		
72			-	-		
73		25	25	15,000		
74			300	35,000		
75				350,000		
76				15,000		
77				32,000		
78				10,000		
79				75,000		
80			400	45,000		
81				10,550		
82				5,850		
83						
84						
85						

TIMING

ARRIVE		UNK	UNK	Jul - Sept		
START				Aug - Oct		
PEAK				Sept - Oct		
END				Oct - E Nov		

REMARKS



NAME OF STREAM FORTIN CREEK
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Pelly River

POSITION 62° 02' 130° 34'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1981 August 15 Flown by Alaska Dept. of Fish and Game -- no fish observed.


ESCAPEMENT RECORD FOR FORTIN CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81		N/O				
82		Not Inspected				
83						
84						
85						

TIMING

ARRIVE						
START						
PEAK						
END						

REMARKS



NAME OF STREAM FORTY MILE RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 64°26' 140°32'
 LENGTH _____ km WIDTH _____ m DRAINAGE 16,600 km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) Near the mouth max 1300 Jun. 17 1982 Min 0.024B Feb 22 1984

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1980 November 8, flown by helicopter. System totally frozen over.


ESCAPEMENT RECORD FOR FORTY MILE RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80				N/O		
81						
82						
83						
84						
85						

TIMING

ARRIVE						
START						
PEAK						
END						

REMARKS 1980 Flown by helicopter. System totally frozen over.



NAME OF STREAM GLADYS RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 120
 LOCATION OF MOUTH Flows E. into Teslin Lake through Gladys and Hall Lakes, Cassiar
 POSITION 59 132 NE. Dist. _____
 LENGTH passable 138 km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____


BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- mainly in riffles on lower portion of river--near Teslin Lake.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 Very good spawning gravel that is not being utilized.1973 Estimated figures from Native food catch at Johnsons Crossing.1978 River inspected from Teslin Lake to Hall Lake, could be some spawning grounds between Hall Lake and Gladys Lake.1980 Approx 60% of streambed is good spawning gravel.1982 Stream has good potential if runs materialize.Some scouring in first 8 km of river and some eagle predation 1973.



Sketch Map of Gladys River,
1971

△ - Chinook spawning grounds

ESCAPEMENT RECORD FOR GLADYS RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71		25				
72		25				
73		25				
74		25				
75		N/O				
76		N/I				
77		N/I				
78		14				
79		N/I				
80		N/I				
81		60				
82		25				
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		Aug				
END		Aug - Sept				

REMARKS



NAME OF STREAM GLENYON RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Pelly River

POSITION 62° 29' 134° 10'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- concentrated in upper river near Earn Lake outlet.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1969 Water very high and discoloured from abnormal rainfall.
 1970 Survey was a little late with most of the spawning completed by August 13.
 1972 Spawning completed at time of inspection.

Predation: 1968 reported bear and eagle.

ESCAPEMENT RECORD FOR GLENLYON RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		75				
67		75				
68		75				
69		N/O				
70		200				
71		75				
72		75				
73		N/I				
74		N/I				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		-				
81		N/I				
82		N/I				
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		L Jul - Aug				
PEAK		Aug				
END		L Aug - Sept				

REMARKS



Hayes River

A horizontal scale bar with vertical end caps. The number '0' is at the left end and '10' is at the right end. Below the bar, the word 'kilometres' is written. A small tick mark is present at the 5-kilometre position.

Disella

NAME OF STREAM HAYES CREEK (Hayes River)
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Teslin Lake

POSITION 62°43' 114°00'

LENGTH passable for 16 km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

>1.00

WETTED AREA _____ m²

SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SOCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

First report

GENERAL REMARKS 1972 No adult salmon observed but hand seining produced chinook fry in stream which appeared to be from the fall of 1971. 5% erosion. Stream has changed its course due to beaver dams and high water.



NAME OF STREAM HOOLE RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Pelly River

POSITION 61° 45' 131° 43'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1981 Flown by Alaska Dept. of Fish and Game.


ESCAPEMENT RECORD FOR HOOLE RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81		233				
82		N/I				
83						
84						
85						

TIMING

ARRIVE		Jul				
START		Jul				
PEAK		Aug				
END		Sept				

REMARKS



NAME OF STREAM HUSKY DOG CREEK
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Macmillan River

POSITION 63° 05' 132° 27'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- in lower 6 km
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 Area burned over this year. Heavy erosion will probably result in the next few years.


ESCAPEMENT RECORD FOR HUSKY DOG CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73		N/I				
74		N/I				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/I				
81		N/I				
82		N/I				
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		Aug 20				
END		Aug 30				

REMARKS



NAME OF STREAM JANET CREEK
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Stewart River

POSITION 63° 37' 135° 28'

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- in upper reaches of stream
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1967 Doubtful assessment. Later reports indicated good seeding.
1967 Weather conditions prevented inspection. No reports from
other sources.


ESCAPEMENT RECORD FOR JANET CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67		200				
68		75				
69		N/I				
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		Aug				
END						

REMARKS



NAME OF STREAM JENNINGS RIVER
 CONSERVATION DISTRICT 10 SUB-DISTRICT 120
 LOCATION OF MOUTH Flows NE. into Teslin Lake, Cassiar Dist.

POSITION 59 132 NE.
 LENGTH passable for 136 km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- scattered over lower 56km.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS Physical conditions: 1972 10% light gravel and fines especially in lower half of river. No change in course, but scouring has occurred in areas which could have been used for spawning.
1972 Very good gravel although may be unstable in periods of high water. This river could facilitate most of the spawning chinook in the Teslin system.
1978 Just the lower portions of Jennings River were inspected, total number of springs to the Jennings River are unknown.
1980 System reported as having very limited spawning gravel.
1981 Poor survey conditions
1982 System has limited spawning gravel.

JENNINGS RIVER see SWIFT RIVER


ESCAPEMENT RECORD FOR JENNINGS RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73		-				
74		N/I				
75		25				
76		N/I				
77		N/I				
78		3				
79		N/I				
80		N/I				
81		250				
82		40				
83		-				
84						
85						

TIMING

ARRIVE	Aug				
START	Aug				
PEAK	Aug				
END	Aug - Sept				

REMARKS



NAME OF STREAM KALZAS RIVER (Big Kalzas River)
 CONSERVATION DISTRICT 10 SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Macmillan River
 POSITION 62° 53' 135° 28'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT log jams scattered in lower sections(1972)

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SCKEYK

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

-majority 1.6km below lake but some scattered for several miles
 downstream.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1971 This survey was done too late. Evidence of dead fish along shoreline,
also evidence of redds in river.


ESCAPEMENT RECORD FOR KALZAS RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71		N/O				
72		75				
73		N/I				
74		N/I				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/I				
81		N/I				
82		N/I				
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug 10				
PEAK		Aug 20				
END		Aug 30				

REMARKS



NAME OF STREAM KLONDIKE RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 64° 03' 139° 26'
 LENGTH _____ km WIDTH _____ m DRAINAGE 7,800 km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) Above Bonanza Creek Max 640 May 29 1972 Min 4.81B Feb.1 1974

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- Throughout S.Klondike -- some seen above YCGC Dam on N. Klondike
COHO	
CHUM	- " " "
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS Physical conditions: 1972 Erosion from highway -- much rechanneling in upper portion. 1978/79 Heavy placer mining operation on tributaries of lower 16km of river is contributing to silting at the mouth of the Klondike where it flows into the Yukon. 1982 N. Klondike has changed its course in one section.

1972 Highways Department has many problems along this river. Road is on flood plain in several places and is attacked each spring. Ice jam caused extensive flooding this spring. Much rechanneling on this river. Contains stretches of excellent gravel. Historically an excellent producer. Should look at means of re-establishing the run.

1974 Evidence that the commercial gillnet fishery at mouth of Klondike River continued.....

Continuation

KLONDIKE RIVER

- 1974 may be having serious effects on the Klondike chinook run. Closure of this area recommended for 1975.
- 1975 Mouth of Klondike River was closed to all domestic and commercial fishing in efforts to build run. Due to manpower shortages unable to evaluate if closure had any beneficial effects.
- 1976 A closure was again established to all commercial and domestic fishing. No benefits were observed this year.
- 1980 Survey was conducted by helicopter. Parts of the river had frozen gravel beds and ice cover. Excellent back water sloughs were still open, but no sign of spawning salmon. Large gravel beds were still exposed. Beaver sticks were seen on gravel bed bottom. A weir at North Fork intake of Y.C.G.C. appears to prevent passage of chum into the system. Reports of chinook carcasses in the system were passed on by local people.
- 1981 Survey for chinook salmon appears to have been made too late this year and only 12 dead fish were seen. A check for chum was made but there was no evidence this year. Both surveys were flown in fixed wing.
- 1982 Tagging programme had one radio tagged chum enter the Klondike River. No survey was done for chum although a flight down the N. Klondike was done on the return from the Fishing Branch River. This was obviously too late in the season. Tagging programme found evidence of bear and eagle kills.


ESCAPEMENT RECORD FOR KLONDIKE RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25		200		
73		Not Inspected				
74		44				
75		N/O				
76		N/O		-		
77		25				
78		25		200		
79		25		100		
80		N/O				
81		N/O				
82		40		N/O		
83						
84						
85						

TIMING

ARRIVE		Jun-Aug		Aug- E Sept	
START		Jul-Aug		Sept	
PEAK		M Aug		L Sept-Oct	
END		L Aug-Sept		Oct	

REMARKS



NAME OF STREAM KLUANE LAKECONSERVATION DISTRICT 10 SUB-DISTRICT 110LOCATION OF MOUTH Flows into Kluane RiverPOSITION 61° 15' 138° 43'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m²SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION UNKNOWN

SPECIES

SECTION OF STREAM USED

SCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1973 Fair numbers of chum reported by Burwash Indians fishing off Mary's Point. One chum caught by sports fisherman at the east end of the lake -- near the Government campground.

1975 No fish were observed spawning in Christmas Bay as was the case in fall '73.


ESCAPEMENT RECORD FOR KLUANE LAKE

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73				UNK		
74				N/O		
75				N/O		
76				N/O		
77				N/I		
78				N/I		
79						
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE				Oct		
START				Oct 15		
PEAK				-		
END				Nov 20		

REMARKS



NAME OF STREAM KLUANE RIVER

CONSERVATION DISTRICT 10 SUB-DISTRICT 110

LOCATION OF MOUTH Flows into Donjek River

POSITION 61° 53' 139° 43'

LENGTH Passable for 160km WIDTH _____ m DRAINAGE 4950 km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) Outlet of Kluane Lake max 382 Aug.14 1971 min 0.538B Feb.27 1971

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	- in mainstream, Kluane Lake and side-channels
"	- mainly in upwelling areas near Mile 1118 of Alaska Hwy.(1982)
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1966 Chums seen spawning in Yukon River near mouth of White River
where this run enters the White-Donjek-Kluane system
1968 Some side channels blocked by beaver dams.
1970/71 Only one or two salmon caught by natives in the area. No others seen.
Usually a fair run in this river. Possibility of an obstruction
somewhere. Ice hinders observation.
1971 On first inspection stream conditions were poor for aerial count.
The second count was done on foot and was a better count. But due to
the vast area to cover only the most desirable spots were checked.

continued.....

continuation

KLUANE RIVER

1975 A poor return in light of the 1971 count of 3000 spawning.

1976 Extremely poor return this year.

1977 American count:	Quill Creek Area	43 live	6 carcasses
	Klondike Lodge Area	95 live	10 carcasses
	Swede Johnson Area	703 live	33 carcasses
		<u>841</u>	<u>49</u>

1978 Americans inspected on Oct. 17 and no fish were observed in the usual sloughs. Canadians inspected again in Nov. 22 and all but one slough was frozen over and no fish were observed.

1979 Chinook were observed spawning near the outlet of Tincup Lake. The spawning this season was the best the locals have seen in recent years. Salmon spawning was reported along the southerly shore of Kluane Lake.

Biological conditions:

1979 Some egg digging in some back channels.

1981/82 Few loose eggs in pool areas.

Light to moderate predation by Wolf and Eagle.


ESCAPEMENT RECORD FOR KLUANE RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66				2,000		
67				3,500		
68				3,500		
69				7,500		
70				N/O		
71				3,500		
72				200		
73				3,500		
74				400		
75				200		
76				20		
77				300		
78				N/O		
79		UNK		100,000		
80				5,000		
81				26,000		
82				10,000		
83						
84						
85						

TIMING

ARRIVE				Sept		
START				L Sept-Oct		
PEAK				Oct - L Oct		
END				Oct - E Nov		

REMARKS



NAME OF STREAM KOIDERN RIVER

CONSERVATION DISTRICT 10 STATISTICAL AREA 110

LOCATION OF MOUTH Flows into White River

POSITION 62° 03' 140° 27'

LENGTH passable for 60 km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1981 Chum were discovered in this system by Foothills Pipeline
personnel during studies of 1979. The area has been studied since
then to determine overwintering and rearing areas. However, this
season, no spawning chums were observed in this system.
1982 Normal fluctuations in water levels.

ESCAPEMENT RECORD FOR KOIDERN RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81					N/O	
82					N/O	
83						
84						
85						

TIMING

ARRIVE						
START						
PEAK						
END						

REMARKS



NAME OF STREAM LAPPIE RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into Pelly River

POSITION 62° 02' 132° 36'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1971 No carcasses visible -- good stream.

1979 Game Branch, Ross River reports one local angler who caught a chinook in the Lappie River below the canyon.

1980 Yukon Fish and Game Branch indicated that the Native Indians have caught Chinook at the confluence of the Lappie and Pelly River. Reports of sports fishermen catching chinook in this area too.

1981 4 Chinook were seen holding in pools awaiting to migrate through canyon
Jul. 31.

1982 Information from COSEP student reveals that Ross River natives fish for salmon in the lower portions of the Lappie River.

ESCAPEMENT RECORD FOR LAPIE RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71		None Observed				
72		-				
73		-				
74		N/I				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/I				
81		N/I				
82		N/I				
83						
84						
85						

TIMING


ARRIVE

START

PEAK

END

REMARKS



NAME OF STREAM LEWIS LAKECONSERVATION DISTRICT 10STATISTICAL AREA SUB - DISTRICT 110LOCATION OF MOUTH Flows into Ross RiverPOSITION 62° 37' 131° 05'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m²SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT Obstruction well above spawning grounds.

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

- first .80km below lake

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1982 Flown by Alaska Dept. of Fish and Game.

ESCAPEMENT RECORD FOR LEWIS LAKE

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		75				
73		N/I				
74		N/I				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/I				
81		N/I				
82		39				
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		Jul - Aug				
PEAK		Aug				
END		L Aug-E Sept				

REMARKS



NAME OF STREAM LITTLE KALZAS RIVER (MacMillan River)
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into Big Kalzas River

POSITION 62° 53' 135° 28'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT passable forest debris 16km from mouth

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- mainly in upper 1.6km to lake --- above and below Big Slough
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	


POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1966 First report. Guardian reports good runs in previous years.

1967 Reports indicate good escapement. Turbid water from flashflooding precluded assessment.

1970 Due to weather inspections were incomplete.

1971 Predation by Grizzly Bears.



Sketch Map of Little Kalzas River, 1966
Not to Scale


ESCAPEMENT RECORD FOR LITTLE KALZAS RIVER (Macmillan River)


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		400				
67		400				
68		200				
69		N/I				
70		400				
71		25				
72		25				
73		N/I				
74		N/I				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/I				
81		N/I				
82		N/I				
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		Aug				
END		Aug - Sept				

REMARKS





NAME OF STREAM LITTLE SALMON RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 62° 03' 135° 40'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25
 0.25 - 0.50
 0.50 - 0.75
 0.75 - 1.00
 > 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- heaviest concentration in upper .50km to lake outlet and
"	scattered throughout river in pockets.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1966 First survey of this stream. - water very low and clear.
1967 New road from Carmacks to Ross River now under construction will open
this entire area to the public in 1968.
1968 Road open. Possibility of Natives fishing here in future.
1969 Water high and discolored throughout the summer. Salmon were scattered
along several miles of the stream but did not appear to be spawning.
1977 This is one of the index streams flown by the Americans.
1978 Actual counts: Americans August 8 - 16 330 chinook 70% Eff.
Canadians August 21 200 chinook 75% of river


continued.....

continuation

LITTLE SALMON RIVER

- 1980 This system was flown by U.S. Fish and Game Branch under poor conditions. 276 live springs, 10 dead.
- 1979 It is estimated that only 25% of the spawners were on redds at the time of the inspection. Many large redds had no fish on them while several large redds had only a few fish on them. This inspection was combined Canadian and American. Actual count: 280 chinook - 5 carcasses.
- 1981 Flown by Alaska Fish and Game.
- 1982 This survey was done by Yukon River Basin study and was a canoe survey.

Predation by eagle and bear.



Sketch Map of Little Salmon
River, 1966.

△ - Chinook spawning grounds.


ESCAPEMENT RECORD FOR LITTLE SALMON RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		200				
67		200				
68		3,500				
69		750				
70		750				
71		400				
72		400				
73		75				
74		N/O				
75		N/I				
76		N/I				
77		258				
78		500				
79		1,100				
80		400				
81		750				
82		350				
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		Jul - Aug				
PEAK		Aug				
END		Aug - Sept				

REMARKS



NAME OF STREAM McNEIL RIVERCONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110LOCATION OF MOUTH Flows into Nisutlin RiverPOSITION 61° 10' 132° 17'LENGTH passable for 12 km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 Poor spawning grounds -- large boulders most of river with very coarse rock.


ESCAPEMENT RECORD FOR McNEIL RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		2				
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		Aug				
END		Aug				

REMARKS



NAME OF STREAM McQUESTEN RIVERCONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110LOCATION OF MOUTH Flows into Stewart RiverPOSITION 63° 33' 137° 27'LENGTH km WIDTH m DRAINAGE 2870 km²COMPOSITION: BEDROCK BOULDER COARSE FINE SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25 0.25 - 0.50 0.50 - 0.75 0.75 - 1.00 > 1.00 WETTED AREA m² SPAWNING AREA m²DISCHARGE (m³/s) Near the mouth Max 237 30 May 1982 Min 4.98B 15 Feb 1981TEMPERATURE (°C) BARRIERS OR POINTS OF DIFFICULT ASCENT

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SOCKEYE CHINOOK - 6 found in N. McQuesten River--the rest distributed between
N. McQuesten and Stewart River.CHUM PINK (ODD YEAR) PINK (EVEN YEAR) STEELHEAD POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS Dan Drummond, conservation officer in Dawson reported 24 chinook
in the McQuesten River between the bridge and the confluence with the
Stewart River.

1980 Flown by helicopter. Fine stretches of open water with good spawning gravel.
A foot patrol of lower reaches in September revealed no chinook. However,
natives from Mayo were fishing at the mouth and caught approx. 800 chinook--
information from Band Chief and Yukon Game Branch.

1981 Only 2 chinook were spotted from the crossing near Elsa to the Stewart
River. Possible that survey was late, or that Native nets at the mouth
of river were catching all returning salmon.

continued.....

continuation

McQUESTEN RIVER

- 1981 Another possibility is that Elsa mines had a spill that may have killed many of the fry in the brood year. Local inhabitants have reported that the grayling fishing has gone down compared with previous years.
- 1982 Survey flown during early evening in fixed wing. Light good.

Eagle predation.


ESCAPEMENT RECORD FOR McQUESTEN RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74		40				
75		N/I				
76		24				
77		N/I				
78		N/I				
79		N/I				
80		N/O				
81		2				
82		38				
83						
84						
85						

TIMING

ARRIVE		Jul				
START		Jul				
PEAK		Aug				
END		Aug - Sept				

REMARKS



NAME OF STREAM MAYO RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA 110
 LOCATION OF MOUTH Flows S. into Stewart at Mayo

POSITION 63° 35' 135° 54'

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) Near Mayo Max 121 Jun 6 1949 - Min 4.98B Feb 15 1981

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable to NCPC Dam.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	<u>-from Dam to junction of Mayo and Stewart Rivers - mostly in area</u>
COHO	<u>just below dam.</u>
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 NCPC Dam is shut down 1 or 2 days each year for inspection.

This results in the complete shut-off of water. Many fish were found lying on the stream bed at a time when chinook fry are abundant in the stream.

1975 It appears the Mayo River has reached its inevitable end and rehabilitation of some sort should be explored.

1980 NCPC had problems with their dam in September and the water levels fluctuated with the opening and closing of the dam. Silt levels were high in the system which made it impossible to do a stream count for chinook.


Helicopter survey in November revealed do chum in the system.

continued.....

continuation

MAYO RIVER

- 1981 Spawning occurs from the dam down stream to the Stewart River. Major area is just below the dam. The report this year indicates that NCPC shut off the dam flow during spawning season leaving salmon stranded.



Sketch Map of Mayo River, 1971

4cm = .80km

△ - Chinook spawning grounds.


ESCAPEMENT RECORD FOR MAYO RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71		400				
72		75		25		
73		N/I				
74		2				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/O		N/O		
81		11				
82		N/I				
83						
84						
85						

TIMING

ARRIVE		Jul - E Aug		E Aug		
START		Aug		E Aug		
PEAK		Aug		M Aug		
END		L Aug - Sept		L Aug		

REMARKS



NAME OF STREAM MICHIE CREEK
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into McClintock River

POSITION 60° 40' 134° 28'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT 1967 passable minor log jams.
1966 passable beaver dams

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	<u>- in 16km downstream from Lake -- mostly at lake outlet.</u>
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS Physical conditions: 1969 2 small landslides on McClintock River
added to turbidity in that stream.
1972 Heavy erosion and bank cutting in McClintock R. Some rechanneling
evident. Highest water levels ever recorded.
1968 Appeared to be heavier concentration of salmon on riffles than in previous
year although escapement through fishway lighter.
1970 Clay banks unstable and sloughing into river.
1971 Did not positively identify 966 chinook that were reported through Whitehorse
Dam. This could be an indication of low aerial counts or that possibly all the
fish registered through dam do not spawn in Mitchie Creek.

continued.....

MICHIE CREEK

- 1972 Count determined by number passing through Whitehorse fish ladder.
This is only known spawning ground above ladder.
- 1973 226 chinook passed through Whitehorse ladder.
- 1974 273 chinook passed through the Whitehorse fishway presumably for
this stream.
- 1975 Stream was looked at quite thoroughly with helicopter and portions
were walked to compare aerial counts.
- 1976 Assessed by helicopter as well as on foot.
- 1977 276 adult chinook utilized the Whitehorse fishway in 1977.
- 1978 Whitehorse fishway final count - 725 adult chinook.
- 1979 " " " " -1184 adult chinook.
- 1980 Flown by Larry Kratt of Environmental Management Associates.

Predators: bear and eagle.


ESCAPEMENT RECORD FOR MICHIE CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		75				
67		75				
68		200				
69		N/O				
70		750				
71		200				
72		400				
73		N/I				
74		N/I				
75		39				
76		22				
77		36				
78		N/I				
79		N/I				
80		1,700				
81						
82						
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		L Aug-Sept				
END		L Aug-Sept				

REMARKS _____



NAME OF STREAM MINER RIVERCONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110LOCATION OF MOUTH Flows into Purcupine RiverPOSITION 66° 30' 138° 25'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m²SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

- lower 64km of river.

- in area immediately below confluence of Fishing Branch River.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1971 Spawning may be limited in this river as bulk of observed
salmon may have been heading to Fishing Branch.1981 River frozen over.

MINER RIVER see FISHING BRANCH


ESCAPEMENT RECORD FOR MINER RIVER


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71				750		
72						
73						
74		75				
75		N/I				
76		N/I				
77		N/I				
78		N/I				
79		N/I				
80		N/I				
81		N/O				
82		N/I				
83						
84						
85						

TIMING

ARRIVE	UNK	-		
START		Sept		
PEAK		Oct 10		
END		Oct 20		

REMARKS





NAME OF STREAM MORLEY RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB- DISTRICT 110
 LOCATION OF MOUTH Flows into Teslin Lake

POSITION 60° 06' 132° 29'
 LENGTH 90 km WIDTH m DRAINAGE km²
 COMPOSITION: BEDROCK BOULDER COARSE FINE
 SILT & SAND UNCLASSIFIED

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA m² SPAWNING AREA m²

DISCHARGE (m³/s)

TEMPERATURE (°C)

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable for 90 km but could be problems
at certain water levels — falls and rapids at 24km.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- Morris L. to Teslin L. heaviest between Morley L. and Teslin L.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM

GENERAL REMARKS A good spawning stream.

1966 A few chinook still active near mouth on Sept 5, Anglers reported taking a few
in August.

1969 High murkey water precluded sightings on spawning grounds on two inspection
flights.

1977 Morley River was inspected from Slim Lake to Morley Bay, the area chinook
were found in was from the Alaska Highway to Strawberry Creek. This area
should become an index area.

1981 Fair survey conditions. Estimated that 100 chinook were taken at the Alaska
Highway Bridge by sports fishermen and poaching.


1982 Only minimal sport fishing pressure noted at the Morley River Bridge this year.

Predators: eagles and bears.



Sketch Map of Morley River,
1981
Scale 1:250000

△ - Chinook spawning
grounds




ESCAPEMENT RECORD FOR MORLEY RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		400				
67		400				
68		200				
69		25				
70		200				
71		25				
72		25				
73		-				
74		N.I.				
75		30				
76		N.I.				
77		32				
78		70				
79		N.I.				
80		300				
81		350				
82		260				
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		Aug				
PEAK		M Aug				
END		Aug-E Sept				

REMARKS



NAME OF STREAM NISUTLIN RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB -DISTRICT 110
 LOCATION OF MOUTH Flows into Teslin Lake

POSITION 60 14' 132 34'
 LENGTH _____ km WIDTH _____ m DRAINAGE 8030 km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) Above Wolf River Max 541 May 29 1981 - Min 9.75B Apl 15 1982

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____



SPAWNING DISTRIBUTION

SPECIES SECTION OF STREAM USED

SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS _____



Sketch Map of Nisutlin River
and Wolf River, 1981

Not to scale


ESCAPEMENT RECORD FOR NISUTLIN RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		1,200				
67		750				
68		1,500				
69		400				
70		1,500				
71		1,500				
72		184				
73		75				
74		150				
75		210				
76		200				
77		111				
78		540				
79		N.I.				
80		2,500				
81		2,700				
82		1,948				
83						
84						
85						

TIMING

ARRIVE		L Jul-E Aug				
START		Jul - Aug				
PEAK		M Aug				
END		Aug - Sept				

REMARKS 1972 Aerial survey
 1973 These figs. from Alaskans out of Anchorage.



NAME OF STREAM NORDENSKIOLD RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB- DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 62° 06' 136° 18'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION OVER 125km AREA

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS New file 1978

1980 Flown by fixed wing. River totally frozen over. Reports state that at one time Native Indians used to have a subsistence fishery for chum and chinook.
1981 No fish were seen on either survey by either method due to meanders and also bad light conditions. Several reports that chinook and chum migrate up this system.

ESCAPEMENT RECORD FOR NORDENSKIOLD RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78		25				
79		N.I.				
80		N.O.				
81		N.O.				
82		N.I.				
83						
84						
85						

TIMING

ARRIVE		L Jul				
START		Aug				
PEAK		Aug				
END		E Sept				

REMARKS



NAME OF STREAM NORTHERN LAKECONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110LOCATION OF MOUTH Flows into North Big Salmon RiverPOSITION 61° 48' 133° 44'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

>1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SOCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

- concentrated at lake outlets and in upper 1.6km

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1969 Only 5 salmon sighted -- no carcasses. 5 total reported by
game guides.


ESCAPEMENT RECORD FOR NORTHERN LAKE

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		50				
67		75				
68		75				
69		25				
70		25				
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		Aug				
PEAK		Aug				
END		L Aug				

REMARKS



NAME OF STREAM (Ollie River)CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110LOCATION OF MOUTH Flows into Hess RiverPOSITION 63 15' 132 35'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SCKEYK

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

- at outlet of both lakes.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS _____


ESCAPEMENT RECORD FOR (Ollie River, Ollie Lakes)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73		N.I.				
74		N.I.				
75		N.I.				
76		N.I.				
77		N.I.				
78		N.I.				
79		N.I.				
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE		Aug				
START		M Aug				
PEAK		L Aug				
END		M Sept				

REMARKS



NAME OF STREAM PELLEY LAKES
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Pelly River

POSITION 62° 04' 130° 17'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

Sockeye

Chinook

Coho

Chum

Pink (odd year)

Pink (even year)

Steelhead

- concentrated at outlet of Lower Lake, also between 1st and 2nd lakes.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

1970

GENERAL REMARKS Difficult to access because of water discoloration.1971 Hard to assess accurately due to lateness of count.1981 Flown by Alaska Dept. of Fish and Game.


ESCAPEMENT RECORD FOR PELLY LAKES

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70		400				
71		200				
72		25				
73		N.I.				
74		N.I.				
75		N.I.				
76		N.I.				
77		N.I.				
78		N.I.				
79		N.I.				
80		-				
81		67				
82		N.I.				
83						
84						
85						

TIMING

ARRIVE		Jul - E Aug				
START		M Aug				
PEAK		L Aug				
END		L Aug				

REMARKS



NAME OF STREAM PELLEY RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 62° 47' 137° 20'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____


BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- in stream between lakes and lower lake outlet.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1968 Water high and discolored at time of inspection.
1980 Conversation with the Pelly Band chief indicates that 1,200 chinook
were caught this year by the band.
1981 A small section was flown while on the way to other river systems. Approx
12 pairs of spawning chinook were seen. Deep water areas were not clear
due to silt loads.




ESCAPEMENT RECORD FOR PELLY RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68		750				
69		N.O.				
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80		N.I.				
81		UNK				
82		N.I.				
83						
84						
85						

TIMING

ARRIVE		Jun - Jul				
START		Jul				
PEAK		Aug				
END		Sept				

REMARKS



NAME OF STREAM PLEASANT CREEK
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Hess River

POSITION 63° 31' 133° 55'

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m²

SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES SECTION OF STREAM USED

SOCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

- just below lake on excellent gravel.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS _____


ESCAPEMENT RECORD FOR PLEASANT CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73		N.I.				
74		N.I.				
75		N.I.				
76		N.I.				
77		N.I.				
78		N.I.				
79		N.I.				
80		N.I.				
81		N.I.				
82		N.I.				
83						
84						
85						

TIMING

ARRIVE						
START						
PEAK		Aug 10				
END						

REMARKS



NAME OF STREAM RED RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Wolf River

POSITION 60° 44' 132° 07'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- throughout river
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 Reasonably clear stream -- good survey.
Eagle predation.

ESCAPEMENT RECORD FOR RED RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE

START

PEAK


END

Aug

Aug

Aug

REMARKS



NAME OF STREAM RIDDELL RIVER (South MacMillan River)
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into South MacMillan River

POSITION 62° 51' 132° 25'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- scattered to headwaters
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1982 Flown by Alaska Dept. of Fish and Game.


ESCAPEMENT RECORD FOR RIDDELL RIVER (South MacMillan River)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73		N.I.				
74		N.I.				
75		N.I.				
76		N.I.				
77		N.I.				
78		N.I.				
79		N.I.				
80		N.I.				
81		N.I.				
82		3				
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		Jul - Aug				
PEAK		Aug				
END		L Aug-Sept				

REMARKS



NAME OF STREAM ROSS RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Pelly River

POSITION 61° 59' 132° 25'
 LENGTH _____ km WIDTH _____ m DRAINAGE 7,250 km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25
 0.25 - 0.50
 0.50 - 0.75
 0.75 - 1.00
 > 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) At Ross River Max 742 Jun 1 1972 - Min 2.83B Mar 7 1974

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable to third lake

SPAWNING DISTRIBUTION

SPECIES SECTION OF STREAM USED

SOCKEYE	
CHINOOK	Heaviest concentration at outlet of Lewis Lake. Spawning between
COHO CHINOOK	Lewis and Field Lakes and Lewis Lake to Big Timber Creek.
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

1971

GENERAL REMARKS Good counting conditions on this stream. A good gradient for chinook spawning in many areas.
1981 Flown by Alaska Dept. of Fish and Game.

ESCAPEMENT RECORD FOR ROSS RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71		1,500				
72		-				
73		-				
74		-				
75		-				
76		-				
77		-				
78		-				
79		-				
80		-				
81		1,102				
82		116				
83						
84						
85						


TIMING

ARRIVE		Jul - Aug				
START		Jul - Aug				
PEAK		Aug				
END		Sept				

REMARKS 1981 Alaska Fish and Game



703 702 15' 701



NAME OF STREAM RUSSEL CREEK (North MacMillan River)
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB- DISTRICT 110
 LOCATION OF MOUTH Flows into MacMillan River
 POSITION 63° 02' 133° 24'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT impassable canyon gorge at 5km.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	<u>below canyon - good gravel</u>
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1981 This area has been subjected to placer mining in the early 1900's and is now being subjected to mining again. If this stock is to survive strict enforcement of controls on the mining operation should be followed.

1981 Spawning occurs below canyon to MacMillan River. Possible for salmon to migrate through canyon although not documented. Placer mining in upper reaches above canyon.


ESCAPEMENT RECORD FOR RUSSEL CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73		N.I.				
74		N.I.				
75		N.I.				
76		N.I.				
77		N.I.				
78		N.I.				
79		N.I.				
80		N.I.				
81		11				
82		N.I.				
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		L Aug				
END		L Aug				

REMARKS



NAME OF STREAM SHELDON LAKE
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Ross River

POSITION 62° 42' 131° 03'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION UNKNOWN

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 This river has about 25km of what appears to be excellent gravel which is not being used by salmon. Hudson Bay Mining are presently developing a mine in this area.


ESCAPEMENT RECORD FOR SHELDON LAKE

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		25				
73		N.I.				
74		N.I.				
75		N.I.				
76		N.I.				
77		N.I.				
78		N.I.				
79		N.I.				
80		N.I.				
81						
82						
83						
84						
85						

TIMING

ARRIVE		Aug				
START		E Aug				
PEAK		M Aug				
END		L Aug				

REMARKS



NAME OF STREAM SMART RIVER (Swift River)
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into Teslin Lake

POSITION 60° 00' 131° 45'
 LENGTH passable for 40 km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SOCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD


- between Cabin Lake and Dorsey Lake.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1981 Fair to poor survey.

1982 Only a marginal salmon producing stream.

1982 Flown by Foothills Pipeline.




ESCAPEMENT RECORD FOR SMART RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80		100				
81		50				
82		N.O.				
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		Aug				
END		Aug				

REMARKS



NAME OF STREAM STEWART RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 63° 19' 139° 24'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1971 Silting from Hess River. Fraser Falls may obstruct passage when water level is low.

ESCAPEMENT RECORD FOR STEWART RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71		N.O.				
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING


ARRIVE

START

PEAK

END

REMARKS



NAME OF STREAM SWEDE JOHNSON CREEK
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into Kluane River

POSITION 61° 36' 139° 24'

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

>1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT Series of beaver dams near highway .80km
from mouth.

SPAWNING DISTRIBUTION

SPECIES SECTION OF STREAM USED

SOCKEYE

CHINOOK

COHO

CHUM

- scattered in lower reaches

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1966 Creek and sidestreams almost dry. Salmon unable to enter at some
points.

1967 Very few sidestream spawners in 1967. All in Kluane River.

1970 No chum salmon were seen in the area by residents. Ice made
observation impossible when stream was checked. Usually a
fair run in this stream. Beaver dams may have blocked the run
with an early freeze.


ESCAPEMENT RECORD FOR SWEDE JOHNSON CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66				800		
67				25		
68				25		
69				25		
70				N.O.		
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE						
START				Sept - Oct		
PEAK				Oct		
END				Oct - Nov		

REMARKS



NAME OF STREAM SWIFT RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110
 LOCATION OF MOUTH Flows into Teslin Lake

POSITION 60° 49' 133° 51'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT Falls and rapids 8 and 32kms from mouth
passable depending on water levels--should be monitored.

1972 Bedrock outcrop 16km from mouth.

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SOCKEYE

CHINOOK

concentrated in upper 5km below Swan Lake outlet and to junction
of Smart River

C. IM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD


POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1967 None sighted during aircraft surveys but reports indicate
fair number in river at later date.

1972 .02% natural erosion.

1977 Swan Lake to junction of Smart River should become in index area and
looked at every year.

Bear predation.



Sketch Map of Gladys River
Swift River
Jennings River,
1981.

Δ - Chinook spawning grounds


ESCAPEMENT RECORD FOR SWIFT RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		200				
67		200				
68		75				
69		N.O.				
70		-				
71		200				
72		25				
73		-				
74		N.I.				
75		N.I.				
76		N.I.				
77		40				
78		15				
79		N.I.				
80		600				
81		350				
82		57				
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		Aug				
PEAK		Aug				
END		Aug - Sept				

REMARKS





NAME OF STREAM TAKHINI RIVERCONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110LOCATION OF MOUTH Flows into Yukon RiverPOSITION 60° 51' 135° 11'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m²SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT Passable for 80km rapids with narrow
channel 48km from mouth.

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SCKEYE

CHINOOK

- in upper 8km below outlet from Arkell(Kusawa)Lake and from
Kusawa Lake down to Haines Junction Hwy. Bridge.

CHUM


PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS Physical conditions:1972 5% erosion and silting -- no change in course1967 Indians not fishing this run for the first time in many years.1970 Sport fishermen caught a chinook in the lake by trolling and several
more were seen near the head of the lake. May be some spawning in this area
that we are not aware of.1972 This stream very hard to survey due to glacial runoff and milky blue colour
of water.1974 This was a late count done by air. 1975 Very good count.1976 Stream very glacial and milky blue -- hard to see fish.1980 Good visibility. 1981 Flown by Sandy Johnson.Predation by bears and birds.




ESCAPEMENT RECORD FOR TAKHINI RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		350				
67		750				
68		750				
69		75				
70		200				
71		-				
72		17				
73		-				
74		80				
75		165				
76		6				
77		88				
78		115				
79		N.I.				
80		950				
81		782				
82		38				
83						
84						
85						

TIMING

ARRIVE		Jul				
START		Jul - Aug				
PEAK		Aug				
END		Aug - Sept				

REMARKS



NAME OF STREAM TATCHUN RIVERCONSERVATION DISTRICT 10 STATISTICAL AREA SUB - DISTRICT 110LOCATION OF MOUTH Flows into Yukon RiverPOSITION 62°17' 136°20'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

>1.00

WETTED AREA _____ m²SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT 1971 Log jam midway between road and lake1974 Impassable beaver dam 365 meters from mouth, recommend removal.1975 Recommend removal of beaver dam. Obstruction report submitted 12.12.751976 Beaver dams monitored.1977 A severe beaver dam problem exists on Tatchun Creek. There are seven dams between mouth and the lake. Some of these were torn out by hand several times in 1977. A powder program is planned for spring of 1978 to rectify this problem.

SPAWNING DISTRIBUTION

SPECIES

SECTION OF STREAM USED

SOCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

- from mouth to Lake, concentrated between highway and Tatchun L.

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1966 Only 1 chinook salmon taken by Indians at the mouth of this stream. Low water conditions altered flow in Yukon River and back eddies disappeared.1967 Native Indians took 250 chinook off mouth of this stream.1968 100 chinook taken by Native Indians at mouth of this stream.1970 Campground is located on this creek also a net fishery is carried on at the mouth by Indians.

continued.....

continuation

TATCHUN RIVER

- 1978 3 beaver dams were blown out of Tatchun Creek — some activity by beavers during summer that had to be removed by hand. Did not present a problem to migrating adults. Chinook were utilizing new areas in 1978.
- 1979 Some beaver problems — inspection was done very late and most spawning had been completed. An intense sport fishery has developed in the Yukon River below Tatchun Creek in 1978 and 1979. This fishery warrants some attention and enforcement in 1980.
- 1980 Heavy sport fishery at the mouth for chinook will have to be patrolled in 1981. Sport fishing for grayling is very popular due to location of Yukon camp ground near highway. 142 live chinook, 71 dead, 9 jacks. Foot patrol.
- 1981 The wash out of two old beaver dams in the upper reaches of the creek has created new spawning gravels which the fish were using this year. At least 50 fish were using one pool. Creek was walked on the 15th one week after part of the lower dam had been removed. It was necessary to remove the dam again on the 24th at which time another 65 fish migrated up stream. Sport fishing has become very heavy at the mouth of the Yukon River and an estimated 520 were taken in this area. One report is that at least 40 fish caught in one day.
- 1982 Beaver dams in lower portions washed out in spring allowing fish passage.

Predators: bear and eagle.


ESCAPEMENT RECORD FOR TATCHUN RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		50				
67		25				
68		25				
69		6				
70		75				
71		200				
72		-				
73		200				
74		192				
75		175				
76		47				
77		150				
78		200				
79		150				
80		400				
81		205				
82		73				
83						
84						
85						

TIMING

ARRIVE		L Jul-E Aug			
START		Jul - Aug			
PEAK		L Aug-Sept			
END		Sept			

REMARKS



NAME OF STREAM TESLIN RIVER
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Yukon River

POSITION 61° 34' 134° 54'
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- upper half of stream for approx 160km (1968)
COHO	- concentrated about 2km below bridge -- random to Roaring Bull
CHUM	(1972)
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1969 Stream high and very turbid during entire season -- due to abnormal precipitation.
1978 Just the top end of the river at Johnsons Crossing was surveyed.
1979 Water was high at time of inspection. Visibility was poor, also an intense sport fishery was developed here in 1979. Warrants some attention in 1980.
1981 Upstream of Teslin Bridge is surveyed N.B.C and Yukon Sub-District. This year survey flown by fixed wing. High winds and poor light conditions made observations difficult.
1982 Although not inspected there was a tagging program on chinook and chum near eagle Alaska and radio tagging was tracked to the Teslin.


ESCAPEMENT RECORD FOR TESLIN RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		3,500		UNK		
67		3,500		UNK		
68		7,500		UNK		
69		1,500		UNK		
70		7,500				
71		-				
72		1,500		75		
73		N.I.				
74		N.I.				
75		N.I.				
76		N.I.				
77		N.I.				
78		500				
79		500				
80		-				
81		45				
82		N.I.				
83						
84						
85						

TIMING

ARRIVE		E Aug				
START		E Aug				
PEAK		Aug - Sept				
END		Sept - Oct				

REMARKS



NAME OF STREAM (Upper Teslin River)
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Teslin Lake

POSITION _____
 LENGTH passable for 30 km WIDTH _____ m DRAINAGE _____ km²
 COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT Small falls and chute at 11 km from mouth.
passable at certain water levels. The obstruction
could create a serious problem to passage of fish
at low water levels. Possibly a minimum amount of
blasting of bedrock would improve conditions.

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- upper portion of river
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 15% light gravel and fines. No change in course but stream bed
has changed recently due to heavy run-offs.

Some eagle predation.


ESCAPEMENT RECORD FOR (Upper Teslin River)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72		10				
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84						
85						

TIMING

ARRIVE		Aug				
START		Aug				
PEAK		Aug				
END		Sept				

REMARKS



NAME OF STREAM WOLF RIVER (Nisutlin River)
 CONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110
 LOCATION OF MOUTH Flows into Teslin Lake

POSITION 60° 17' 132° 33'
 LENGTH passable for 128 km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____
 SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25	
0.25 - 0.50	
0.50 - 0.75	
0.75 - 1.00	
> 1.00	

WETTED AREA _____ m² SPAWNING AREA _____ m²

DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION

SPECIES	SECTION OF STREAM USED
SOCKEYE	
CHINOOK	- concentrated in upper .50km to Wolf Lake and from Wolf Lake to below Fish Lake.
COHO	
CHUM	
PINK (ODD YEAR)	
PINK (EVEN YEAR)	
STEELHEAD	

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1972 Lower half of river composed of fine sands and silts not suitable for spawning.

1975 Actual age composition of Wolf River run as yet unknown.

1977 Area inspected was from Nisutlin River to Fish Lake (64km) and only 12 chinook were observed. The second area inspected was from Wolf Lake to approx 12km downstream and 25 chinook were observed. The second area should become an index area.

1978 Area inspected was from Wolf Lake to approx 13km down river. The remainder of the river was not inspected.

1981 Poor to fair weather conditions during survey.

1982 Flown by A.F.G.

Predation by bears and birds

ESCAPEMENT RECORD FOR WOLF RIVER


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66		75				
67		75				
68		75				
69		25				
70		200				
71		750				
72		45				
73		-				
74		N.I.				
75		40				
76		N.I.				
77		37				
78		44				
79		N.I.				
80		800		200		
81		502				
82		350				
83						
84						
85						

TIMING

ARRIVE		Jul - Aug				
START		Jul - Aug				
PEAK		Aug				
END		Aug - Sept				

REMARKS

kilometres





NAME OF STREAM YUKON RIVERCONSERVATION DISTRICT 10 STATISTICAL AREA SUB-DISTRICT 110LOCATION OF MOUTH Flows into Bering Sea, mouth 62 164 NW.POSITION 64° 41' 141° 00'LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²

COMPOSITION: BEDROCK _____ BOULDER _____ COARSE _____ FINE _____

SILT & SAND _____ UNCLASSIFIED _____

PERCENT GRADIENT

0.00 - 0.25

0.25 - 0.50

0.50 - 0.75

0.75 - 1.00

> 1.00

WETTED AREA _____ m²SPAWNING AREA _____ m²DISCHARGE (m³/s) _____

TEMPERATURE (°C) _____

BARRIERS OR POINTS OF DIFFICULT ASCENT _____

SPAWNING DISTRIBUTION SEE GENERAL COMMENTS.

SPECIES

SECTION OF STREAM USED

SCKEYE

CHINOOK

COHO

CHUM

PINK (ODD YEAR)

PINK (EVEN YEAR)

STEELHEAD

- in deep water of Ingersol Islands, Yukon Crossing, outlet Lake Laberge
- mainly in sloughs (73) sloughs dry(1982) spawning in mainstem

POTENTIAL OF INACCESSIBLE PORTION OF STREAM _____

GENERAL REMARKS 1973 Spawning chinook located in deep water off Ingersol Islands.Location could only be determined by drifting nets over the spawning grounds.Water is fast and deep. Real possibility that mainstem spawning is extensive.Chum spawning in mainstem appears to be restricted to sloughs.1974 Population estimates for chinook and chum based on the Yukon River tagging project conducted this season. Spawning grounds located at Yukon Crossing, Ingersol Island and the outlet of Lake Laberge.1975 This year the tag recovery program was not continued so total population estimates are not available. The recorded 600 chinook and 7,000 chum were located between Minto and Fr. Selkirk. Total count over Whitehorse Fishway 309.

continued.....

continuation

YUKON RIVER

- 1976 The figure of 5-10,000 is an estimate for the Yukon System most of which is thought to be mainstem
- 1979 This escapement of 60,000 includes the total Yukon system.
- 1980 No actual survey of the Yukon River was done this year except on November 11. At this time the water was too green to see any fish and no carcasses were seen.
- 1981 This years escapement figures are based on index stream returns and commercial and domestic catch date. Main stem spawning does occur, but due to dirty water it is impossible to determine exactly where and how many. Hoochekoo Bluff has a slough area near it on the east side of the Yukon River, and 250 chum were spotted on 31st September. One week earlier chum salmon were seen jumping every three minutes at Minto Landing for a half hour period.
- 1982 Boat survey of Ingerson Island area by Yukon River Basin Study.

ESCAPEMENT RECORD FOR YUKON RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73		UNK		UNK		
74		25,000		23,000		
75		600		7,000		
76		7,500				
77		45,000				
78		90,000				
79		60,000				
80		70,000	5,000	10,000		
81		50,000		70,000		
82		UNK		1,020		
83						
84						
85						

TIMING

ARRIVE		Jun - Jul	Oct	Aug-Sept		
START		Jun - Aug	Oct	Aug-Sept		
PEAK		E Jul-Aug	Oct	Sept-Oct		
END		Sept - Oct	Nov	Oct - Nov		

REMARKS

METRIC EQUIVALENTS

CONVERSION FACTORS

The following list of convenient equivalents of measure gives the relationship between imperial units and the International System of Units (SI).

1 inch equals 2.54 cm (centimetres)	1 pouce vaut 2.54 cm (centimètres)
1 foot equals 0.3048 m (metre)	1 pied vaut 0.3048 m (mètre)
1 statute mile equals 1.6093 km (kilometres)	1 mille terrestre équivaut à 1.6093 km (kilomètre)
1 cm (centimetre) equals 0.393 70 inch	1 cm (centimètre) équivaut à 0.393 70 pouce
1 m (metre) equals 3.2808 feet	1 m (mètre) équivaut à 3.2808 pieds
1 km (kilometre) equals 0.621 37 mile	1 km (kilomètre) équivaut à 0.621 37 mille
1 acre equals 43 560 square feet	1 acre vaut 43 560 pieds carrés
1 acre equals 0.404 69 ha (hectare)	1 acre équivaut à 0.404 69 ha (hectare)
1 square mile equals 640 acres	1 mille carré vaut 640 acres
1 square mile equals 2.5900 km ² (square kilometres)	1 mille carré équivaut à 2.5900 km ² (kilomètres carrés)
1 square mile equals 259.0 ha (hectares)	1 mille carré équivaut à 259.0 ha (hectare)
1 ha (hectare) equals 10 000 m ² (square metres)	1 ha (hectare) vaut 10 000 m ² (mètres carrés)
1 ha (hectare) equals 2.4710 acres	1 ha (hectare) équivaut à 2.4710 acres
1 km ² (square kilometre) equals 0.386 10 square mile	1 km ² (kilomètre carré) équivaut à 0.386 10 mille carré
1 cubic foot equals 6.2288 imperial gallons	1 pied cube équivaut à 6.2288 gallons impériaux
1 imperial gallon equals 4.546 09 L (litres)	1 gallon impérial vaut 4.546 09 L (litres)
1 imperial gallon equals 1.2010 U.S. gallons	1 gallon impérial équivaut à 1.2010 gallon américain
1 U.S. gallon equals 0.133 68 cubic foot	1 gallon américain équivaut à 0.133 68 pied cube
1 cubic foot equals 0.068 317 m ³ (cubic metre)	1 pied cube vaut 0.068 317 m ³ (mètre cube)
1 m ³ (cubic metre) equals 35.315 cubic feet	1 m ³ (mètre cube) vaut 35.315 pieds cubes
1 cubic foot per second for one day equals 1.9835 acre-feet	1 pied cube par seconde pendant un jour équivaut à 1.9835 acre-pied
1 cubic foot per second for one day covers one square mile to a depth of 0.037 19 inch	1 pied cube par seconde pendant un jour équivaut à un volume d'un mille carré par 0.037 19 pouce
1 acre-foot equals 1.2335 dam ³ (cubic decametres)	1 acre-pied équivaut à 1.2335 dam ³ (décamètres cubes)
1 m ³ /s (cubic metre per second) for one day equals 86.4 dam ³ (cubic decametre)	1 m ³ /s (mètre cube par seconde) pour un jour vaut 86.4 dam ³ (décamètre cube)
1 m ³ /s (cubic metre per second) for one day covers one square kilometre to a depth of 0.0864 m (metre)	1 m ³ /s (mètre cube par seconde) pour un jour couvre un kilomètre carré à une profondeur de 0.0864 m (mètre)
1 foot per second equals 0.6818 mile per hour	1 pied par seconde équivaut à 0.6818 mille par heure
1 mile per hour equals 1.467 feet per second	1 mille par heure équivaut à 1.467 pied par seconde
1 m/s (metre per second) equals 3.6 km/h (kilometre per hour)	1 m/s (mètre par seconde) vaut 3.6 km/h (kilomètre par heure)
1 km/h (kilometre per hour) equals 0.2778 m/s (metre per second)	1 km/h (kilomètre par heure) vaut 0.2778 m/s (mètre par seconde)
1 cubic foot per second equals 0.028 317 m ³ /s (cubic metre per second)	1 pied cube par seconde équivaut à 0.028 317 m ³ /s (mètre cube par seconde)
1 m ³ /s (cubic metre per second) equals 35.315 cubic feet per second	1 m ³ /s (mètre cube par seconde) équivaut à 35.315 pieds cubes par seconde
1 pound equals 0.453 59 kg (kilogram)	1 livre équivaut à 0.453 59 kg (kilogramme)
1 kg (kilogram) equals 2.2046 pounds	1 kg (kilogramme) équivaut à 2.2046 livres
1 short ton (2000 pounds) equals 0.907 18 t (tonne)	1 tonne courte (2000 livres) équivaut à 0.907 18 t (tonne)
1 t (tonne) equals 2204.6 pounds	1 t (tonne) équivaut à 2204.6 livres
degrees Celsius = 5/9 (degrees Fahrenheit - 32)	dégrés Celsius = 5/9 (degrés Fahrenheit - 32)
degrees Fahrenheit = 9/5 (degrees Celsius) + 32	dégrés Fahrenheit = 9/5 (degrés Celsius) + 32

FACTEURS DE CONVERSION

Voici une liste des unités de mesure impériales et leurs équivalences dans le Système international d'unités (SI).

1 pouce vaut 2.54 cm (centimètres)	1 pied vaut 0.3048 m (mètre)	1 mille terrestre équivaut à 1.6093 km (kilomètre)
1 cm (centimètre) équivaut à 0.393 70 pouce	1 m (mètre) équivaut à 3.2808 pieds	1 km (kilomètre) équivaut à 0.621 37 mille
1 acre vaut 43 560 pieds carrés	1 acre équivaut à 0.404 69 ha (hectare)	1 mille carré vaut 640 acres
1 mille carré équivaut à 2.5900 km ² (kilomètres carrés)	1 mille carré équivaut à 259.0 ha (hectare)	
1 ha (hectare) vaut 10 000 m ² (mètres carrés)	1 ha (hectare) équivaut à 2.4710 acres	1 km ² (kilomètre carré) équivaut à 0.386 10 mille carré
1 pied cube équivaut à 6.2288 gallons impériaux	1 gallon impérial vaut 4.546 09 L (litres)	1 gallon impérial équivaut à 1.2010 gallon américain
1 gallon américain équivaut à 0.133 68 pied cube	1 pied cube vaut 0.068 317 m ³ (mètre cube)	1 m ³ (mètre cube) vaut 35.315 pieds cubes
1 pied cube par seconde pendant un jour équivaut à 1.9835 acre-pied	1 pied cube par seconde pendant un jour équivaut à un volume d'un mille carré par 0.037 19 pouce	1 acre-pied équivaut à 1.2335 dam ³ (décamètres cubes)
1 m ³ /s (mètre cube par seconde) pour un jour vaut 86.4 dam ³ (décamètre cube)	1 m ³ /s (mètre cube par seconde) pour un jour couvre un kilomètre carré à une profondeur de 0.0864 m (mètre)	
1 pied par seconde équivaut à 0.6818 mille par heure	1 mille par heure équivaut à 1.467 pied par seconde	1 m/s (mètre par seconde) vaut 3.6 km/h (kilomètre par heure)
1 km/h (kilomètre par heure) vaut 0.2778 m/s (mètre par seconde)		
1 pied cube par seconde équivaut à 0.028 317 m ³ /s (mètre cube par seconde)	1 m ³ /s (mètre cube par seconde) équivaut à 35.315 pieds cubes par seconde	
1 livre équivaut à 0.453 59 kg (kilogramme)	1 kg (kilogramme) équivaut à 2.2046 livres	
1 tonne courte (2000 livres) équivaut à 0.907 18 t (tonne)	1 t (tonne) équivaut à 2204.6 livres	
dégrés Celsius = 5/9 (degrés Fahrenheit - 32)	dégrés Fahrenheit = 9/5 (degrés Celsius) + 32	