Table of Content

S. No.	Title	Page No.
	Transformer-Based Spatio-Temporal Deep Learning Models for Solar Irradiance Forecasting in Smart Grid Applications	
1	Introduction	1
2	Literature Review	4
3	Research Gap	10
4	Objectives of the Study	11
5	Hypotheses	11
6	Statement of the Research Problem	12
7	Significant Contribution of the Proposed Research to the Field	12
8	Research Methodology	13
9	Delimitations of the Study	17
10	Proposed Plan of Research Work (Sample, Sampling Techniques, Research Tools and Techniques, Variables)	18
11	Expected Outcomes of the Proposed Research	19
12	Proposed Chapterization	20
13	Suggestions for Future Research	21
14	References	22

1. Introduction

The worldwide shift to low-carbon and sustainable energy systems has further increased the inclusion of renewable energy sources into the present-day power grids, with the solar energy turning out to be one of the most promising and fast-growing sources. The relatively decreasing prices of photovoltaic (PV) technologies coupled with the supportive government policies and increasing environmental consciousness have resulted in the wide usage of solar power generation in both the developed and the developing economies (Bouquet et al., 2024). Nonetheless, the natural fluctuation and intermittence of solar energy which are mainly caused by atmospheric and meteorological factors are major drawbacks of reliable functioning and stability of power systems. In smart grid settings, where high-quality sensing, communication and control technologies are used to increase efficiency and resiliency, proper solar irradiance prediction can be vital in determining the maximum energy management, load balancing and grid stability (Shahat et al., 2024).



Figure 1 Application of Smart grid

The productivity of solar PV systems is directly dependent on the intensity of the sunlight reaching the surface known as solar irradiance. Irregularity of irradiance caused by clouds movement, temperature, humidity changes and seasonal changes may cause a lot of uncertainty in generation of power. Wrong predictions can lead to poor planning of the traditional power plants, the use of more expensive energy reserves, unstable voltage and operational expenses. Therefore, accurate and predictive forecasting of solar irradiance

over a broad range of time frames between minutes and hours, but extending to days in the future is a major prerequisite of successful smart grid planning and functioning. This has necessitated massive research being done in order to come up with better forecasting methods, which can be able to capture the intricate dynamics of the solar energy production (Bhutta et al., 2024).

Physical and statistical methods have been the most widely used in the traditional solar irradiance forecasting methods. Physical models rely on atmospheric physics and numerical weather prediction (NWP) methods that need much computer resources and fine-scale measurements of the atmosphere. Although these models are useful in medium and long term forecasting, they tend to lose their accuracy at finer time scales and local scales. Statistical techniques such as autoregressive integrated moving average (ARIMA), regression-based models are simple and interpretable yet they cannot model non-linear relationships and complex dependencies that are imposed by the solar irradiance time series. This has encouraged the transition to more data inspired methods, especially machine learning and deep learning models, which have the ability to automatically identify patterns in past data (Suanpang, 2024).

Techniques of machine learning including the support vector machines, decision trees, and ensemble methods have shown better forecasting capabilities in past years by being able to identify non-linear relationships between the meteorological variables and solar irradiance. Based on these developments, recurrent neural networks (RNNs), long short-term memory (LSTM) and gated recurrent units (GRUs) models have become popular since they can capture time-dependent characteristics of time-series data. These models have been quite successful in the short time prediction of solar irradiance but there are some limitations associated with them. RNN based architectures are prone to vanishing gradient problems, low levels of parallelization, and decreased efficiency with respect to long input sequences. In addition, even current studies in deep learning are more centered on temporal relations in small locations neglecting the existence of spatial correlations between geographically spread solar stations (Xu et al., 2024).

Space dimension is especially relevant in smart grid implementation, where the solar power is generated in various zones. The weather systems and cloud movements are

spatially continuous, that is, irradiance conditions at a single point tend to be correlated with the adjacent positions. Such spatial interactions may cause inefficient forecasting results particularly in regional and grid scales when ignored. Although there have been studies that have tried to utilize the spatial information with the use of convolutional neural networks or graph-based models, the methods tend to be complex to implement and still may not be able to effectively capture the long-range temporal dependencies. Consequently, a dire requirement exists to have spatial and time dependent forecasting structures that are scalable and efficient in modeling both dependencies (Tajjour et al., 2023).

Deep learning models which are transformer-based and initially adapted to perform natural language processing tasks have recently become useful tools to model sequential data. Transformers is fundamentally an innovation of the self-attention mechanism, which allows the model to prioritize the value of various components in a sequence in a dynamic way, with respect to the relative positions (Chen et al., 2023). Transformers unlike recurrent architectures are computationally efficient and high scalable since they provide parallel processing of input sequences. These qualities put Transformer models in a good position to handle the long range dependencies and complex interactions in time-series data. Although proven to be effective in other areas like language translation, speech recognition, and computer vision, the use of Transformer-based architecture in solar irradiance forecasting particularly in a spatio-temporal setting is comparatively underutilized (Aslam et al., 2021).

The capability of Transformers to reproduce not only long-term time trends but also spatial patterns at different places offers a great chance to improve the accuracy and reliability of the forecast. Transformer-based spatio-temporal models are able to use the combination of data in geographically dispersed solar stations and also include the meteorological variables to make more precise and stronger forecasts of the solar irradiance (Bamisile et al., 2020). These advancements relate directly to the smart grid operation, such as rationality in energy dispatch, reserves, enhanced integration of renewable resources and the stability of the grid. In addition, precise forecasting can aid

decision-making in grid operators, policymakers, and participants in the energy markets, which serves the bigger objectives of sustainable energy transition (Technologies, 2020).

In this environment, the current research suggests the creation and testing of the Transformer-based spatio-temporal deep learning frameworks to predict the solar irradiance in smart grid operations. Based on high-resolution, multi-site solar irradiance and meteorological data provided in the National Solar Radiation Database, the research will undertake a systematic exploration of the utility of self-attention systems in the dominant spatial and temporal dependencies (Approach, 2019). The suggested models undergo a strict comparison with the conventional statistical frameworks, machine learning algorithms and recurrent deep-learning models to show their respective strengths. This study can help to solve the existing weaknesses in the solar irradiance prediction and improve the current achievements of deep learning to develop intelligent data-driven solutions to the next-generation smart grids and make the successful integration of solar energy into the existing power systems (Wan et al., 2015).

2. Review of Literature

(Aljohani, 2026) discuss the drawbacks of conventional statistical forecasting tools to simulate the nonlinear and weather-sensitive behaviour of renewable energy systems. To overcome this, the authors suggest an AI-based spatio-temporal model that includes CNNLSTM on the forecasting of renewable generation, Gradient Boosting Machine on the forecasting of demand, and Particle Swarm Optimization on the cost-effective allocation of energy. It was shown that the framework is strongly generalized as it was validated on national hourly energy data in Spain and also tested on the energy market in Germany. The findings indicate a high predictive accuracy with solar generation having an R 2 of 99.03. Also, the PSO-related optimization minimized fossil fuel dependency and enhanced the efficiency of the grid, highlighting the importance of AI in the energy management of the smart city.

(Saravana et al., 2026) offer an extensive literature review of Transformer-based models of multivariate time-series forecasting that resolves the criticisms of standard deep learning models in their ability to deal with long-term dependencies and large data

volumes. The paper methodically provides a review of early Transformer networks, efficient attention models, decomposition and patch-based models, hybrid models, and large-scale Transformer pretrained models. A hierarchical dataset taxonomy, metrics of evaluation and domains of application are provided, which provides researchers with a clear picture. Of interest is also the discussion of the major challenges, such as computational complexity, interpretability as well as data sparsity as well as future research directions such as spatio-temporal modeling, physics-informed learning, and federated frameworks which makes the survey very relevant in the research of renewable energy forecasting.

(Sarkar, 2025) propose a deep learning-based time-series forecasting framework to capture nonlinear relationships, spatio-temporal relationships, and changing patterns in time. The paper combines ensemble learning, guided networks, interactive convolution, self-supervised learning, Transformer networks, meta-learning strategies, and Bayesian Optimization. A new directed ensemble method, known as GATE, is a form of unsupervised learning that adapts a contribution of the models in real time to enhance model stability and prediction errors. The results of the experiments show great performance gains, as the performance is up to 17.5 percent higher than that of RNNs and 20.9 percent higher than LSTMs. The results highlight the usefulness of ensemble-based Transformer-based models with complex forecasting problems.

(Rastgoo & Amjady, 2025) present a probabilistic forecasting system of ultra-short-term photovoltaic power prediction using cloud images based on diffusion. The suggested Cross Branch Visual Informer (CB-ViInf) is a combination of the multi-resolution sky image patch-based solution with the Vision Informer architecture to boost the feature extraction and time modeling. Dual block temporal encoder using Spatiotemporal Ridgelet Transform and spatio-temporal attention is effective in capturing the rapid irradiance changes due to the dynamic nature of clouds. Also, there is a new ELBO loss function that enhances the quantification of uncertainty. Wide comparisons with 64 baseline models validate a higher level of forecasting accuracy and strength making the method a state of the art solution to uncertainty-aware power forecasting of the sun.

(Cavus, 2025) present ST-CALNet, a spatio-temporal CNN-LSTM encoder-decoder framework, an inbuilt attention to forecast energy demand and renewable generation. CNN element acquires spatial relationships of multivariate meteorological and generation input, and LSTM coefficients acquire temporal connections within historical data. The attention system is a dynamically time-weighted mechanism that makes it easier to interpret and robust to demand changes. Real-world dataset assessment showed that the predictive accuracy of the proposed models was superior to the baseline models, and both MAE and the RMSE values were low in predicting electricity demand, PV output, and wind generation. The findings indicate that the model is applicable to smart grid systems when it comes to adaptive and transparent forecasting.

Table 2.1 Literature Summary

Author / Year	Methodology	Key Findings	Research Gap Identified	Limitations
(Yang et al., 2025)	Proposed Wavelet–Graph Learning (WGL) framework integrating learnable wavelet shrinkage, temporal multi-scale fusion, graph self-attention, and factorized spatio-temporal attention for multi-site PV forecasting	Joint multi-site forecasting significantly outperformed deep and graph-based baselines for intra-hour PV prediction; effectively captured spatio-temporal correlations	Focused primarily on PV power output; limited exploration of generalized solar irradiance forecasting across diverse smart grid horizons	High architectural complexity; computational overhead may limit scalability for real-time smart grid deployment

(Li et al., 2025)	Data-driven predictive analytics framework using decision trees, deep neural networks, and ensemble	Achieved up to 40% MAE reduction and 25% R^2 improvement over traditional models; improved grid resilience forecasting	Did not explicitly model long-range spatio-temporal dependencies using attention-based architectures	Transformer-based or spatial attention mechanisms were not explored; interpretability remains limited
(Kim et al., 2025)	Transfer learning-based CL-Transformer integrating weather and satellite data across multi-country PV plants	Significant R^2 improvements (up to 67.3%) across different plant scales; strong generalization across regions	Emphasis on PV capacity prediction rather than continuous irradiance forecasting for smart grid operations	Requires pretrained source-domain datasets; limited evaluation on multi-horizon grid forecasting
(Ramu et al., 2025)	Transformer-based climate forecasting with hierarchical graph representation and dynamic temporal graph attention, optimized using	Improved accuracy, recall, and F1-score with reduced training time; effective for climate forecasting	Model applied to temperature prediction, not solar irradiance or smart grid energy forecasting	Validation restricted to a single climatic region; limited renewable energy application

	hybrid metaheuristics			
(Saeed et al., 2025)	Transformer-based spatio-temporal framework combining Chebyshev graph convolution and multi-head attention for natural gas demand forecasting	Achieved high R^2 (0.93–0.96) and strong MAE performance under climatic volatility	Focused on gas demand forecasting rather than solar irradiance or renewable integration	Sector-specific application; lacks direct evaluation in solar or smart grid contexts
(Xu et al., 2024)	Vision Transformer (ViT) + GRU framework using sky-imager cloud images for ultra-short-term PV forecasting	Outperformed baseline models for ultra-short-term PV generation prediction	Limited to ultra-short-term forecasting and image-based data; lacks long-term spatio-temporal modelling	Requires specialized imaging infrastructure; limited generalization to non-image-based smart grids
(Abubakar et al., 2024)	Machine learning and RNN-based models (RNN, LSTM, GRU) for solar power	LSTM achieved highest accuracy (≈97%), improving solar power	Did not explore advanced attention-based or Transformer architectures	High hardware requirements; weak modeling of long-range dependencies

	prediction in smart grids	generation efficiency		and spatial correlations
(Boussif & Massaroli, 2023)	Spatio-temporal deep learning using satellite imagery and time-series data with uncertainty-aware forecasting	Demonstrated robust GHI forecasting and zero-shot generalization to unseen locations	Limited use of Transformer-based self-attention for explicit multi-site temporal learning	Computational cost of satellite data processing; limited smart grid operational validation

3. Research Gap

Proper operation of smart grids is based on to predict solar irradiance efficiently, requiring the historic accuracy of smart grids in the high penetration of solar photovoltaic generation. The available literature has talked at length about statistical models, machine learning models and recurrent deep learning models like LSTM and GRU as a predictor to solar irradiance. Although the performance of these approaches has been found to be reasonable on short-term forecasting, there are some severe limitations that have not been addressed.

To begin with, most studies that are currently available are mostly concerned with the temporal modeling of individual locations, thus overlooking the spatial interdependence of geographically dispersed sun sites. These spatial correlates play a very important role in smart grid setting where weather systems and cloud dynamics affect numerous locations at a time. Second, recurrent neural network-based models have disadvantages of long-range dependency learning, vanishing gradients, and lack of parallelization that limit their ability to predict long-term and large data sets.

Moreover, even though Transformer-based architectures have shown impressive performance in the modeling of long-range dependencies in other tasks, they have not

been used yet to forecast spatio-temporal solar irradiance using large-scale, multi-year databases like the National Solar Radiation Database (NSRDB). No detailed comparative studies are also found to systematically compare Transformer-based models with traditional statistical, machine learning, and recurrent deep learning under realistic conditions of smart grids.

Thus, there is an apparent research gap in creating and testing Transformer-based spatio-temporal deep learning systems that will successfully combine spatial and temporal data to enhance the accuracy of solar irradiance forecasts and enable smart grid activities.

4. Objectives of the Study

The primary objective of this study is to develop and evaluate Transformer-based spatio-temporal deep learning models for accurate solar irradiance forecasting to enhance smart grid operations. The specific objectives are as follows:

1. To analyze the temporal and spatial characteristics of solar irradiance and associated meteorological parameters in smart grid environments.
2. To design a Transformer-based spatio-temporal deep learning architecture capable of capturing long-term temporal dependencies and spatial correlations among multiple solar measurement sites.
3. To compare the performance of the proposed Transformer-based models with conventional forecasting approaches, including machine learning, and RNN-based deep learning models.
4. To evaluate the forecasting accuracy of the proposed models across multiple time horizons (short-term and long-term).
5. To assess the applicability of the proposed forecasting framework in smart grid operations such as energy management, load balancing, and grid stability.

5. Hypotheses

Based on the objectives and research gap identified, the following hypotheses are formulated:

H_1: Transformer-based spatio-temporal deep learning models provide significantly higher forecasting accuracy for solar irradiance compared to traditional statistical and machine learning models.

H_2: Incorporation of spatial information from multiple geographically distributed locations improves solar irradiance forecasting performance compared to single-site temporal models.

H_3: Transformer-based models outperform recurrent deep learning models (LSTM and GRU) in capturing long-term temporal dependencies in solar irradiance data.

H_4: Improved solar irradiance forecasting accuracy positively enhances smart grid operations, including energy management, load balancing, and grid stability. .

6. Statement of the Research Problem

The growing demand to incorporate solar energy in smart grids presents a lot of trouble in terms of its operation because solar irradiance is intermittent and most volatile. Imprecise prediction of the irradiance of the sun causes inefficient scheduling of energy, more dependence on reserve generation of power, and decreased grid stability. Current methods of forecasting do not sufficiently characterize complicated spatio-temporal relationships and long-term time trends found in multi-location solar irradiance observations.

The core research problem addressed in this study is:

How can Transformer-based spatio-temporal deep learning models be effectively designed and applied to improve the accuracy and reliability of solar irradiance forecasting for smart grid applications?

7. Significant Contribution of the Proposed Research to the Field

The proposed research makes several significant contributions to the field of renewable energy forecasting and smart grid analytics:

1. Development of a Transformer-based spatio-temporal forecasting framework tailored for solar irradiance prediction.
2. Integration of multi-location solar irradiance and meteorological data to explicitly model spatial dependencies.
3. Comprehensive comparative evaluation against traditional statistical, machine learning, and recurrent deep learning models.
4. Empirical validation using the National Solar Radiation Database (NSRDB), ensuring robustness and reproducibility.
5. Practical insights into the application of advanced deep learning models for smart grid energy management and decision support.

8. Research Methodology

The current research proposal is a systematic systematic and data-intensive research approach that will design, develop, and test the Transformer-based spatio-temporal deep learning models to predict solar irradiance in smart grid systems. The methodology combines the multi-source acquisition of data, intensive pre-processing, exploratory data analysis, baseline model benchmarking, advanced Transformer-based architecture training, model training and optimization, and specific performance analysis. The general system design is a framework that is scientifically sound, reproducible and relevant in a practical sense in terms of forecasting renewable energies and intelligent grid management.

3.1 Research Design

The research is based on the quantitative, experimental and comparative research design as it is concerned with predictive modeling utilizing real-world solar irradiance and meteorological data. It is an exploratory study because it examines whether Transformer-based architectures can be used to model complex spatio-temporal relationships, and comparative because they assess their accuracy in comparison to existing statistical, machine learning, and deep learning forecasting models.

Experimental workflow is both sequential and iterative with the initial phase being dataset selection and preprocessing, exploratory data analysis, feature engineering, baseline development and training, validation and performance analysis. All the stages depend on each other, so the data understanding is followed by systematic development of the models of verification and practical interpretation to be applied in the smart grid.

3.2 Dataset Selection and Data Sources

The main source of data that is used in this research is the National Solar Radiation Database (NSRDB). The reason behind the choice of NSRDB is its extensive spatio-temporal coverage, excellent data quality, and great popularity in the research on renewable energy and smart grids.

Dataset Link: https://developer.nrel.gov/docs/solar/nsrdb/

The dataset offers long-term and high resolution solar irradiance data, i.e. Global Horizontal Irradiance (GHI), Direct Normal irradiance (DNI), and Diffuse Horizontal Irradiance (DHI), and related meteorological parameters, i.e. ambient temperature, relative humidity, wind speed, wind direction, surface pressure, and cloud-related parameters. Information is present at hourly time scales and over several years making it to effectively model daily, seasonal and interannual changes.

In order to obtain the spatial dependencies to be used in the spatio-temporal forecasting, data are taken out of various geographically dispersed areas, which reflect varying weather and weather patterns of the sun. This multi-site architecture facilitates the creation of spatial attention processes in Transformer-based architectures and concurs with smart grid scenarios in real-world settings with distributed solar generation.

3.3 Data Preprocessing

Raw data that is retrieved through NSRDB can be missing as well as have noise and inconsistencies due to sensor restrictions or satellite estimation. Thus, a powerful data preprocessing pipeline is provided in order to improve the data quality and model reliability.

Time-series-intelligent imputation methods like linear interpolation and rolling statistical estimations are used to deal with missing values based on the duration and frequency of the missing chunks. The statistical methodologies applied to detect outliers are the interquartile range (IQR) method and the z-score analysis of the irradiance or the meteorological data to identify extreme values. Outliers are identified and eliminated to avoid distortion of learning of models.

All continuous variables are scaled between Min and Max to normalize them or they are scaled to Z-score in order to guarantee the numerical stability and effective convergent of model training. Temporal alignment is done to align all variables across locations and time so that there is a consistency in the representation of spatio-temporal input.

3.4 Exploratory Data Analysis (EDA)

Exploratory Data Analysis is performed in order to get an idea on the statistical and temporal peculiarities of the NSRDB data. Descriptive statistics are calculated to investigate the distribution, variability and central tendencies of the solar irradiance and meteorological variables. The time-series visualizations are employed to recognize the daily cycles, seasonal patterns, and the long-term trends.

To test how the irradiance components relate to meteorological variables, correlation analysis is done. Seasonal decomposition methods are used to extract trend, seasonal, and residual term of the irradiance series. Besides, spatial correlation analysis at various sites is performed to measure inter-site dependencies, which proves the necessity of spatio-temporal modeling. The EDA results support the choice of features, lag window, and architecture of forecasting models.

3.5 Feature Engineering and Input Representation

It is essential to have effective feature engineering to increase the accuracy of predictions. Lagged values of solar irradiance and meteorological parameters are developed in this analysis to represent the time dependencies. Categorical features that are based on the time of the day, in the week or month of the year are represented to capture the periodic and seasonal trends.

In the case of spatial representation, the data in various locations are integrated into multi-dimensional input tensors, with each geographic location being the spatial node. Positional encoding is used to retain the information of the temporal order, which is needed by Transformer-based self-attention mechanisms. The resulting input structure is a combination of the spatial and temporal dimension, which allows the model to learn complex cross-location and long-range temporal interactions.

3.6 Baseline Model Development

In order to determine the performance benchmarks, a number of baseline forecasting models are established and tested on the NSRDB dataset. These are traditional statistical models like the AutoRegressive Integrated Moving Average (ARIMA), machine learning models like the Support Vector Machines (SVM) and Random Forests, and recurrent neural network models, like the Long Short-Term Memory (LSTM) and the Gated Recurrent Unit (GRU).

These models are chosen based on the fact that they experience wide application in the literature of forecasting solar irradiance. The entire baseline models are trained and tested on the same data partitions and are measured based on the same performance measures to obtain an impartial and equal comparison to the proposed Transformer-based models.

3.7 Transformer-Based Spatio-Temporal Model Architecture

The major work of this study is the Transformer-based spatio-temporal forecasting models. Transformers use self-attention mechanisms that enable the model to dynamically balance the significance of various time-steps and spatial positions to enable successful learning of long-range relationships.

The suggested structure is made of multi-head self-attention, position-wise feed-forward, residual, and layer normalization. The learning of spatial dependencies involves processing of inputs across different geographic positions concurrently to enable the attention heads to acquire inter-site relationships. The temporal positional encoding is included to retain order of sequences information. Such architecture is highly scalable and parallel as compared to recurrent models.

3.8 Model Training and Optimization

Adam optimizer is used as a model training by its adaptive learning rate and convergence speed features. Temporal splitting is used to split the dataset into the training, validation, and testing subsets to avoid the leakage of any information and make the forecasting evaluation realistic.

The grid search or cross-validation does hyperparameter tuning, which targets parameters: learning rate, batch size, attention heads, and Transformer layers and hidden dimensions. Such regularization methods as dropout, early stopping, and weight decay are used to reduce overfitting and improve overall generalization.

3.9 Performance Evaluation Metrics

All the models are compared in terms of the standard regression measures such as Mean Absolute Error (MAE), Root Mean Square Error, Mean Absolute Percentage Error (MAPE) and the coefficient of determination (R 2). These measures offer complementary measurements of prediction accuracy, magnitude of error and explanatory power. It is done through comparative analysis over various forecasting horizons and seasonal conditions to determine the strength and uniformity of the model.

3.10 Result Analysis and Smart Grid Implications

The last phase of the methodology will be the analysis of the experimental data and interpretation of the implications on the application of the smart grid. Transformer-based models are improved to assess their advancements in terms of increased forecasting accuracy, reliability, and scalability. The practical application of precise irradiance

prediction is addressed as to the importance of energy scheduling, grid stability, integration of renewables, and smart grid operators.

This holistic research approach will make sure the proposed Transformer-based spatio-temporal models are strictly proved and more feasible to be applied to next-generation smart energy systems.

9. Delimitations of the Study

The study is subject to the following delimitations:

1. The analysis is limited to solar irradiance forecasting and does not directly model photovoltaic power output.
2. Only historical data from the NSRDB dataset are considered; real-time data integration is outside the scope of this study.
3. Forecasting is performed at hourly temporal resolution; ultra-short-term (minute-level) forecasting is not addressed.
4. The study focuses on selected geographic locations and does not cover all climatic regions globally.
5. Economic and policy-related impacts of improved forecasting are not explicitly analyzed.

10. Proposed Plan of Research Work

The plan of research work proposes the systematic approach, which should be followed to ensure that the proposed methodology is rigorous and analytically sound. It includes the sampling approach, the tools and methods used to do the research and the selection of the variables used to come up with, test and prove the developed Transformer-based spatio-temporal models in predicting solar irradiance in smart grid applications.

Sampling Techniques

The purposive sampling method is taken to select several locations that are geographically separated based on the NSRDB dataset. Places are selected to be representative of varying climatic and solar conditions of relevance to smart grid activities.

Research Tools and Techniques

1. Programming Environment: Python
2. Deep Learning Frameworks: TensorFlow / PyTorch
3. Statistical Tools: NumPy, Pandas, SciPy
4. Visualization: Matplotlib, Seaborn
5. Models: ARIMA, SVM, Random Forest, LSTM, GRU, Transformer

Variables

1. **Dependent Variable:** Solar irradiance (GHI, DNI, DHI)
2. **Independent Variables:** Temperature, humidity, wind speed, pressure, cloud cover, temporal features
3. **Control Variables:** Geographic location, temporal resolution, data preprocessing techniques

11. Expected Outcomes of the Proposed Research

the results of this study are supposed to contribute to the sector of renewable energy forecasting and smart grid management both theoretically and practically. To begin with, the research is likely to realize far better prediction results of solar irradiance over various time horizons such as short-term and long-term predictions. Realistic predictions of irradiance will facilitate credible power scheduling and decrease the unpredictability of generation of solar energy.

Second, the study is likely to provide empirical proof of the superiority of the Transformer-based spatio-temporal deep learning models over the traditional forecasting models, including statistical models, machine learning algorithms, and the models that are

based on recurrent neural networks. The effectiveness of self-attention mechanism in conveying long-range temporal dependencies and spatial relations will be pointed out through the systematic performance comparison provided by the study.

Third, the research will aid in the understanding of complex spatio-temporal patterns that the solar irradiance data have by investigating the interdependencies between various locations with geographical distribution and meteorological variables. This enhanced understandability will enhance model design policies towards renewable energy forecasting.

The study is likely to provide a scalable, robust and versatile forecasting model that could be implemented into smart grid settings. Lastly, the results will offer viable ideas to grid operators and energy planners so that they can integrate renewable energy in a better manner, optimize grid operations, increase reliability, and make effective decisions regarding sustainable energy systems.

12. Proposed Chapterization

To achieve the objectives of the study, the proposed research work is well structured in five well-organized chapters to achieve logical flow, conceptual clarity, and methodological rigor.

Chapter 1: Introduction This chapter presents the research problem, which has led to increased relevance of proper solar irradiance prediction in smart grid applications. It gives the background of the integration of renewable energy, the reasons why Transformer-based spatio-temporal models have been adopted, the research objectives, hypotheses, scope and relevance of the research. The structure of the thesis is also shown in the chapter.

Chapter 2: Review of Literature This chapter is a thorough literature review on the available literature on solar irradiance forecasting, smart grids, deep learning models, and Transformer-based architectures. It is a critical analysis of the conventional statistical methods, machine learning solutions, and advanced spatio-temporal models that illustrate the gaps and limitations in a research that support the proposed study.

Chapter 3: Research Methodology This chapter provides information on the methodological framework that the study has adopted such as research design, data set selection, pre-processing of data, exploratory data analysis, feature engineering, base model construction, Transformer-based model architecture, training plan, and evaluation metrics. There is focus in the reproducibility and scientific rigor.

Chapter 4: Results and Discussion The chapter shows the results of the experiment using baseline and proposed models. The performance analysis is made based on various forecasting horizons, and the nature of the study then presented is the discussion of the spatio-temporal learning behavior, model robustness, and implications of the model to the smart grid operations.

Chapter 5: Conclusion and Future Scope The concluding chapter will summarize the significant findings, dwell on the key contributions of the paper, limitation, and future research directions in the more advanced renewable energy forecasting and intelligent grid control.

13. Suggestions for Future Research

Future research recommendations are given to broaden the current research and to combat new challenges in the solar irradiance forecasting and smart grid application. These guidelines are used to increase the model strength, flexibility, and application in real-world energy systems.

1. Included in multi-modal sources of data that can be incorporated into future studies are satellite images, sky camera images, numerical weather prediction images, and streams of Internet of Things sensors. The combination of heterogeneous information through the sophisticated attention systems can enhance the depiction of atmospheric dynamics and the accuracy in forecasting of the weather given the rapidly varying weather patterns.

2. Transformer-based architectures can be enhanced by incorporating the physical concepts of the solar radiation and atmospherical processes to enhance the interpretability and reliability of the model. Deep learning methods based on

physics can minimize the need of huge datasets and improve extrapolation to new climatic areas.

3. Additional studies could be done on real-time and online learning models which can allow continuous adjustment of the model to change in environmental and grid conditions. These adaptive models would be more appropriate in the operational smart grid applications that would need timely and dynamic forecasting.

4. The potential future research is to consider uncertainty measurement and probabilistic prediction techniques as a means of producing confidence levels and point estimates. This would facilitate making informed decisions about energy scheduling, grid stability and integration of renewable.

5. Further studies can examine scalable, computationally efficient Transformer architectures to be used in large scale and edge deployment, which is practical but still achieves high forecasting performance.

References

Abubakar, M., Che, Y., Faheem, M., Bhutta, M. S., & Mudasar, A. Q. (2024). *Intelligent Modeling and Optimization of Solar Plant Production Integration in the Smart Grid Using Machine Learning Models*. *2300160*, 1–19. https://doi.org/10.1002/aesr.202300160

Aljohani, R. M. (2026). *Predictive Models for Renewable Energy Generation and Demand in Smart Cities : A Spatio-Temporal Framework*.

Approach, L. (2019). *Day-Ahead Solar Irradiance Forecasting for Microgrids Using a Long Short-Term Memory Recurrent Neural Network : A Deep*.

Aslam, S., Herodotou, H., & Muhammad, S. (2021). *A Survey on Deep Learning Methods for Power Load and Renewable Energy Forecasting in Smart Microgrids*.

Bamisile, O., Oluwasanmi, A., Obiora, S., Osei-, E., Asoronye, G., & Huang, Q. (2020). *Environmental Effects Application of deep learning for solar irradiance and solar photovoltaic multi-parameter forecast*.

Bhutta, M. S., Li, Y., Abubakar, M., Almasoudi, F. M., Alatawi, K. S. S., Altimania, M. R., & Barashi, M. Al. (2024). Optimizing solar power efficiency in smart grids using hybrid machine learning models for accurate energy generation prediction. *Scientific Reports*, 1–25. https://doi.org/10.1038/s41598-024-68030-5

Bouquet, P., Jackson, I., Nick, M., & Kaboli, A. (2024). AI-based forecasting for optimised solar energy management and smart grid efficiency ABSTRACT. *International Journal of Production Research*, *62*(13), 4623–4644. https://doi.org/10.1080/00207543.2023.2269565

Boussif, O., & Massaroli, S. (2023). *Improving day-ahead Solar Irradiance Time Series Forecasting by Leveraging Spatio-Temporal Context*. *NeurIPS*, 1–26.

Cavus, M. (2025). *Spatio-Temporal Attention-Based Deep Learning for Smart Grid Demand Prediction*. 1–24.

Chen, Y., Bhutta, M. S., Abubakar, M., Xiao, D., Almasoudi, F. M., Naeem, H., & Faheem, M. (2023). *Evaluation of Machine Learning Models for Smart Grid Parameters : Performance Analysis of ARIMA and Bi-LSTM*.

Kim, B., Member, A., & Belkilani, K. (2025). Integrated Spatiotemporal Hybrid Solar PV Generation Forecast Between Countries on Different Continents Using Transfer Learning Method. *IEEE Access*, *13*(October 2024), 2486–2502. https://doi.org/10.1109/ACCESS.2024.3514098

Li, Z., Liu, S., & Liu, G. (2025). Improving Power Grid Disaster Resilience Through Predictive Analytics Based on Data-Driven Photovoltaic Power Plant and Distribution Network Power Forecasting. *IEEE Access*, *13*(August), 176157–176175. https://doi.org/10.1109/ACCESS.2025.3611104

Ramu, T. B., Kocherla, R., Sirisha, G. N. V. G., & Chetana, V. L. (2025). *Transformer based models with hierarchical graph representations for enhanced climate*

forecasting. 1–22.

Rastgoo, R., & Amjady, N. (2025). A Diffusion-Based Probabilistic Ultra-Short-Term Solar Power Prediction Using the Sky Image Sequences. *IEEE Access*, *13*(August), 149532–149553. https://doi.org/10.1109/ACCESS.2025.3600713

Saeed, M. S., Raza, S., & Ahmad, H. (2025). Transformer-Based Spatio-Temporal Framework for Forecasting Natural Gas Demand in the Power Sector. *IEEE Access*, *13*(November), 188432–188446. https://doi.org/10.1109/ACCESS.2025.3627150

Saravana, M. K., Member, S., & Roopa, M. S. (2026). Transformers for Multivariate Time Series Forecasting : Comprehensive Analysis , Challenges , Research Opportunities , and Future Prospects. *IEEE Access*, *14*(December 2025), 11424–11457. https://doi.org/10.1109/ACCESS.2026.3654408

Sarkar, M. R. (2025). *Deep Learning Techniques Improvement for Time Series Forecasting*.

Shahat, E., Data, B., Shahat, D. El, Tolba, A., Abouhawwash, M., & Basset, M. A. (2024). Machine learning and deep learning models based grid search cross validation for short - term solar irradiance forecasting. *Journal of Big Data*. https://doi.org/10.1186/s40537-024-00991-w

Suanpang, P. (2024). *Machine Learning Models for Solar Power Generation Forecasting in Microgrid Application Implications for Smart Cities*.

Tajjour, S., Chandel, S. S., Alotaibi, M. A., Malik, H., Member, S., Pedro, F., Márquez, G., & Afthanorhan, A. (2023). *Short-Term Solar Irradiance Forecasting Using Deep Learning Techniques : A Comprehensive Case Study*. *September*, 119851–119861. https://doi.org/10.1109/ACCESS.2023.3325292

Technologies, C. (2020). *Advanced Deep Learning Methodologies for Renewable Energy Sources*.

Wan, C., Zhao, J., Member, S., & Song, Y. (2015). *Photovoltaic and Solar Power Forecasting for Smart Grid Energy Management*. *1*(4), 38–46.

Xu, S., Zhang, R., Ma, H., Ekanayake, C., & Cui, Y. (2024). On vision transformer for ultra-short-term forecasting of photovoltaic generation using sky images. *Solar Energy*, *267*(November 2023), 112203. https://doi.org/10.1016/j.solener.2023.112203

Yang, G., Xiao, J., Zhang, C., Yang, D., & Li, C. (2025). *A Real-Time Photovoltaic Power Estimation Framework Based on Multi-Scale Spatio-Temporal Graph Fusion*. 1–19.