

OPERATION D


Add and subtract fractions with friendly but unlike denominators (e.g., 2 and 10), using models and symbols

SAMPLE 1

$\frac{5^{\times 2}}{8} 8(16) \mid \frac{0}{16} + \frac{3}{16} + \frac{8}{16} - \frac{21}{16} + 8 \text{ wholes} = 8 \frac{21}{16}$
 $\frac{13^{\times 4}}{16} (16)$
 $\frac{821}{16} - 9 \frac{8}{16} = 1 \frac{13}{16}$
 $\frac{7^{\times 8}}{2} 2468101214(16)$
 $\frac{10^{\times 8}}{2} 2468101214(16)$
 $\frac{5^{\times 4}}{4} 4812(16)$
 $\frac{821}{16} - 5 \frac{12}{16} = 3 \frac{9}{16}$


then it might have been much easier to subtract the final numbers.

SAMPLE 2



Students used both the number line and common denominators for their proof. They used mixed, improper, and proper fractions interchangeably and were able to correctly place the thicknesses additively onto the number line, using the smaller thicknesses first and finishing with the largest thickness.

SAMPLE 3



These students were successfully able to place the various thicknesses on the number line and find the unknown thickness without finding common denominators. They moved fluently between fractions with friendly but unlike denominators. They started with the largest thickness first and moved to the smaller thicknesses.