

1 **Running head:** CRANIAL MORPHOLOGICAL DIVERSITY IN  
2 TENRECS

3 Morphological diversity in tenrecs  
4 (Afrosoricida, Tenrecidae): Comparing  
5 tenrec skull diversity to their closest  
6 relatives

7 Sive Finlay<sup>1,2</sup> and Natalie Cooper<sup>1,2,\*</sup>

8 <sup>1</sup> School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland.

9 <sup>2</sup> Trinity Centre for Biodiversity Research, Trinity College Dublin, Dublin 2, Ireland.

10 \*Corresponding author: ncooper@tcd.ie; Zoology Building, Trinity College Dublin,  
11 Dublin 2, Ireland.

12 Tel: +353 1 896 1926.

13 **Keywords:** geometric morphometrics, golden moles, morphological  
14 diversity, tenrecs

## 15 Introduction

16 Analysing patterns of morphological diversity has important implications  
17 for our understanding of ecological and evolutionary traits. For example,  
18 from a functional ecology perspective, morphological characteristics of  
19 limbs inform us about locomotory style (e.g. Bou et al., 1987) and the  
20 trophic niches associated with particular dental morphologies affect  
21 speciation and diversification rates through time (Price et al., 2012).  
22 Morphological diversity is also an important aspect of evolutionary  
23 patterns such as adaptive radiations and convergent evolution. High  
24 morphological diversity is a unifying (Losos and Mahler, 2010; Olson and  
25 Arroyo-Santos, 2009), although not defining (Glor, 2010; Olson and  
26 Arroyo-Santos, 2009), characteristic of adaptive radiations. Furthermore,  
27 analysing morphological convergences in groups such as freshwater  
28 cichlid fish (Muschick et al., 2012) and anole lizards (Mahler et al., 2013)  
29 gives interesting insights into the relative repeatability of evolution (Losos,  
30 2011).

31 Although studies of morphological diversity have clear implications  
32 for our understanding of ecological and evolutionary patterns, apart from  
33 a few examples (e.g. Ruta et al., 2013; Goswami et al., 2011; Brusatte et al.,  
34 2008), it is still common to study morphological diversity from a  
35 qualitative rather than quantitative perspective. However, we need to  
36 quantify the morphological similarities and differences among species to  
37 gain a better understanding of their ecological interactions and  
38 evolutionary history. Unfortunately, morphological diversity is difficult to  
39 quantify. Studies are inevitably constrained to measure the diversity of  
40 specific traits rather than overall morphologies (Roy and Foote, 1997). In

41 addition, our perception of morphological diversity is influenced by the  
42 trait being used. One study of pterosaurs demonstrated that comparing  
43 the diversity of different morphological traits using varying methods  
44 produced similar results (Foth et al., 2012). However, it remains unclear  
45 whether this finding can be applied to all vertebrate groups: in some  
46 species, comparing the relative diversity of cranial and limb morphologies  
47 may yield different results (Foth et al., 2012). Furthermore, linear  
48 measurements of morphological traits can restrict our understanding of  
49 overall morphological variation. A distance matrix of measurements  
50 between specific points is unlikely to give a completely accurate  
51 representation of a three dimensional structure (Rohlf and Marcus, 1993).

52 These are important limitations to consider but geometric  
53 morphometric approaches help to overcome some of the issues associated  
54 with traditional morphological studies (Adams et al., 2004).  
55 Morphometric studies based on caliper measurements of particular  
56 features can only describe a limited set of distances, ratios and angles  
57 which often fail to capture the overall shape of a specific structure (Slice,  
58 2007). Geometric morphometrics circumvents these issues by using a  
59 system of Cartesian landmark coordinates to define anatomical points.  
60 This method captures more of the true, overall anatomical shape of  
61 particular structures (Mitteroecker and Gunz, 2009). These more detailed  
62 approaches are useful tools for studying patterns of morphological  
63 diversity.

64 Here we apply geometric morphometric techniques to quantify  
65 morphological diversity in a Family of small mammals, the tenrecs.  
66 Tenrecs (Afrosoricida, Tenrecidae) are a morphologically diverse group  
67 that is commonly cited as an example of both convergent evolution and an

68 adaptive radiation (Soarimalala and Goodman, 2011; Eisenberg and  
69 Gould, 1969). The Family is comprised of 34 species, 31 of which are  
70 endemic to Madagascar (Olson, 2013). Body masses of tenrecs span three  
71 orders of magnitude (2.5 to  $\geq 2,000$ g); a greater range than all other  
72 Families, and most Orders, of living mammals (Olson and Goodman,  
73 2003). Within this vast size range there are tenrecs which convergently  
74 resemble shrews (*Microgale* tenrecs), moles (*Oryzorictes* tenrecs) and  
75 hedgehogs (*Echinops* and *Setifer* tenrecs, Eisenberg and Gould, 1969). Their  
76 similarities include examples of morphological, behavioural and  
77 ecological convergence (Soarimalala and Goodman, 2011). Tenrecs are one  
78 of only four endemic mammalian clades in Madagascar and the small  
79 mammal species they resemble are absent from the island (Garbutt, 1999).  
80 Therefore, it appears that tenrecs represent an adaptive radiation of  
81 species which filled otherwise vacant ecological niches (Soarimalala and  
82 Goodman, 2011). The similarities among tenrecs and other small  
83 mammals are even more remarkable when you consider their  
84 phylogenetic history. Tenrecs were originally classified within the general  
85 "Insectivora" clade and only molecular studies revealed their true  
86 phylogenetic affinities within the Afrotherian mammals (Stanhope et al.,  
87 1998). Therefore, despite initial appearances, tenrecs are more closely  
88 related to elephants, manatees and aardvarks than they are to shrews,  
89 moles or hedgehogs.

90 Although tenrecs are often cited as an example of both an adaptive  
91 radiation and exceptional convergent evolution, these claims have not  
92 been investigated quantitatively. There are qualitative similarities among  
93 the hind limb morphologies of tenrecs and several other unrelated species  
94 with similar locomotory styles (Salton and Sargis, 2009) but the degree of

95 morphological similarity has not been established. Morphological  
96 diversity is an important feature of adaptive radiations (Losos and Mahler,  
97 2010) and it also informs our understanding of convergent phenotypes  
98 (Muschick et al., 2012). Therefore, it is important to quantify patterns of  
99 morphological diversity in tenrecs to gain an insight into their evolution.

100 We present the first quantitative study of patterns of morphological  
101 diversity in tenrecs. We use geometric morphometric techniques (Rohlf  
102 and Marcus, 1993) to compare cranial morphological diversity in tenrecs  
103 to that of their closest relatives, the golden moles (Afrosoricida,  
104 Chrysochloridae). We expect tenrecs to be more morphologically diverse  
105 than golden moles because tenrecs occupy a wider variety of ecological  
106 niches. The tenrec Family includes terrestrial, semi-fossorial, semi-aquatic  
107 and semi-arboreal species (Soarimalala and Goodman, 2011). In contrast,  
108 all golden moles occupy very similar, fossorial ecological niches (Bronner,  
109 1995). Greater ecological variety is often (though not always) correlated  
110 with higher morphological diversity (Losos and Mahler, 2010). However,  
111 our results reveal that, in skulls at least, morphological diversity in tenrecs  
112 is not as great as it first appears.

## 113 **Materials and Methods**

114 The methods we used involved several steps of i) data collection, ii)  
115 geometric morphometric analyses and iii) estimating morphological  
116 diversity. For clarity, Figure 1 summarises all of these steps and we  
117 describe them in detail below.

## 118 Data collection

119 One of us (SF) used the collections of five museums: the Natural History  
120 Museum, London (BMNH), the Smithsonian Institute Natural History  
121 Museum, Washington D.C. (SI), the American Museum of Natural  
122 History, New York (AMNH), the Museum of Comparative Zoology,  
123 Cambridge M.A. (MCZ) and the Field Museum of Natural History,  
124 Chicago (FMNH). We recorded species names as they were written on  
125 museum specimen labels and then corrected them to match the taxonomy  
126 in Wilson and Reeder's Mammal Species of the World (2005). For recently  
127 identified species, which are not included in Wilson and Reeder (2005), we  
128 used the taxonomy recorded on the specimen labels. Wilson and Reeder  
129 (2005) record 30 species of tenrec but more recent studies indicate that  
130 there are now 34 species (Olson, 2013). The additional species belong to  
131 the shrew tenrec (*Microgale*) Genus and represent either recognition of  
132 cryptic species boundaries (Olson et al., 2004) or discovery of new species  
133 (Goodman et al., 2006; Olson and Arroyo-Santos, 2009). Only one of these  
134 four recent additions, *M. jobihelyi*, was present in the museum collections  
135 and therefore we could not include the three other newly recognised  
136 species in the analyses. We photographed all of the tenrec and golden  
137 mole skulls available in the collections. This included 31 of the 34 species  
138 in the tenrec Family and 12 of the 21 species of golden moles (Wilson and  
139 Reeder, 2005).

140 We took pictures of the skulls using photographic copy stands  
141 consisting of a camera attachment with an adjustable height bar, a flat  
142 stage on which to place the specimen and an adjustable light source. To  
143 take possible light variability into account, on each day we took a

144 photograph of a white sheet of paper and used the custom white balance  
145 function on the camera to set the image as the baseline "white"  
146 measurement for those particular light conditions.

147 We photographed the specimens with a Canon EOS 650D camera fitted  
148 with a EF 100 mm f/2.8 Macro USM lens. We used a remote control  
149 (Hähnel Combi TF) to take the photos to avoid shaking the camera and  
150 distorting the images. We photographed the specimens on a black  
151 material background with a light source in the top left-hand corner of the  
152 photograph. We used small bean bags as necessary to hold the specimens  
153 in position while being photographed to ensure that they lay in a flat  
154 plane relative to the camera and did not tilt in any direction. We used the  
155 grid-line function on the live-view display screen of the camera to position  
156 the specimens in the centre of each image.

157 We photographed the skulls in three views: dorsal (top of the  
158 cranium), ventral (underside of the skull with the palate roof facing  
159 upwards) and lateral (right side of the skull) (Figure 1). When the right  
160 sides of the skulls were damaged or incomplete we photographed the left  
161 sides and later reflected the images so that they could be compared to  
162 pictures of the right sides (e.g. Barrow and Macleod, 2008).

163 We converted the raw files to binary (grey scale) images and re-saved  
164 them as TIFF files (uncompressed files preserve greater detail, RHOI,  
165 2013). Photographs of the specimens from the American Museum of  
166 Natural History and the Smithsonian Institute are available on figshare in  
167 separate file sets for the dorsal (Finlay and Cooper, 2013a), ventral (Finlay  
168 and Cooper, 2013c) and lateral (Finlay and Cooper, 2013b) skull pictures.  
169 Copyright restrictions from the other museums prevent public sharing of

170 their images but they are available on request.

## 171 **Geometric morphometric analyses**

172 We used a combination of landmark and semilandmark analysis  
173 approaches to assess the shape variability in the skulls. We used the TPS  
174 software suite (Rohlf, 2013) to digitise landmarks and curves on the  
175 photos. We set the scale on each image individually to standardise for the  
176 different camera heights used when photographing the specimens. We  
177 created separate data files for each of the three morphometric analyses  
178 (skulls in dorsal, ventral and lateral views). One of us (SF) digitised  
179 landmarks and semilandmark points on every image individually. Some  
180 specimens were too damaged to use in particular views so there were a  
181 different total number of images for each analysis. Our final data sets  
182 included photographs of 182 skulls in dorsal view (148 tenrecs and 34  
183 golden moles), 173 skulls in ventral view (141 tenrecs and 32 golden  
184 moles) and 171 skulls in lateral view (140 tenrecs and 31 golden moles).

185 When using semilandmark approaches there is a potential problem of  
186 over - sampling: simpler structures will require fewer semilandmarks to  
187 accurately represent their shape (MacLeod, 2012). To ensure that we  
188 applied a uniform standard of shape representation to each outline  
189 segment (i.e. that simple structures would not be over-represented and  
190 more complex features would not be under-represented), we followed the  
191 method outlined by MacLeod (2012). For each data set we chose a random  
192 selection of photos of specimens which represented the breadth of the  
193 morphological data (i.e. specimens from each sub-group of species). We  
194 drew the appropriate curves on each specimen and over-sampled the



195 number of points on the curves. We measured the length of the line and  
196 regarded that as the 100%, true length of that outline. We then re-sampled  
197 the curves with decreasing numbers of points and measured the length of  
198 the outlines. We calculated the length of each re-sampled curve as a  
199 percentage of the total length of the curve and then found the average  
200 percentage length for that reduced number of semilandmark points across  
201 all of the specimens in my test file. We continued this process until we  
202 found the minimum number of points that gave a curve length which was  
203 at least 95% accurate. We repeated these curve-sampling tests for each  
204 analysis to determine the minimum number of semilandmark points  
205 which would give accurate representations of morphological shape.

206 Figure 2 depicts that landmarks and curves which we used for each of  
207 the sets of photographs. For landmarks which are defined by dental  
208 structures, we used published dental sources (Repenning, 1967; Eisenberg  
209 and Gould, 1969; Nowak, 1983; MacPhee, 1987; Knox Jones and Manning,  
210 1992; Davis and Schmidly, 1997; Qu  rouil et al., 2001; Nagorsen, 2002;  
211 Wilson and Reeder, 2005; Goodman et al., 2006; Karata   et al., 2007;  
212 Hoffmann and Lunde, 2008; Asher and Lehmann, 2008; Muldoon et al.,  
213 2009; Lin and Motokawa, 2010) where available to identify the number  
214 and type of teeth in each species. Detailed descriptions of the landmarks  
215 can be found in the supplementary material.

216 After creating the files with the landmarks and semilandmarks placed  
217 on each photograph, we used TPSUtil (Rohlf, 2012) to create "sliders" files  
218 that defined which points in the TPS files should be treated as  
219 semilandmarks (Zelditch et al., 2012). We combined the landmarks and  
220 taxonomic identification files into a single morphometrics data object and  
221 carried out all further analyses in R version 3.1.1 (R Core Team, 2014).

222 Data and code for all of our analyses is available on GitHub (Finlay and  
223 Cooper, 2015).

224 At this stage, we either used the full data set (31 species of tenrec and  
225 12 species of golden mole) or a reduced data set with just 17 species of  
226 tenrec (Figure 1). We created this reduced data set because the majority of  
227 tenrec species (19 out of 31 in our data) belong to the *Microgale*  
228 (shrew-like) Genus that has relatively low morphological diversity  
229 (Soarimalala and Goodman, 2011; Jenkins, 2003). This may mask signals  
230 of higher morphological diversity among other tenrecs. To test this, we  
231 created a subset of the tenrec data that included just five of the *Microgale*  
232 species, each representing one of the five sub-divisions of *Microgale*  
233 outlined by Soarimalala and Goodman (2011), i.e. small, small-medium,  
234 medium, large and long-tailed species. We compared the morphological  
235 diversity of this subset of tenrecs (n=17: five *Microgale* and 12  
236 non-*Microgale* species) to that of the 12 species of golden moles (dashed  
237 arrows in Figure 1). After this selection stage, all further steps in the  
238 analyses were the same.

239 For each analysis, we used the `gpgen` function in the `geomorph`  
240 package (Adams et al., 2013) to run a general Procrustes alignment (Rohlf  
241 and Marcus, 1993) of the landmark coordinates while sliding the  
242 semilandmarks by minimising Procrustes distance (Bookstein, 1997). We  
243 used these Procrustes-aligned coordinates of all specimens to calculate  
244 average shape values for each species which we then used for a principal  
245 components (PC) analysis with the `plotTangentSpace` function (Adams  
246 et al., 2013). We selected the number of principal component (PC) axes  
247 that accounted for 95% of the variation in the data (Figure 1) and used  
248 these axes to estimate morphological diversity in each Family.

## Estimating morphological diversity

We grouped the PC scores for tenrecs and golden moles separately so that we could estimate the diversity of each Family and then compare the two groups (Figure 1). We compared morphological diversity in two ways.

First, we used non parametric multivariate analysis of variance (npMANOVA; Anderson, 2001) to test whether tenrecs and golden moles occupied significantly different positions within the morphospaces defined by the PC axes that accounted for 95% of the overall variation in the data (e.g. Serb et al., 2011; Ruta et al., 2013). A significant difference between the two Families would indicate that they have unique morphologies which do not overlap. Second, we compared morphological diversity within tenrecs to the diversity within golden moles. We define morphological diversity as the mean Euclidean distance (sum of squared differences) between each species and its Family centroid (Figure 3). This is summarised in the equation below where  $n$  is the number of species in the Family,  $i$  is the number of PC axes and  $c$  are the average PC scores for each axis (the centroid).

$$Disparity = \frac{\sqrt{\sum (PCn_i - PCc_i)^2}}{n} \quad (1)$$

If tenrecs are more morphologically diverse than golden moles, then they should be more dispersed within the morphospaces and have, on average, higher values of mean Euclidean distance.

One possible issue with these analyses is that the two Families have unequal sample sizes: 31 (or a subset of 17) tenrec species compared to just 12 golden mole species. Morphological diversity is usually decoupled

272 from taxonomic diversity (e.g. Ruta et al., 2013; Hopkins, 2013) so larger  
273 groups are not necessarily more morphologically diverse. However,  
274 comparing morphological diversity in tenrecs to the diversity of a smaller  
275 Family could still bias the results. We used pairwise permutation tests to  
276 account for this potential issue.

277 We tested the null hypothesis that tenrecs and golden moles have the  
278 same morphological diversity (the same mean Euclidean distance to the  
279 Family centroid). If this is true, when we randomly assign the group  
280 identity of each species (i.e. shuffle the "tenrec" and "golden mole" labels)  
281 and then re-compare the morphological diversity of the two groups, there  
282 would be no significant difference between these results and those  
283 obtained when the species are assigned to the correct groupings. We  
284 performed this shuffling procedure (random assignation of group  
285 identity) 1000 times and calculated the difference in morphological  
286 diversity between the two groups for each permutation. This generated a  
287 distribution of 1000 values which are calculations of the differences in  
288 morphological diversity under the assumption that the null hypothesis  
289 (equal morphological diversity in the two Families) is true. This method  
290 automatically accounts for differences in sample size because shuffling of  
291 the group labels preserves the sample size of each group: there will  
292 always be 12 species labelled as "golden mole" and then, depending on  
293 the analysis, either 31 or 17 species labelled as "tenrec". Therefore, the  
294 1000 permuted values of differences in morphological diversity create a  
295 distribution of the expected difference in diversity between a group of  
296 sample size 31 (or 17 in the case of the subsetted tenrec data) compared to  
297 a group of sample size 12 under the null hypothesis that the two groups  
298 have the same morphological diversity. We compared the observed

measures of the differences in morphological diversity between the two Families to these null distributions to determine whether there were significant differences after taking sample size into account (two-tailed t test).

## Results

Figure 4 depicts the morphospaces defined by the first two principal component (PC) axes from our principal components analyses (PCAs) of skull and mandible morphologies. The PCAs are based on the average Procrustes -superimposed shape coordinates for skulls in three views (dorsal, ventral and lateral). To compare morphological diversity in the two Families, we used the PC axes which accounted for 95% of the cumulative variation in each of the skull analyses: dorsal (n=6 axes), ventral (n=7 axes) and lateral (n=7 axes). First, we compared the position of each Family within the morphospace plots. Tenrecs and golden moles occupy significantly different positions in the dorsal (npMANOVA:  $F_{1,42}=68.13$ ,  $R^2=0.62$ ,  $p=0.001$ ), ventral (npMANOVA:  $F_{1,42}=103.33$ ,  $R^2=0.72$ ,  $p=0.001$ ) and lateral (npMANOVA:  $F_{1,42}=76.7$ ,  $R^2=0.65$ ,  $p=0.001$ ) skull morphospaces, indicating that the Families have very different, non-overlapping cranial and mandible morphologies (Figure 4).

Second, we compared the morphological diversity within each Family. Based on our measures of mean Euclidean distance to the Family centroids, tenrec skulls are more morphologically diverse than golden mole skulls when they are measured in lateral view but not in dorsal or ventral view (Table 1). In contrast, when we analysed morphological

323 diversity of skulls within the sub-sample of 17 tenrecs (including just five  
324 *Microgale* species) compared to the 12 golden mole species, we found that  
325 tenrec skulls were significantly more morphologically diverse than golden  
326 moles in all analyses (Table 1).

327 The pairwise permutation tests for each analysis confirmed that  
328 differences in morphological diversity were not artefacts of differences in  
329 sample size (Table 2)

## 330 Discussion

331 Tenrecs are often cited as an example of a mammalian group with high  
332 morphological diversity (Olson, 2013; Soarimalala and Goodman, 2011;  
333 Eisenberg and Gould, 1969). They are also more ecologically diverse than  
334 their closest relatives (Soarimalala and Goodman, 2011; Bronner, 1995) so  
335 we predicted that they would be more morphologically diverse than  
336 golden moles. However, our results do not support our original  
337 prediction, highlighting the importance of quantitative tests of perceived  
338 morphological patterns.

339 In our full analysis, tenrecs only had higher morphological diversity  
340 than golden moles when the skulls were measured in lateral view (Table  
341 1). There was no difference in morphological diversity when we analysed  
342 the skulls in dorsal or ventral views. This is most likely due to our choice  
343 of landmarks. The two outline curves in lateral view (Figure 2) emphasise  
344 morphological variation in the back and top of the skulls. These curves  
345 summarise overall shape variation but they do not identify clear  
346 anatomical differences because they are defined by relative features rather

347 than homologous structures (Zelditch et al., 2012). Therefore, high  
348 morphological diversity in tenrecs when analysed in this view may not  
349 indicate biologically or ecologically relevant variation. These lateral  
350 aspects of the skull morphology were not visible in the dorsal and ventral  
351 photographs so they could not be included in those analyses. In contrast,  
352 our landmarks in the dorsal, and particularly ventral, views focus on  
353 morphological variation in the overall outline shape of the sides of the  
354 skull and palate (Figure 2). The result that tenrecs are no more diverse  
355 than golden moles in these areas makes intuitive sense: most tenrecs have  
356 broad, non-specialised diets (Olson, 2013) so there is no obvious  
357 functional reason why they should have particularly diverse palate  
358 morphologies. The different results for our analysis of lateral skull  
359 morphologies compared to dorsal and ventral views highlight the  
360 importance of using multiple approaches when studying 3D  
361 morphological shape using 2D geometric morphometrics techniques  
362 (Arnqvist and Mårtensson, 1998). Landmark choice and placement will  
363 inevitably influence the results of a geometric morphometrics study. Our  
364 interest in broad-scale, cross-taxonomic comparisons of cranial  
365 morphology constrained our choice of landmarks to those that could be  
366 accurately identified in many different species (e.g. Ruta et al., 2013;  
367 Goswami et al., 2011; Wroe and Milne, 2007). In contrast, studies that use  
368 skulls to characterise morphological variation within species (e.g.  
369 Blagojević and Milošević-Zlatanović, 2011; Bornholdt et al., 2008) or to  
370 delineate species boundaries within a clade (e.g. Panchetti et al., 2008)  
371 tend to focus on more detailed, biologically homologous landmarks  
372 (Zelditch et al., 2012). Repeating our analyses with a narrower taxonomic  
373 focus may give greater insight into the specific morphological differences

374 among subgroups of tenrecs and golden moles.

375 In addition to the differences among the three skull views, our results  
376 indicate that, in skulls at least, the overall morphological diversity within  
377 tenrecs is not as large as is often assumed (e.g. Eisenberg and Gould, 1969;  
378 Olson, 2013). Studies of morphological variation are sensitive to the  
379 sampling used. If a particular morphotype is over-represented then the  
380 similarities among those species will reduce the overall morphological  
381 variation within the group (Foote, 1991). This appears to be the case for  
382 our data; it was only when we included a sub-sample of *Microgale* tenrecs  
383 that we found higher morphological diversity in tenrecs compared to  
384 golden moles across all three skull analyses (Table 1). While there are  
385 clear physical differences among Family members (Olson, 2013; Eisenberg  
386 and Gould, 1969), the majority of tenrecs are very morphologically similar  
387 (Jenkins, 2003) so morphological diversity in the Family as a whole is not  
388 as large as it first appears. The goal of our study was to quantify  
389 morphological variation in tenrecs instead of relying on subjective  
390 assessments of their high morphological diversity (Olson, 2013;  
391 Soarimalala and Goodman, 2011; Eisenberg and Gould, 1969). However, it  
392 is difficult to quantify overall morphological diversity because any study  
393 is inevitably constrained by its choice of specific traits (Roy and Foote,  
394 1997). Variation in skull shape is only one aspect of overall morphology.  
395 Quantifying variation in other morphological traits could yield different  
396 patterns. Therefore future work should extend our approach beyond  
397 skulls to gain a more complete understanding of the overall morphological  
398 diversity of tenrecs and golden moles. While recognising these limitations,  
399 our results provide valuable insights into the differences between  
400 subjective and quantitative assessments of morphological diversity.



## 401 **Conclusions**

402 We have presented the first quantitative investigation of morphological  
403 diversity in tenrecs. Our results indicate that, overall, tenrec skulls are not  
404 more morphologically diverse than golden moles and that similarities  
405 among the species rich *Microgale* tenrecs mask signals of higher  
406 morphological diversity among the rest of the Family. Of course the  
407 results presented here are restricted to just one axis of morphological  
408 variation and further analysis of other traits is required. However, our  
409 findings provide a foundation for future investigations and represent a  
410 significant step towards a more quantitative understanding of patterns of  
411 morphological and evolutionary diversity in tenrecs.

## 412 **Acknowledgements**

413 We thank Thomas Guillaume, François Gould and the members of NERD  
414 club for their insightful discussions and comments. Thank you to  
415 museum staff and curators for facilitating our access to their collections:  
416 Leona Leonard and Nigel Monaghan (Natural History Museum, Ireland),  
417 Roberto Portela Miguez and Paula Jenkins (Natural History Museum,  
418 London), Esther Langan (Smithsonian Institute), Eileen Westwig  
419 (American Museum of Natural History), Judy Chupasko (Museum of  
420 Comparative Zoology) and Bill Stanley and Steve Goodman (Field  
421 Museum).

## References

- Adams, D. C., E. Otárola-Castillo, and E. Paradis. 2013. geomorph: an R package for the collection and analysis of geometric morphometric shape data. *Methods in Ecology and Evolution* 4:393–399.
- Adams, D. C., F. J. Rohlf, and D. Slice. 2004. Geometric morphometrics: Ten years of progress following the “revolution”. *Italian Journal of Zoology* 71:5–16.
- Anderson, M. 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26:32–46.
- Arnqvist, G. and T. Mårtensson. 1998. Measurement error in geometric morphometrics; empirical strategies to assess and reduce its impact on measures of shape. *Acta Zoologica Academiae Scientiarum Hungaricae* 44:73–96.
- Asher, R. J. and T. Lehmann. 2008. Dental eruption in Afrotherian mammals. *BMC Biology* 6:14.
- Barrow, E. and N. Macleod. 2008. Shape variation in the mole dentary (Talpidae: Mammalia). *Zoological Journal of the Linnean Society* 153:187–211.
- Blagojević, M. and S. Milošević-Zlatanović. 2011. Sexual shape dimorphism in Serbian roe deer (*Capreolus capreolus* L.). *Mammalian Biology - Zeitschrift für Säugetierkunde* 76:735–740.
- Bookstein, F. 1997. Landmark methods for forms without landmarks: morphometrics of group differences in outline shape. *Medical Image Analysis* 1:225–243.

- 446 Bornholdt, R., L. R. Oliveira, and M. E. Fabián. 2008. Size and shape  
447 variability in the skull of *Myotis nigricans* (schinz, 1821) (chiroptera:  
448 Vespertilionidae) from two geographic areas in brazil. Brazilian Journal  
449 of Biology 68:623–631.
- 450 Bou, J., A. Casinos, and J. Ocaña. 1987. Allometry of the limb long bones  
451 of insectivores and rodents. Journal of Morphology 192:113–123.
- 452 Bronner, G. 1995. Systematic revision of the golden mole genera  
453 *Amblysomus*, *Chlorotalpa* and *Calcochloris* (Insectivora:  
454 Chrysochloromorpha; Chrysochloridae). Ph.D. thesis.
- 455 Brusatte, S., M. Benton, M. Ruta, and G. Lloyd. 2008. Superiority,  
456 competition and opportunism in the evolutionary radiation of  
457 dinosaurs. Science 321:1485–1488.
- 458 Davis, W. and D. Schmidly. 1997. The Mammals of Texas - Online Edition.  
459 <http://www.nsrl.ttu.edu/tmot1/Default.htm>.
- 460 Eisenberg, J. F. and E. Gould. 1969. The Tenrecs: A Study in Mammalian  
461 Behaviour and Evolution. Smithsonian Contributions to Zoology  
462 27:1–152.
- 463 Finlay, S. and N. Cooper. 2013a. "Insectivore" mammal skulls, dorsal view.  
464 <http://dx.doi.org/10.6084/m9.figshare.705863>.
- 465 Finlay, S. and N. Cooper. 2013b. "Insectivore" mammal skulls, lateral  
466 view. <http://dx.doi.org/10.6084/m9.figshare.715890>.
- 467 Finlay, S. and N. Cooper. 2013c. "Insectivore" mammal skulls, ventral  
468 view. <http://dx.doi.org/10.6084/m9.figshare.715841>.

- 469 Finlay, S. and N. Cooper. 2015. GitHub data and code repository.  
470 [https://github.com/SiveFinlay/Diversity\\_Paper](https://github.com/SiveFinlay/Diversity_Paper).
- 471 Foote, M. 1991. Morphological and taxonomic diversity in a clade's  
472 history: the blastoid record and stochastic simulations. University of  
473 Michigan, Museum of Paleontology Contributions 28:101–140.
- 474 Foth, C., S. Brusatte, and R. Butler. 2012. Do different disparity proxies  
475 converge on a common signal? Insights from the cranial morphometrics  
476 and evolutionary history of *Pterosauria* (Diapsida: Archosauria). Journal  
477 of Evolutionary Biology 25:904–915.
- 478 Garbutt, N. 1999. Mammals of Madagascar. Pica Press, Sussex.
- 479 Glor, R. 2010. Phylogenetic insights on adaptive radiation. Annual Review  
480 of Ecology, Evolution, and Systematics 41:251–270.
- 481 Goodman, S. M., C. J. Raxworthy, C. P. Maminirina, and L. E. Olson. 2006.  
482 A new species of shrew tenrec (*Microgale jobihely*) from northern  
483 Madagascar. Journal of Zoology 270:384–398.
- 484 Goswami, A., N. Milne, and S. Wroe. 2011. Biting through constraints:  
485 cranial morphology, disparity and convergence across living and fossil  
486 carnivorous mammals. Proceedings of the Royal Society B: Biological  
487 Sciences 278:1831–1839.
- 488 Hoffmann, R. and D. Lunde. 2008. Order Erinaceomorpha Pages 192–297.  
489 Princeton University Press, Oxfordshire, UK.
- 490 Hopkins, M. 2013. Decoupling of taxonomic diversity and morphological  
491 disparity during decline of the Cambrian trilobite family  
492 Pterocephaliidae. Journal of Evolutionary Biology 26:1665–1676.

- 493 Jenkins, P. 2003. *Microgale*, shrew tenrecs Pages 1273–1278. The University  
494 of Chicago Press, Chicago.
- 495 Karataş, A., M. Mouradi Gharkheloo, and T. Kankiliç. 2007. Cranial  
496 features and karyotypes of two hedgehogs (Insectivora: Erinaceidae)  
497 from Iran. *Anatomia, Histologia, Embryologia* 36:419–423.
- 498 Knox Jones, J. and R. Manning. 1992. *Insectivores* Page 75. Texas Tech  
499 University Press, United States of America.
- 500 Lin, L.-K. and M. Motokawa. 2010. *Mammals of Taiwan, Volume 1.*  
501 *Soricomorpha*. <http://mammal.biota.biodiv.tw/>.
- 502 Losos, J. 2011. Convergence, adaptation and constraint. *Evolution*  
503 65:1827–1840.
- 504 Losos, J. B. and D. Mahler. 2010. Adaptive radiation: the interaction of  
505 ecological opportunity, adaptation and speciation Pages 381–420.  
506 Sinauer Association, Sunderland, MA.
- 507 MacLeod, N. 2012. Going Round the Bend ii: Extended Eigenshape  
508 Analysis. <http://www.palass.org>.
- 509 MacPhee, R. 1987. The shrew tenrecs of Madagascar: Systematic revision  
510 and holocene distribution of *Microgale* (Tenrecidae, Insectivora).  
511 *American Museum Novitates* Number 2889:1–45.
- 512 Mahler, D., T. Ingram, L. Revell, and J. Losos. 2013. Exceptional  
513 convergence on the macroevolutionary landscape in island lizard  
514 radiations. *Science* 341:292–295.
- 515 Mitteroecker, P. and P. Gunz. 2009. Advances in geometric morphometrics.  
516 *Evolutionary Biology* 36:235–247.

- 517 Muldoon, K., D. de Blieux, E. Simons, and P. Chatrath. 2009. The  
518 subfossil occurrence and paleoecological significance of small mammals  
519 at Ankilitelo Cave, Southwestern Madagascar. *Journal of Mammalogy*  
520 90:111–1131.
- 521 Muschick, M., A. Indermaur, and W. Salzburger. 2012. Convergent  
522 evolution within an adaptive radiation of cichlid fishes. *Current Biology*  
523 22:1–7.
- 524 Nagorsen, D. 2002. An identification manual to the small mammals of  
525 British Columbia. Ministry of Sustainable Resource Management,  
526 Ministry of Water, Land and Air Protection, Biodiversity Branch and  
527 Royal BC Museum.
- 528 Nowak, R. 1983. Walker's Mammals of the World, 4th edition vol. 1. Johns  
529 Hopkins University Press, Baltimore.
- 530 Olson, L. E. 2013. Tenrecs. *Current Biology* 23:R5–R8.
- 531 Olson, L. E. and S. M. Goodman. 2003. Phylogeny and biogeography of  
532 tenrecs Pages 1235–1242. The University of Chicago Press, Chicago.
- 533 Olson, L. E., S. M. Goodman, and A. D. Yoder. 2004. Illumination of  
534 cryptic species boundaries in long-tailed shrew tenrecs (Mammalia:  
535 Tenrecidae; *Microgale*), with new insights into geographic variation and  
536 distributional constraints. *Biological Journal of the Linnean Society*  
537 83:1–22.
- 538 Olson, M. E. and A. Arroyo-Santos. 2009. Thinking in continua: beyond  
539 the "adaptive radiation" metaphor. *BioEssays* 31:1337–1346.

540 Panchetti, F., M. Scalici, G. Carpaneto, and G. Gibertini. 2008. Shape and  
 541 size variations in the cranium of elephant-shrews: a morphometric  
 542 contribution to a phylogenetic debate. *Zoomorphology* 127:69–82.

543 Price, S., S. S. B. Hopkins, K. K. Smith, and L. Roth. 2012. Tempo of  
 544 trophic evolution and its impact on mammalian diversification.  
 545 *Proceedings of the National Academy of Sciences, USA* 109:7008–7012.

546 Quéroutil, S., P. Hutterer, M. Colyn, J. Kerbis Peterhans, and E. Verheyen.  
 547 2001. Phylogeny and evolution of African shrews (Mammalia:  
 548 Soricidae) inferred from 16s rRNA sequences. *Molecular Phylogenetics*  
 549 *and Evolution* 20:185–195.

550 R Core Team. 2014. R: A language and environment for statistical  
 551 computing. <http://www.R-project.org/>.

552 Repenning, C. 1967. Subfamilies and Genera of the Soricidae. Geological  
 553 Survey Professional Paper 565 United States Government Printing  
 554 Office, Washington.

555 RHOI. 2013. Revealing Hominid Origins Initiative Fossil Photography  
 556 Protocol, U.C Berkeley. [http:](http://rhoi.berkeley.edu/RHOI_photo/RHOI_Photography_Protocol.html)  
 557 [//rhoi.berkeley.edu/RHOI\\_photo/RHOI\\_Photography\\_Protocol.html](http://rhoi.berkeley.edu/RHOI_photo/RHOI_Photography_Protocol.html).

558 Rohlf, F. 2012. Tpsutil ver 1.53. <http://life.bio.sunysb.edu/morph/>.

559 Rohlf, F. 2013. Tpsdig2 ver 2.17. <http://life.bio.sunysb.edu/morph/>.

560 Rohlf, J. and L. Marcus. 1993. A revolution in morphometrics. *Trends in*  
 561 *Ecology and Evolution* 8:129–132.

562 Roy, K. and M. Foote. 1997. Morphological approaches to measuring  
 563 biodiversity. *Trends in Ecology and Evolution* 12:277–281.

- 564 Ruta, M., K. Angielczyk, J. Fröbisch, and M. Benton. 2013. Decoupling of  
565 morphological disparity and taxic diversity during the adaptive  
566 radiation of anomodont therapsids. *Proceedings of the Royal Society B:*  
567 *Biological Sciences* 280:20131071.
- 568 Salton, J. A. and E. Sargis. 2009. Evolutionary morphology of the  
569 Tenrecoidea (Mammalia) hindlimb skeleton. *Journal of Morphology*  
570 270:367–387.
- 571 Serb, J., A. Alejandrino, E. Otárola-Castillo, and D. Adams. 2011.  
572 Morphological convergence of shell shape in distantly related scallop  
573 species (Mollusca: Pectinidae). *Zoological Journal of the Linnean*  
574 *Society* 163:571–584.
- 575 Slice, D. 2007. Geometric morphometrics. *Annual Review of*  
576 *Anthropology* 36:261–281.
- 577 Soarimalala, V. and S. Goodman. 2011. *Les petits mammifères de*  
578 *Madagascar. Guides sur la diversité biologique de Madagascar*  
579 *Association Vahatra, Antananarivo, Madagascar.*
- 580 Stanhope, M., V. Waddell, O. Madsen, W. de Jong, S. Hedges, G. Cleven,  
581 D. Kao, and M. Springer. 1998. Molecular evidence for multiple origins  
582 of insectivora and for a new order of endemic african insectivore  
583 mammals. *Proceedings of the National Academy of Sciences, USA*  
584 95:9967–9972.
- 585 Wilson, D. and D. Reeder. 2005. *Mammal species of the world. A*  
586 *taxonomic and geographic reference (3rd edition). Johns Hopkins*  
587 *University Press.*



- 588 Wroe, S. and N. Milne. 2007. Convergence and remarkably consistent  
589 constraint in the evolution of carnivore skull shape. *Evolution*  
590 61:1251–1260.
- 591 Zelditch, M., D. Swiderski, and D. Sheets. 2012. *Geometric Morphometrics*  
592 *for Biologists*, 2nd edition. Academic Press, Elsevier.

## 593 **List of Figures**

594	1	Flowchart diagram of data collection and analysis . . . . .	27
595	2	Skulls: dorsal, ventral and lateral landmarks . . . . .	28
596	3	Calculating diversity as mean Euclidean distance to Family	
597		centroid. . . . .	29
598	4	Morphospace (principal components) plot of morphological	
599		diversity in tenrec and golden mole skulls. . . . .	30




Figure 1: Summary of the main steps in our data collection, processing and analysis protocol. Note that the analyses were repeated separately for each set of photographs: skulls in dorsal, ventral and lateral views. The dashed arrows refer to the stage at which we selected a subsample of the tenrecs (including just five species of the *Microgale* Genus) so that we could compare the morphological diversity of this reduced subsample of tenrec species to the diversity of golden moles.




Figure 2: Landmarks (numbered points) and curves (outlines) for the skulls in dorsal,ventral and lateral view. See the supplementary material for detailed landmark descriptions. The skulls are two different specimens of *Potamogale velox* (otter shrew tenrec), museum accession numbers AMNH 51327 (dorsal picture) and BMNH 1934.6.16.2 (ventral and lateral pictures).




Figure 3: Estimating morphological diversity as the mean Euclidean distance between each species and the Family centroid. Every species had scores on the principal components (PC) axes that accounted for 95% of the variation in the principal components analysis. The number of axes (PCn) varied for each analysis but they were the same within a single analysis. PC scores were used to calculate the Euclidean distance from each species to the Family centroid (average ( $\bar{x}$ ) PC scores for the entire Family). Morphological diversity of the Family is the average value of these Euclidean distances.




Figure 4: Principal components plots of the morphospaces occupied by tenrecs (triangles,  $n=31$  species) and golden moles (circles,  $n=12$  species) for the skulls in dorsal (top left), ventral (top right) and lateral (bottom left) views. Each point represents the average skull shape of an individual species. Axes are principal component 1 (PC1) and principal component 2 (PC2) of the average scores from principal components analyses of mean Procrustes shape coordinates for each species.

## 600 **List of Tables**

<small>601</small>	<b>1</b>	Comparing morphological diversity in tenrecs and golden	
<small>602</small>		moles. . . . .	<b>32</b>
<small>603</small>	<b>2</b>	Results of the permutation tests . . . . .	<b>33</b>

Table 1: Morphological diversity in tenrecs compared to golden moles (12 species). N is the number of tenrec species: 31 species or 17 species including just five representatives of the *Microgale* Genus. Morphological diversity of the Family is the mean Euclidean distance from each species to the Family centroid. Significant differences between the two Families ( $p < 0.05$ ) from two-tailed t-tests are highlighted in bold.

N	Analysis	Morphological diversity		$t_{df}$	p value
		Tenrecs (mean $\pm$ s.e)	Golden moles (mean $\pm$ s.e)		
31	Skulls dorsal	0.036 $\pm$ 0.0029	0.029 $\pm$ 0.0032	-1.63 <sub>29.88</sub>	0.11
	Skulls ventral	0.048 $\pm$ 0.0034	0.044 $\pm$ 0.0041	-0.68 <sub>26.99</sub>	0.51
	Skulls lateral	0.044 $\pm$ 0.0041	0.032 $\pm$ 0.0037	-2.16 <sub>35.03</sub>	<b>0.04</b>
17	Skulls dorsal	0.044 $\pm$ 0.0025	0.029 $\pm$ 0.0032	-3.62 <sub>22.75</sub>	<b>&lt;0.01</b>
	Skulls ventral	0.054 $\pm$ 0.0039	0.042 $\pm$ 0.0041	-2.23 <sub>25.46</sub>	<b>0.04</b>
	Skulls lateral	0.054 $\pm$ 0.0053	0.031 $\pm$ 0.0037	-3.47 <sub>26.31</sub>	<b>&lt;0.01</b>



Table 2: Results of the permutation analyses comparing the observed differences in morphological diversity to a null distribution of expected results. Morphological diversity of the Family is the mean Euclidean distance from each species to the Family centroid. Results are shown for both the full (N=31 species of tenrec compared to 12 species of golden mole) and reduced (N=17 species of tenrec compared to 12 golden moles) data sets. Significant values ( $p < 0.05$ ) indicate that the observed morphological diversity is different to the expected differences under a null hypothesis of equivalent diversities in the two Families.

N	Analysis	Morphological diversity					p value
		Measured values			Permuted values		
		Tenrecs	Golden moles	Difference	Min.	Max.	
31	Dorsal	0.036	0.029	0.007	-0.011	0.009	0.013
	Ventral	0.048	0.044	0.004	-0.014	0.013	0.023
	Lateral	0.044	0.032	0.012	-0.012	0.011	0.001
17	Dorsal	0.044	0.029	0.015	-0.011	0.014	0.001
	Ventral	0.054	0.042	0.013	-0.017	0.019	0.023
	Lateral	0.054	0.031	0.022	-0.018	0.019	0.001