

HOD informed prior of PNG bias

Siyi Zhao

Contents

1	Introduction	1
2	Mocks and their Power Spectra	1
2.1	HOD model	1
2.1.1	velocity bias	1
2.1.2	Secondary properties: concentration	1
2.1.3	Secondary properties: environment	1
2.2	HOD fittings	1
2.3	lightcone mocks based on HOD	1
2.4	Measuring the power spectrum from the mocks	1
3	Inference	2
3.1	The PNG Model in desilike	2

1 Introduction

2 Mocks and their Power Spectra

2.1 HOD model

2.1.1 velocity bias

$$v_{\text{cent, z}} = v_{\text{L2, z}} + \alpha_c \delta v (\sigma_{\text{LoS}}) \quad (1)$$

2.1.2 Secondary properties: concentration

2.1.3 Secondary properties: environment

2.2 HOD fittings

2.3 lightcone mocks based on HOD

2.4 Measuring the power spectrum from the mocks

Following ?, we estimate the power spectrum multipoles using the Yamamoto estimator. The multipoles of the power spectrum are estimated as

$$\hat{P}_\ell(k_\mu) = \frac{2\ell+1}{AV_{k_\mu}} \int_{V_{k_\mu}} d\mathbf{k} \int d\mathbf{x}_1 \int d\mathbf{x}_2 e^{i\mathbf{k} \cdot (\mathbf{x}_2 - \mathbf{x}_1)} \mathcal{F}(\mathbf{x}_1) \mathcal{F}(\mathbf{x}_2) \mathcal{L}_\ell(\hat{\mathbf{k}} \cdot \hat{\mathbf{x}}_1) - \mathcal{N}_\ell, \quad (2)$$

where $\mathcal{F}(\mathbf{x})$ is the FKP weight field estimated from the data and the random catalogues. The normalization factor A is given by

$$A = \int d\mathbf{x} \bar{n}_g^2(\mathbf{x}), \quad (3)$$

Table 1: Priors used for LRG and QSO HOD fits. We use broad Gaussian priors on all parameters. We also quote the bounds we impose in addition to the Gaussian priors. Units of mass are in $h^{-1}M_{\odot}$.

	LRG (w/o dv)		QSO (w/ dv)	
Params	Prior	Bounds	Prior	Bounds
$\log M_{\text{cut}}$	flat	[11, 15]	$\mathcal{N}(12.7, 1.0)$	[11.2, 14.0]
$\log M_1$	flat	[10, 18]	$\mathcal{N}(15.0, 1.0)$	[12.0, 16.0]
σ	log	[0.0001, 1]	$\mathcal{N}(0.5, 0.5)$	[0.0, 3.0]
α	flat	[-1, 3]	$\mathcal{N}(1.0, 0.5)$	[0.3, 2.0]
κ	flat	[0, 6]	$\mathcal{N}(0.5, 0.5)$	[0.3, 3.0]
α_c	flat	[0, 3]	$\mathcal{N}(1.5, 1.0)$	[0.0, 2.0]
α_s	flat	[0, 3]	$\mathcal{N}(0.2, 1.0)$	[0.0, 2.0]

and the shot noise term is removed with

$$\mathcal{N}_{\ell} = \frac{\delta_{\ell 0}}{A} \int d\mathbf{x} \bar{n}_g(\mathbf{x}) w^2(\mathbf{x}) \mathcal{L}_{\ell}(\hat{\mathbf{k}} \cdot \hat{\mathbf{x}}). \quad (4)$$

By introducing

$$F_{\ell}(\mathbf{k}) = \int d\mathbf{x} e^{i\mathbf{k} \cdot \mathbf{x}} \mathcal{F}(\mathbf{x}) \mathcal{L}_{\ell}(\hat{\mathbf{k}} \cdot \hat{\mathbf{x}}), \quad (5)$$

Eq. 2 reduces to

$$\hat{P}_{\ell}(k_{\mu}) = \frac{2\ell + 1}{AV_{k_{\mu}}} \int_{V_{k_{\mu}}} d\mathbf{k} F_0(\mathbf{k}) F_{\ell}(-\mathbf{k}) - \mathcal{N}_{\ell}. \quad (6)$$

3 Inference

3.1 The PNG Model in desilike

We perform the inference with desilike.

The model is as follows, the power spectrum of biased tracers is

$$P(k, z) = \left(b_1(z) + f_{\text{NL}}^{\text{loc}} \frac{b_{\Phi}(z)}{T_{\Phi \rightarrow \delta}(k, z)} \right)^2 P_{\text{lin}}(k, z), \quad (7)$$

where b_1 is the linear bias, b_{Φ} is the PNG bias, P_{lin} is the linear matter power spectrum, and $T_{\Phi \rightarrow \delta}$ is the transfer function from the primordial gravitational potential field Φ to matter overdensity δ and can be computed from CLASS by

$$T_{\Phi \rightarrow \delta}(k, z) \equiv \sqrt{\frac{P_{\text{lin}}(k, z)}{P_{\Phi}(k)}} \quad \text{with} \quad P_{\Phi}(k) = \frac{9}{25} \frac{2\pi^2}{k^3} A_s \left(\frac{k}{k_{\text{pivot}}} \right)^{n_s - 1}, \quad (8)$$

where P_{Φ} is the primordial potential (normalised to $3/5\mathcal{R}$ to match the usual definition [TBD]) power spectrum, n_s is the spectral index and A_s is the amplitude of initial power spectrum at $k_{\text{pivot}} = 0.05 \text{ Mpc}^{-1}$ [ToBeCheck].

[ToBeCheck] There are also an usually used equation show the relation between $T_{\Phi \rightarrow \delta}(k, z)$ and the matter transfer function $T_m(k, z)$,

$$T_{\Phi \rightarrow \delta}(k, z) = \frac{2}{3\Omega_m} \frac{k^2}{H_0^2} (1+z) T_m(k, z). \quad (9)$$

The full model should include RSD effect, we adopt a simpler model including the Kaiser effect and a damping factor for small scales as

$$P(k, \mu) = \frac{\left(b_1(z_{\text{eff}}) + f_{\text{NL}}^{\text{loc}} \frac{b_{\Phi}(z_{\text{eff}})}{T_{\Phi \rightarrow \delta}(k, z_{\text{eff}})} + f(z_{\text{eff}}) \mu^2 \right)^2}{\left[1 + \frac{1}{2} (k\mu\Sigma_s)^2 \right]^2} P_{\text{lin}}(k, z_{\text{eff}}) + s_{\text{n},0}. \quad (10)$$

This is the model used in desilike, an example of the usage can be found in https://github.com/cosmodesi/desilike/blob/hmc/nb/png_examples.ipynb f is the linear growth rate. Σ_s is the amount of damping at small scales. $s_{n,0}$ is a potential residual shot noise which should be close to 0 as we always remove the shot noise contribution from our power spectrum measurements.

Finally, the power spectrum is expanded in Legendre multipoles as

$$P_\ell(k) = \frac{2\ell+1}{2} \int_{-1}^1 d\mu P(k, \mu) \mathcal{L}_\ell(\mu). \quad (11)$$

We consider the monopole ($\ell = 0$) and quadrupole ($\ell = 2$) in our analysis. Since the statistical error on the hexadecapole ($\ell = 4$) is too big in large scales, we do not include it in the inference.

For the PNG bias b_Φ , a usual relation is

$$b_\Phi(z) = 2\delta_c(b_1(z) - p), \quad (12)$$

where $\delta_c = 1.686$ is the critical overdensity for spherical collapse, and p quantifies the merger history of the tracer. We study the redshift evolution of p in [TBD].

References