

---

# Climate Change and Its Impact on Thermal Ecology and Biodiversity: A Survey

---

[www.surveyx.cn](http://www.surveyx.cn)

## Abstract

This survey paper explores the complex interplay between climate change, thermal ecology, and biodiversity, emphasizing their interconnected impacts on ecological systems. It highlights the critical role of temperature dynamics, such as daily minimum temperatures and fluctuations, in shaping ecosystem health and species interactions. Advanced modeling techniques, including Bayesian hierarchical models, are instrumental in predicting climate impacts and providing early warning signals for ecological transitions. Thermal ecology significantly influences species interactions, with temperature fluctuations affecting metabolic rates and population dynamics, particularly in insect populations. Biodiversity acts as a buffer against environmental changes, essential for maintaining planetary climate stability. The paper underscores the potential of crop diversification as an adaptation strategy and the promise of Quantum Artificial Intelligence in advancing climate science. Despite challenges, no evidence was found for increasing precipitation extremes in dry regions, highlighting the need for refined data analysis. Future research should expand time series data, refine analytical methods, and integrate interdisciplinary perspectives to address climate impacts on ecosystems. Innovative solutions, such as bioacoustic communication in pest management, are crucial for enhancing biodiversity resilience. Overall, the survey calls for continued research and innovation to mitigate climate impacts and ensure ecosystem resilience, advocating for integrating socioeconomic considerations into risk assessment and management strategies.

## 1 Introduction

### 1.1 Climate Change Overview

Climate change is a multifaceted global challenge, necessitating interdisciplinary approaches to address its complex impacts [1]. Driven primarily by anthropogenic activities such as fossil fuel combustion and deforestation, rising atmospheric CO<sub>2</sub> levels significantly enhance the greenhouse effect, resulting in global warming. This warming correlates with increased frequency and intensity of extreme weather events, including severe heat waves and floods, which jeopardize Earth's habitability [2].

The variability in atmospheric conditions, especially in mid-latitude regions, complicates climate dynamics and challenges the accuracy of climate models [3]. These models are essential for simulating interactions among climate system components and for understanding changes in precipitation patterns, particularly in arid regions where statistical artifacts may have previously led to overestimated trends.

The global implications of climate change extend to terrestrial and freshwater ecosystems, with observed and projected shifts threatening biodiversity and ecosystem services. Additionally, climate change intensifies the challenges posed by invasive species, highlighting the need for sustainable alternatives to traditional control methods, such as chemical pesticides [4]. Research into climate




Figure 1: chapter structure

system behaviors, including ice age dynamics and temperature fluctuations, is vital for developing sustainable strategies that address the economic, environmental, and social complexities of climate change, thus emphasizing the necessity for global cooperation and policy interventions.

## 1.2 Relevance to Thermal Ecology and Biodiversity

Climate change significantly influences thermal ecology and biodiversity by altering temperature regimes and atmospheric conditions, which in turn affect species interactions, distributions, and survival strategies. This is particularly evident in regions like the Mediterranean Basin, where climate change threatens water resources, biodiversity, and food security [5]. Understanding these interactions is crucial for predicting ecosystem responses and developing effective adaptation strategies [6]. The global redistribution of species due to changing climate conditions disrupts ecosystems and human societies, underscoring the importance of studying these dynamics within thermal ecology and biodiversity frameworks [7].

Shifts in population boundaries driven by environmental gradients are critical for forecasting biodiversity changes [8]. The feedback mechanisms linking the biosphere and planetary climate systems emphasize the intricate relationship between climate stability and biodiversity [9]. This interconnectedness is vital for maintaining ecological balance and habitability, both of which are threatened by ongoing climatic shifts [10].

Global warming contributes to the expansion and weakening of the tropical atmospheric component, disrupting temperature patterns and atmospheric circulation, which may alter ecological networks and species interactions [11]. Rising temperatures and increased atmospheric CO<sub>2</sub> levels, along with altered precipitation patterns, influence agricultural pest dynamics, necessitating adaptive pest management strategies [8]. These changes have cascading effects on ecosystem health and biodiversity, impacting food security and ecosystem services [5].

Furthermore, climate change affects soil microbiomes, essential for biogeochemical cycling and overall ecosystem health [12]. The limitations of current numerical models in accurately predicting climate change due to sensitivity to initial conditions and potential tipping points pose additional challenges [13]. Discrepancies between current global warming trends and future projections from general circulation models raise questions about their reliability [14].

---

Habitat fragmentation, exacerbated by climate change, threatens adaptive genetic variation and alters mating systems, complicating ecosystem resilience [10]. The inadequate understanding of global warming, stemming from biases in geographical representation and temporal limitations in existing studies, underscores the need for standardized methods to analyze temperature variations [13]. Additionally, the potential influence of cosmic rays on climate forcing, as suggested by correlations between cosmic ray variations and climate changes documented in cosmogenic isotope archives, adds complexity to climate dynamics [11].

The survival of cold-adapted species, such as *Nebria germari* in high-altitude habitats, is jeopardized by climate change-induced alterations in their habitats and phenology [10]. Moreover, a long-term decrease in coloration of Mediterranean blue tit subspecies over the past 15 years illustrates the impact of climate on biodiversity, emphasizing the need for ongoing monitoring and research [7]. Addressing these multifaceted challenges requires an interdisciplinary approach to understand the intricate interactions among climate change, thermal ecology, and biodiversity.

### 1.3 Structure of the Survey

This survey is organized into several key sections, each addressing distinct aspects of climate change and its complex relationship with thermal ecology and biodiversity. The introductory section provides an overview of climate change, emphasizing its global significance and relevance to thermal ecology and biodiversity, while also outlining the survey's structure.

The second section delves into the background and definitions of core concepts such as daily minimum temperature, temperature fluctuation, interspecific interaction, thermal ecology, insect metabolism, and biodiversity, establishing a foundational understanding for subsequent topics.

Section three examines climate change and temperature dynamics, detailing how climate change affects temperature patterns, including daily minimum temperatures and fluctuations, and discussing the implications for ecosystems and species interactions. Subsections within this section explore modeling temperature fluctuations, the impact of temperature extremes on ecosystem health, and predictive climate modeling techniques.

The fourth section investigates interspecific interactions and thermal ecology, analyzing how temperature fluctuations influence species interactions within ecosystems, highlighting thermal ecology's role in understanding these dynamics and the challenges posed by climate change. Methodological approaches to studying interspecific interactions are also reviewed.

In section five, the survey explores the effects of climate change on insect metabolism, considering how temperature changes can alter metabolic rates and impact insect populations, with broader implications for ecosystem functioning and biodiversity, including population dynamics and adaptation strategies.

The sixth section examines biodiversity and ecosystem resilience, analyzing the impact of climate change on biodiversity and the resilience of ecosystems to temperature changes and interspecific interactions. It explores strategies for conserving biodiversity and mitigating climate change impacts, discussing the synergistic effects of climate change on biodiversity and ecosystem stability.

The conclusion synthesizes the principal findings of the survey, emphasizing the intricate relationships between climate change, thermal ecology, and biodiversity. It underscores the urgent need for ongoing research to address these interconnections and identifies specific areas for future investigation, such as the impacts of climate variability on agricultural sustainability, biodiversity loss, and public health challenges related to climate-induced disease patterns [1, 15, 16]. The following sections are organized as shown in Figure 1.

## 2 Background and Definitions

### 2.1 Key Concepts and Their Interconnections

Climate change, marked by persistent shifts in global temperature and weather patterns, poses significant challenges to thermal ecology and biodiversity due to its intricate and dynamic nature. The chaotic dynamics of climate, coupled with inherent natural variability, complicate the identification of statistically significant trends, with traditional extreme value analysis often failing to account

---

for nonstationary behavior in temperature distributions [17]. Daily minimum temperature, a crucial climate indicator, reflects climate variability and influences physiological processes and ecosystem dynamics. These fluctuations, exacerbated by anthropogenic activities like increased CO<sub>2</sub> emissions, highlight the need for precise climate sensitivity models, which remain challenging to estimate due to varying feedback mechanisms across different climate states [3].

Interspecific interactions, involving relationships between different species, are profoundly influenced by temperature dynamics. Environmental gradients and local heterogeneities shape population boundaries, illustrating the complexity of ecological systems under changing climatic conditions [4]. Traditional models often inadequately capture nonlinear interactions within these complex systems, lacking the capability to account for intricate feedback mechanisms [1]. Thermal ecology, which examines the effects of temperature on organisms and their environments, provides a framework for understanding these interactions and their broader ecological implications. Habitat fragmentation, especially in forest ecosystems, can hinder species' evolutionary responses to environmental changes, affecting their climate adaptation.

Insect metabolism, encompassing the biochemical processes that sustain insect life, is sensitive to temperature fluctuations, altering metabolic rates and impacting population dynamics and ecosystem functioning [3]. The dynamics of insect infestations, particularly in boreal forests, demonstrate how climate influences forest health, with extreme weather events, changing precipitation patterns, and rising temperatures posing threats to food security and agricultural practices.

Biodiversity, the variety of life forms within an ecosystem, is integral to ecosystem resilience and stability. The interplay between climate change and biodiversity is mediated through feedback mechanisms linking the biosphere and planetary climate systems [4]. Accurately representing the nonlinear characteristics of climate models is essential for understanding climate sensitivity and its effects on biodiversity. The entropy budget of the climate system, focusing on material processes, is crucial for assessing climate stability and the role of biodiversity in maintaining it [2]. A limited understanding of how microorganisms influence and are influenced by climate change may jeopardize efforts toward an environmentally sustainable future.

Recent theoretical advancements, such as integrating climate network and percolation theories, provide insights into the evolution of climate patterns and their responses to global warming. Approaches proposed by Huang et al. yield reliable estimates of rare temperature extremes using extensive climate model runs, overcoming the limitations of short observational records [18]. Theoretical tools combining statistical physics and applied mathematics enhance the understanding of climate dynamics, addressing the challenge of distinguishing climate variability from climate change signals amidst unsteady external forcings and complex internal dynamics. The theory suggesting that climatic variations are significantly influenced by solar and planetary oscillations, often inadequately represented in current General Circulation Models (GCMs), further complicates the understanding of climate dynamics.

The inadequacy of current metrics for measuring temporal turnover, failing to capture the unidirectional nature of community dynamics and the ecological processes driving changes in community composition over time, presents a challenge. Existing metrics, such as the Jaccard and Bray-Curtis indices, were designed for spatial contexts and do not adequately account for the unique characteristics of temporal turnover [18]. The benchmark established for accurately estimating daily global CO<sub>2</sub> emissions from fossil fuel combustion and cement production during 2020 underscores the need for real-time data to reduce uncertainties in global emissions assessments [2]. Many existing benchmarks rely on outdated national inventories, failing to capture real-time changes in emissions, leading to significant uncertainties. Urban cooling demand trends, categorized by climate drivers and urban characteristics, emphasize the role of extreme heat events and urban heat islands in shaping energy demands. The proposition that climate sensitivity increases with warming challenges previous assumptions of constant sensitivity across different climate states [19]. Effectively teaching complex climate concepts, such as the greenhouse effect and planetary albedo, through interactive tools is crucial for enhancing understanding. Extreme temperatures adversely affect agricultural productivity, leading to reduced crop yields and altered farming practices among subsistence farmers. Organizing current methods into stages of energy resource utilization, policy development, and technological innovation, while focusing on their interdependencies and impacts on climate change, is critical for addressing these challenges. The necessity to identify and confirm the existence of a 60-year oscillation in global climate data, inadequately represented in current GCMs, further complicates the

understanding of climate dynamics [3]. These concepts and their interconnections underscore the complexity of climate change and its pervasive impact on thermal ecology and biodiversity.

In recent years, the significance of advanced modeling techniques in the study of climate change has gained considerable attention. These models not only facilitate a deeper understanding of temperature dynamics but also play a crucial role in predicting future climate scenarios. To illustrate this complexity, Figure 2 presents a hierarchical classification of these advanced modeling techniques and their applications. The figure categorizes the approaches into three primary areas: modeling temperature fluctuations, assessing the impacts of temperature extremes on ecosystem health, and employing predictive climate modeling techniques. Each category highlights key methodologies and underscores their respective roles in enhancing climate predictions, thereby providing a comprehensive framework for researchers and policymakers alike.




Figure 2: This figure illustrates the hierarchical classification of advanced modeling techniques and their applications in understanding climate change and temperature dynamics. It categorizes the approaches into modeling temperature fluctuations, impacts of temperature extremes on ecosystem health, and predictive climate modeling techniques, highlighting key methodologies and their roles in enhancing climate predictions.

### 3 Climate Change and Temperature Dynamics

#### 3.1 Modeling Temperature Fluctuations

Understanding temperature fluctuations is crucial for assessing climate change impacts on ecological and atmospheric systems. Traditional deterministic models often fall short in capturing the stochastic and nonlinear nature of temperature dynamics, prompting the development of advanced methodologies. The Bayesian hierarchical model (HCM-RBE) effectively addresses spatial and temporal dependencies in record-breaking temperature events, offering insights into regional trends, such as those in peninsular Spain [20]. The multi-level spatio-temporal model (MLSTM) improves predictive accuracy by incorporating autoregressive terms and spatial processes, emphasizing the critical need to consider both spatial and temporal dimensions in climate models [21].

As illustrated in Figure 3, advanced modeling techniques for understanding temperature fluctuations and climate dynamics can be categorized into three main approaches: Bayesian hierarchical models, machine learning techniques, and flexible parametric models. This figure highlights key methodologies such as HCM-RBE, the DCR method utilizing eXtreme Gradient Boosting (XGBoost) models, and the BATs distribution model. The DCR method enhances predictions of historical daily CO<sub>2</sub> emissions, enriching our understanding of climate dynamics [22]. These methodologies demonstrate the integration of advanced data analytics in climate modeling, improving predictions and interpretations of climatic impacts across sectors, including agriculture [23]. The BATs distribution model, a flexible parametric tool, accommodates both extreme and bulk temperature data, providing a comprehensive framework for analyzing temperature variability [17]. Hierarchical frameworks like WOMBAT v2.0, employing spatially varying time-series decomposition, underscore the importance of spatial and temporal variability in climate modeling [24].

These advanced modeling techniques are essential for understanding temperature fluctuations, highlighting their role in shaping climate change impacts and guiding effective adaptation strategies. By enhancing the precision of climate predictions through sophisticated statistical methodologies, these approaches provide valuable insights into the ecological and atmospheric consequences of temperature variations, thereby enriching discussions on climate change across various platforms, including social media [1, 15, 25].




Figure 3: This figure illustrates advanced modeling techniques for understanding temperature fluctuations and climate dynamics, categorized into Bayesian hierarchical models, machine learning techniques, and flexible parametric models, highlighting key methodologies such as HCM-RBE, DCR Method, and BATs Model.

#### 3.2 Temperature Extremes and Ecosystem Health

Temperature extremes, marked by significant deviations from average climatic conditions, pose substantial challenges to ecosystem health and stability. These extremes can disrupt species distributions and community dynamics, vital for maintaining ecological balance [28]. Advanced statistical models are crucial for understanding temperature trends and their ecological ramifications. The TML model captures the non-normality of temperature residuals and regime-switching behavior, representing a significant advancement in modeling temperature dynamics [29]. Table 1 provides a comprehensive summary of various modeling techniques employed to analyze temperature extremes and their impact on ecosystem resilience and climate-ecosystem interactions.

Method Name	Modeling Techniques	Ecosystem Resilience	Climate-Ecosystem Interaction
NSM[26]	Neural Shooting Method	-	Climate Change Dynamics
MLSTM[21]	Hierarchical Bayesian Methods	Maintaining Stability	Complex Spatial Dependencies
ACF[27]	Analog Curve Fitting	-	Climate Change Impacts
CMLA[8]	Double Machine Learning	Adaptation Strategy	Climatic Stressors Impacts

Table 1: Overview of modeling techniques and their applications in understanding ecosystem resilience and climate-ecosystem interactions. This table summarizes the methods used, including Neural Shooting, Hierarchical Bayesian, Analog Curve Fitting, and Double Machine Learning, highlighting their contributions to climate change dynamics, maintaining stability, adaptation strategies, and understanding complex spatial dependencies.

The synchronization of climatic oscillations with astronomical cycles highlights the limitations of current General Circulation Models (GCMs) in replicating these patterns, which are crucial for predicting extreme temperature events [30]. The neural shooting approach, combined with the Onsager-Machlup action functional, enhances predictive capabilities by identifying probable transition pathways in the carbon cycle [26].

In arid ecosystems, the self-organization and clustering of vegetation patterns are significantly influenced by environmental inhomogeneities exacerbated by temperature extremes [14]. This process is vital for understanding ecosystem resilience under climate-induced stress. The negative feedback effect of low clouds on climate sensitivity underscores the complexity of climatic responses to temperature extremes [31].

The rapid decrease in zonal surface temperature gradients due to warming, as explained by a simplified surface energy budget, illustrates the intricate interactions between temperature extremes and climate dynamics [32]. Integrating spatially varying coefficients and temporal autoregressive effects in modeling daily maximum temperatures provides a comprehensive framework for analyzing temperature trends and their impacts [21].

Systemic growth constants identified in CO<sub>2</sub> emission trajectories suggest significant temperature increases with profound implications for ecosystem health [27]. These projections highlight the urgency of developing effective adaptation strategies to enhance ecosystem resilience. Notably, crop diversification has been shown to increase net primary productivity by 2.8%, demonstrating a viable strategy for mitigating the impacts of temperature extremes on agricultural systems [8].

Explicitly considering temperature's effect on shear strength in slope stability analyses represents a significant advancement over previous methods [2]. This approach is essential for understanding the broader ecological implications of temperature extremes. The complex interplay between temperature extremes and ecosystem stability necessitates a multidisciplinary approach that integrates atmospheric, oceanic, and biological data. Advanced modeling techniques and theoretical frameworks are critical for accurately predicting the impacts of extreme temperatures, guiding effective adaptation strategies, and ensuring ecosystem resilience in the face of increasing climate variability.

### 3.3 Predictive Climate Modeling Techniques

Predictive climate modeling techniques are essential for accurately forecasting the impacts of climate change on temperature dynamics, crucial for developing effective adaptation strategies. Traditional General Circulation Models (GCMs) have struggled to fully capture the inherent uncertainties in climate systems, leading to the integration of probabilistic modeling approaches that better reflect these uncertainties [33]. The incorporation of stochastic parameterizations represents a significant advancement, enabling models to account for the effects of unresolved scales on climate dynamics, thereby enhancing prediction accuracy [21].

The integration of machine learning and statistical methods into climate modeling has demonstrated promise in improving predictive capabilities. Bayesian hierarchical models, incorporating spatial correlation and inter-model dependence, yield more precise uncertainty quantifications and enhanced accuracy in climate projections [33]. This approach emphasizes the importance of considering both spatial variability and temporal dependencies for a comprehensive understanding of temperature dynamics.

Innovative techniques, particularly Quantum Artificial Intelligence (QAI), have the potential to significantly enhance modeling capabilities in climate science by leveraging advanced computational

---

methods that efficiently process vast datasets. As climate change continues to pose severe challenges, including extreme weather events, integrating QAI into numerical modeling could lead to improved spatial and temporal forecasting, facilitating more effective adaptation strategies and contributing to societal benefits through better predictions and mitigative measures [34, 35]. These advancements underscore the evolving landscape of climate modeling, where the fusion of traditional and cutting-edge methodologies provides a robust framework for understanding climate dynamics.

The development of analytical methods that replicate the main periods of Milankovitch cycles offers reliable alternatives to traditional numerical integration methods, highlighting the need for a diverse array of tools in predictive modeling. The integration of data assimilation techniques and advanced numerical simulations to assess the climate system exemplifies ongoing efforts to enhance the precision and reliability of climate models. These methodologies, informed by empirical decomposition of climatic data and statistical analyses, enable researchers to better understand the interactions between natural climate cycles and anthropogenic influences, refining predictions of future climate scenarios. Notably, the work of Klaus Hasselmann has paved the way for innovative approaches in model response analysis and detection of climate change signals, further advancing climate science [3, 25, 36].

Recent advancements include visual analysis and prediction methods utilizing classic models adapted to data characteristics, simplifying complexity and reducing the need for advanced feature engineering. Spatial quantile clustering methods effectively address the distributional characteristics of climate data, including asymmetry and outliers, leading to more reliable clustering outcomes [21].

These predictive climate modeling techniques, integrating stochastic processes, machine learning, and innovative statistical methods, are crucial for advancing our understanding of climate dynamics. As climate models evolve, their enhanced ability to accurately predict temperature dynamics will be vital for developing effective strategies to mitigate climate change impacts, safeguarding ecosystem resilience and biodiversity. Utilizing biophysical models that capture the intricate thermodynamic interactions between organisms and their environments enables better forecasting of species' responses to climate fluctuations. This understanding is essential for managing risks such as biodiversity loss, shifts in ecosystem structures, and changes in predator-prey dynamics, all exacerbated by climate variability. Furthermore, accurate temperature forecasting can inform integrated resource management approaches, ultimately supporting the sustainability of ecosystems and the services they provide to human well-being [16, 37, 38, 39, 7].

## **4 Interspecific Interactions and Thermal Ecology**

### **4.1 Role of Thermal Ecology in Species Interactions**

Thermal ecology plays a pivotal role in understanding species interactions and ecosystem stability amid climate change, which alters temperature regimes and ecological boundaries. The adaptive capacity of species to these changes is crucial for survival, and a framework for analyzing climate connectivity provides insights into emerging patterns and potential tipping points, essential for understanding the interplay between climate variables and species dynamics [40].

Temperature fluctuations also impact human health, notably through cardiovascular mortality, underscoring the broader implications on ecological and human systems [5]. This necessitates frameworks that integrate ecological and anthropogenic factors for comprehensive climate change impact assessments. Microhabitat characteristics, such as supraglacial debris, serve as refugia for cold-adapted species, maintaining ecological stability under climatic shifts [11].

Incorporating interspecies interactions into predictive models enhances species distribution predictions, offering insights into temperature's influence on species dynamics and ecosystem resilience [41]. Frameworks considering community dynamics and temporal turnover, distinct from spatial turnover, are vital for capturing ecological interaction nuances [42]. Thermal ecology provides a critical lens for examining species-environment interactions, yielding insights into ecosystem resilience, vital for developing adaptive strategies that enhance stability and biodiversity conservation [3].



## 4.2 Methodological Approaches to Studying Interspecific Interactions

Robust methodologies are essential for understanding interspecific interactions within thermal ecology. Reaction-diffusion models, incorporating logistic growth and Allee effects, are foundational for describing population dynamics across space and time, crucial for understanding species responses to climate change [43]. Integrating these models with empirical data allows simulation of species distribution dynamics, accounting for shifting climate envelopes, species mobility, and extinction risks during migration [44, 45, 46, 38].

As illustrated in Figure 4, which depicts the primary methodological approaches to studying interspecific interactions in thermal ecology, these methodologies encompass modeling techniques, data integration, and conservation strategies. Advanced statistical and computational techniques, including agent-based models and network analysis, have gained prominence in studying interspecific interactions. Agent-based models simulate individual behaviors and interactions, providing insights into species responses to temperature changes at the micro-level, while network analysis reveals temperature as a critical driver of interaction networks [47, 48, 40]. Machine learning algorithms analyze extensive ecological datasets, uncovering intricate relationships and emerging trends in thermal ecology, enhancing understanding of climate change impacts on interspecific dynamics [48, 42, 37].

Remote sensing technologies provide valuable spatial and temporal data on habitat conditions and species distributions, facilitating comprehensive assessments of temperature variations' impact on interspecific interactions. Integrating diverse methodologies, including ecological assessments, climate modeling, and socio-economic analyses, aids in developing effective conservation strategies and policy frameworks addressing biodiversity loss and ecosystem service disruptions [49, 16, 7].




Figure 4: This figure illustrates the primary methodological approaches to studying interspecific interactions in thermal ecology, highlighting modeling techniques, data integration, and conservation strategies.

## 4.3 Climate Change and Interspecific Interaction Challenges

Climate change significantly challenges interspecific interactions, impacting biodiversity and ecosystem dynamics. Biodiversity reduction threatens planetary climate systems' stability, leading to ecological imbalances and altered species interactions [50]. Predicting species redistribution outcomes is complex due to varying responses to climate change, resulting in unpredictable ecological consequences [49].

Inadequate historical datasets and logistical sampling challenges, particularly in inaccessible regions, impede accurate assessments of interspecific interactions [51]. This data gap limits understanding of climate change impacts, while existing datasets' low temporal resolution restricts models' ability to capture dynamic ecosystem changes, such as mangrove distribution [52]. Biases in climate models further complicate understanding interspecific interactions under changing conditions [53].

Addressing these challenges requires multifaceted approaches, including nonparametric methods for clearer insights into extreme temperature trends [54]. Significant reductions in fossil fuel emissions and effective CO<sub>2</sub> extraction strategies are crucial to mitigate future catastrophic climate impacts [55]. Enhancing data collection, improving model accuracy, and fostering interdisciplinary research

---

are vital for devising strategies to mitigate climate change impacts on interspecific interactions and ensure ecosystem resilience.

Fragmentation of climate strategies diminishes effectiveness and collaboration among stakeholders [56]. Limited use of biophysical models in ecology, coupled with data availability issues, restricts comprehensive solution development [37]. However, sustainable bio-control methods, such as enhancing predator efficacy, present promising avenues for addressing these challenges [4]. Infrastructure, traditionally seen as detrimental, may aid population persistence under climate change by supporting ecological adaptation [45]. Intermediate mutation and adaptation levels can enhance survival, providing insights into evolutionary strategies buffering against climate-induced extinction risks [57]. Recognizing ecological grief is crucial for understanding the emotional dimensions of climate change impacts, vital for fostering resilience and adaptation [18].

## 5 Insect Metabolism and Climate Change

### 5.1 Temperature Fluctuations and Metabolic Rates

Temperature fluctuations critically affect insect metabolic rates, pivotal for understanding population dynamics and ecological processes. As ectotherms, insects rely on external temperatures for metabolic regulation, where variations impact growth, reproduction, and survival, subsequently influencing ecological dynamics [5]. Accurate climate models are essential for predicting these impacts. Advanced modeling techniques, like machine learning, enhance climate model precision by integrating observed data with predictions, improving our grasp of climate change on insect populations [58]. Solar models further elucidate the relationship between solar activity and climate change, affecting temperature dynamics and insect metabolism [59].

Rainfall patterns, closely linked to temperature changes, shape insect population dynamics, notably impacting species such as *Aedes aegypti*, with significant ecosystem and public health implications [4]. This interconnectedness underscores the collective influence of climatic factors on insect metabolism and ecological outcomes. Historical climate data indicate a 60-year oscillation, likely due to solar-astronomical factors, contributing to global warming and affecting insect metabolic rates [60]. These insights are vital for understanding long-term impacts on insect metabolism and ecosystem stability. Methodologies estimating changes in linear restoring rates based on Langevin-type dynamics offer valuable tools for detecting critical ecological transitions [61].

### 5.2 Population Dynamics and Adaptation

Insect populations adapt diversely to climate-induced metabolic changes, crucial for ecological balance and biodiversity. Their adaptive capacity is closely tied to climate change rates, with thresholds beyond which extinction risks rise if adaptation fails. Understanding interactions between climate dynamics and insect metabolism is vital for predicting population shifts and addressing issues like deforestation and pest management, threatening food security and ecological stability [62, 63]. Advanced methodologies, such as the GKLT algorithm, enhance extreme heatwave season predictions, providing insights into climatic event impacts on insect populations [64]. Wavelet spectral analysis (WSA) aids in identifying shifts in extreme events, offering perspectives on insect responses to climate dynamics [65].

To better illustrate these complex relationships, Figure 5 presents a hierarchical categorization of key concepts related to insect population dynamics and adaptation in the context of climate change. This figure highlights the impact of climate on insect metabolism, species interactions, and adaptation strategies in agriculture, underscoring the multifaceted nature of these interactions.

Species interactions, like predator-prey dynamics, are significantly affected by climate variability, where shifts between favorable and unfavorable conditions alter these interactions, influencing population stability [38]. Understanding these dynamics is crucial for predicting broader ecological implications. Habitat fragmentation challenges insect adaptation by disrupting climate adaptation processes and reducing drought tolerance, as seen in *Primula elatior* [66]. This leads to decreased flowering investment and altered mating systems, impacting biodiversity and ecosystem resilience. Research often faces limitations due to sampling biases and body size distribution variability, obscuring true ecological trends [10].

Non-autonomous nonlinear models, as proposed for honey bee-parasite interactions, highlight seasonality's role in reproductive strategies and population dynamics [67]. These models provide insights into insect life cycle adaptations to environmental changes. The interaction between selection pressure and mutation rates is critical for predicting insect responses to climate change [57]. Understanding these interactions informs conservation and biodiversity management strategies. In agriculture, extreme heat impacts productivity, prompting subsistence farmers to adapt by altering crop types, partially offsetting declines [68]. Future research should refine temperature dynamics models and explore applications across climatic regions or other weather variables affecting agriculture [29]. These efforts are crucial for developing comprehensive adaptation strategies ensuring ecosystem and agricultural resilience.




Figure 5: This figure illustrates the hierarchical categorization of key concepts related to insect population dynamics and adaptation in the context of climate change. It highlights the impact of climate on insect metabolism, species interactions, and adaptation strategies in agriculture.

### 5.3 Case Studies: Specific Insect Populations

Climate change profoundly impacts insect populations, affecting distribution, abundance, and ecological roles. Case studies on specific populations reveal insights into climate change's multifaceted impacts, illustrating how rising temperatures, altered precipitation, and increased CO<sub>2</sub> levels influence insect dynamics. These studies show geographic range expansions of pests, life cycle changes, and invasion risks, threatening agricultural productivity and food security. They highlight the need for adaptive management and innovative pest control measures to address these climatic challenges [69, 67, 63, 62, 15].

The honey bee (*Apis mellifera*) exemplifies climate change impacts through seasonality and parasitism. Research shows seasonality can enhance survival or cause collapse, depending on environmental conditions and parasitic threats [67]. This underscores the intricate balance between climatic variables and biological interactions, necessitating adaptive management for pollinator survival. Leaf litter decomposition, mediated by insect detritivores, is another climate-influenced area. Future research should refine models by incorporating factors like nitrogen content and site fertility, affecting decomposition rates [70]. Understanding micro-scale processes in broader climatic shifts is vital for nutrient cycling and ecosystem health.

Annual plant-insect interactions further illustrate climate change effects. Studies suggest annual plants thrive in hotter, drier conditions due to climate variability and anthropogenic influences [71]. This shift alters associated insect population composition and behavior, impacting ecosystem dynamics and biodiversity. These case studies emphasize ongoing research into climate change's multifaceted impacts on insects. Analyzing specific climate-driven changes provides insights into ecosystem dynamics, such as biodiversity loss, shifts in structure, and altered hydrological regimes. This understanding enables targeted conservation strategies to mitigate adverse effects on ecosystem services, enhancing resilience to climate impacts [1, 15, 7].

---

## 6 Biodiversity and Ecosystem Resilience

### 6.1 Impacts on Ecosystem Functioning

Climate change significantly alters ecosystem functioning by influencing both biotic and abiotic components through shifts in temperature, precipitation, and extreme weather events. The state-dependent nature of climate sensitivity results in diverse ecosystem responses across climatic regimes, necessitating a nuanced understanding for predicting adaptations and recoveries from climatic perturbations [19]. Advanced forecasting models indicate that while immediate action may seem less urgent, it remains critical to prevent surpassing vital temperature thresholds [72].

Biodiversity is crucial for ecosystem resilience, providing functional traits that buffer against climate variability and sustain ecosystem functionality. Small changes in carbon levels can lead to significant shifts in climate dynamics, emphasizing the interconnectedness of climate change and biodiversity [19]. Mountain ecosystems, for instance, display varied responses to climate change, with some regions experiencing increased vegetation cover and microbial biomass, while others face declines in freshwater quality. These differential responses underscore the complexity of ecosystem dynamics under climate stress, highlighting the importance of localized studies to identify specific vulnerabilities. The continuous spatiotemporal approach enhances accuracy by incorporating spatial heterogeneity and managing missing data, improving estimations of climate change impacts compared to traditional methods [2].

High-altitude ice-related landforms act as refugia for cold-adapted species, mitigating extinction risks associated with climate warming. The geometry of environmental boundaries and Allee effects critically influence population persistence, revealing that high mobility does not guarantee survival amid rapid climate change. This necessitates sophisticated modeling approaches that integrate spatial and temporal dynamics to predict ecosystem responses accurately, particularly given the complex interactions among species and environmental changes [73, 37, 42, 12, 46].

Ecosystem resilience to climate change is intricately linked to biodiversity, which enhances adaptation and recovery from disturbances. Preserving biodiversity is vital for maintaining essential ecosystem services, such as food security, water quality, and carbon sequestration, while bolstering ecological resilience against rapid climatic shifts. As climate change alters ecological community composition, it affects both ecosystem functioning and human well-being, underscoring the need to integrate biodiversity considerations into adaptation and mitigation strategies for sustainable development [49, 7].

### 6.2 Conservation and Mitigation Strategies

Conservation and mitigation strategies are essential for addressing the multifaceted impacts of climate change on biodiversity and ecosystems. These strategies require an integrated approach combining biological, environmental, and socio-economic factors to manage the intricate interactions within the climate system. The Optimal Enhancing Resilience Model (OERM) has been proposed as a strategic framework for safeguarding critical ecological components, emphasizing the maintenance of ecosystem resilience amid climatic perturbations [12]. Moreover, precision agriculture and the development of stress-tolerant crop varieties are vital for enhancing agricultural resilience to climate variability and sustaining ecosystem services.

Figure 6 illustrates the hierarchical structure of conservation and mitigation strategies, highlighting key categories such as enhancing resilience, advanced climate modeling, and public engagement. Each category encompasses specific strategies and innovations aimed at addressing climate change impacts on biodiversity and ecosystems. Incorporating real-time data streams into climate models can significantly improve their adaptability and accuracy, facilitating better decision-making in policy and management [33]. The potential of Quantum Artificial Intelligence (QAI) to enhance climate models and forecasts emphasizes the need for further exploration in machine learning methods [34]. Public education and awareness are critical for fostering support for conservation and mitigation efforts, with enhanced carbon pricing mechanisms driving substantial reductions in greenhouse gas emissions [74].

Integrating biophysical models can deepen understanding of species responses to climate change, highlighting the necessity for improved training and data infrastructure to broaden their application [6]. Efficient analysis of large datasets can bolster strategies for conserving biodiversity in urban

environments. Future research should focus on refining existing frameworks, exploring data-driven approaches, and addressing current models' limitations in capturing the multiscale nature of climate dynamics [75]. By combining advanced modeling techniques, conservation practices, and public engagement, these strategies can effectively mitigate climate change impacts, ensuring biodiversity preservation and ecosystem resilience.

Future research could explore sophisticated methods to combine ecological data across species and investigate subsets that respond differently to climate models [62]. Additionally, refining models to incorporate additional oscillations and integrating detailed empirical data can enhance predictive accuracy [3]. Increasing the density of weather stations in underrepresented areas and integrating remote sensing data can improve evapotranspiration estimation, crucial for understanding ecosystem responses to climate change [76]. Comprehensive multi-attribution analyses are necessary for a complete understanding of climate dynamics. The ecological role of cold-adapted species and their interactions with competitors and prey should also be considered in conservation strategies [77].

Innovative pest management approaches align with conservation strategies by reducing reliance on harmful pesticides and supporting biodiversity [4]. Future research should focus on in-depth analyses of specific climate change topics and emerging trends in adaptation and mitigation strategies [1]. Furthermore, developing fully-coupled thermo-hydro-mechanical models and conducting case studies to validate proposed methods under various conditions is essential for advancing conservation efforts [2].




Figure 6: This figure illustrates the hierarchical structure of conservation and mitigation strategies, highlighting key categories such as enhancing resilience, advanced climate modeling, and public engagement. Each category includes specific strategies and innovations aimed at addressing climate change impacts on biodiversity and ecosystems.

### 6.3 Synergistic Effects of Climate Change on Biodiversity

The synergistic effects of climate change on biodiversity and ecosystem stability are characterized by complex interactions between climatic variables and ecological processes, potentially leading to abrupt shifts in population boundaries. Such shifts often occur when gradual environmental changes mask ecological responses until significant changes are triggered, which can lead to misinterpretations of stability [78]. These dynamics underscore the potential for sudden and unpredictable changes in biodiversity as ecosystems respond to climatic pressures.

Wavelet Spectral Analysis (WSA) has proven effective in detecting significant shifts in climate signals, crucial for understanding how these changes impact biodiversity and ecosystem stability [65]. These shifts can disrupt ecological networks, leading to cascading effects on species interactions and ecosystem functions. However, gaps remain in understanding the long-term effects of climate change on pest dynamics, complicating management strategies and emphasizing the need for comprehensive approaches to mitigate these impacts [63].

Current climate models face limitations in accurately predicting future temperatures, particularly due to the complex role of solar influences, which are not fully understood [79]. The potential role of cosmic rays in modulating climate further complicates these predictions, suggesting that additional research is needed to elucidate these influences [80]. These uncertainties highlight the importance

---

of refining climate models to incorporate a broader range of climatic factors and improve regional process representations [81].

Technological uncertainties, high costs, and the lack of comprehensive policy frameworks present significant challenges to the large-scale implementation of climate mitigation strategies [82]. Addressing these challenges is critical for developing effective conservation strategies that enhance biodiversity and ecosystem resilience. Future research should focus on integrating additional ecological variables and conducting field studies to validate theoretical predictions, thereby improving understanding of the synergistic effects of climate change [67].

Quantum Mechanical principles offer a promising avenue for gaining deeper insights into climate change dynamics, potentially leading to a better understanding of climatic transitions and their impacts on biodiversity [83]. The gravitational influences of celestial bodies, as described by Milankovitch cycles, further complicate long-term climate dynamics, affecting biodiversity and ecosystem stability [84]. Addressing these complexities will enable researchers to develop more accurate models and strategies to mitigate the synergistic effects of climate change on biodiversity, ultimately ensuring ecosystem resilience amid ongoing climatic shifts.

#### **6.4 Role of Biodiversity in Climate System Stability**

Biodiversity is vital for sustaining climate system stability and ecosystem health, providing a diverse array of functional traits that enhance resilience to environmental changes. The ongoing redistribution of species due to climate change alters ecological community composition, affecting ecosystem functioning, food security, and carbon sequestration processes. This dynamic underscores the critical need to integrate biodiversity considerations into climate adaptation and mitigation strategies, as biodiversity loss can undermine ecosystem services essential for human well-being [85, 49, 16, 7]. The interplay between biodiversity and climate dynamics is underscored by the capacity of diverse ecosystems to enhance resilience, thereby sustaining functionality under varying climatic conditions. A wide range of species within an ecosystem contributes to its ability to adapt and recover from disturbances, ensuring ecological balance and the continued provision of ecosystem services.

The significance of biodiversity in climate system stability is further illustrated by feedback mechanisms linking biological diversity with climate processes. For example, diverse plant communities can influence atmospheric carbon levels through varying rates of photosynthesis and respiration, thereby affecting climate regulation. This interaction emphasizes biodiversity's essential role in influencing climate dynamics and alleviating climate change effects, as species redistribution driven by climate change alters critical ecosystem functions for food security, disease transmission, and carbon sequestration, impacting human well-being and the overall climate system [16, 6, 15, 49, 7].

Existing research has established connections between solar activity and climate, enhancing understanding of historical climate patterns and the role of astronomical factors [79]. These insights into the natural variability of the climate system underscore biodiversity's importance in buffering against both natural and anthropogenic climatic changes. Biodiverse ecosystems are better equipped to absorb shocks and maintain stability amid external pressures, contributing to the overall resilience of the climate system.

Moreover, biodiversity supports ecosystem health by facilitating essential processes such as nutrient cycling, soil formation, and pollination. These processes are crucial for maintaining ecosystem integrity and their capacity to support life. The decline in biodiversity, driven largely by human-induced climate change, can disrupt these essential ecological processes, diminishing ecosystem resilience and increasing susceptibility to climate impacts. This loss not only alters species distributions and community compositions but also adversely affects critical ecosystem services such as food production, disease regulation, and carbon sequestration. Consequently, the ongoing biodiversity crisis poses serious risks to ecosystem integrity and human well-being, necessitating urgent integration of biodiversity considerations into climate adaptation and mitigation strategies [49, 7].

## **7 Conclusion**

This survey provides a comprehensive exploration of the intricate relationships between climate change, thermal ecology, and biodiversity, highlighting their critical interdependencies and diverse ecological impacts. The dynamics of temperature, especially the fluctuations in daily minimum

---

temperatures, play a crucial role in determining ecosystem health and the nature of species interactions. The application of sophisticated modeling techniques, such as Bayesian hierarchical models and the MC-LR method, enhances our ability to attribute climate changes and predict complex ecological processes. These models are instrumental in identifying critical slowing down (CSD) and issuing early warnings for potential critical transitions, particularly in environments characterized by red noise.

The role of thermal ecology in mediating species interactions is underscored by the influence of temperature variations on metabolic rates and population dynamics, especially within insect communities. This necessitates ongoing research into adaptive strategies and the refinement of models that incorporate complex ecological interactions and potential tipping points. Biodiversity serves as a vital buffer against environmental fluctuations, contributing significantly to the stability of the planetary climate system. The survey also emphasizes the potential of crop diversification as an adaptation strategy to enhance agricultural resilience, particularly in vulnerable regions like Cyprus.

The potential of Quantum Artificial Intelligence to advance climate science is notable, offering improved pattern recognition and simulation capabilities, despite existing technological hurdles. The survey identifies a lack of evidence for increasing precipitation extremes in arid regions when employing precise definitions and robust data processing methods, underscoring the importance of advancing data analysis techniques for accurate climate assessments.

Future research directions should focus on extending time series datasets, refining analytical approaches, and examining the impacts of emerging climate patterns on global food security. Interdisciplinary research that integrates ecological, evolutionary, and neurobiological insights is crucial for understanding the broader implications of climate change, particularly its effects on nervous systems. Additionally, investigating bioacoustic communication in insect infestation dynamics presents new avenues for detection and management strategies. The integration of additional food resources with spatial dynamics offers promising avenues for effective pest management, highlighting the need for innovative solutions in addressing the challenges posed by climate change and biodiversity conservation.

---

## References

- [1] Robin Haunschild, Lutz Bornmann, and Werner Marx. Climate change research in view of bibliometrics, 2016.
- [2] Marco Loche and Gianvito Scaringi. Temperature continuously controls the stability of clay slopes, 2022.
- [3] Craig Loehle and Nicola Scafetta. Climate change attribution using empirical decomposition of climatic data, 2012.
- [4] Urvashi Verma, Aniket Banerjee, and Rana D. Parshad. T(w)o patch or not t(w)o patch: A novel additional food model, 2024.
- [5] Pierre Masselot, Fateh Chebana, Taha B. M. J. Ouarda, Diane Bélanger, André St-Hilaire, and Pierre Gosselin. A new look at weather-related health impacts through functional regression, 2018.
- [6] Arianna Pera and Luca Maria Aiello. Shifting climates: Climate change communication from youtube to tiktok, 2024.
- [7] Camille Parmesan, Mike D Morecroft, and Yongyut Trisurat. *Climate change 2022: Impacts, adaptation and vulnerability*. PhD thesis, GIEC, 2022.
- [8] Georgios Giannarakis, Ilias Tsoumas, Stelios Neophytides, Christiana Papoutsas, Charalampos Kontoes, and Diofantos Hadjimitsis. Understanding the impacts of crop diversification in the context of climate change: a machine learning approach, 2023.
- [9] Adam Frank and Woodruff Sullivan. Sustainability and the astrobiological perspective: Framing human futures in a planetary context, 2013.
- [10] Grant M. Connette, John A. Crawford, and William E. Peterman. Climate change and shrinking salamanders: Alternative mechanisms for changes in plethodontid salamander body size, 2014.
- [11] Keno Riechers, Georg Gottwald, and Niklas Boers. Glacial abrupt climate change as a multi-scale phenomenon resulting from monostable excitable dynamics, 2024.
- [12] Yu Sun, Teng Liu, Shang Wang, Jun Meng, Yongwen Zhang, Saini Yang, Xiaosong Chen, Deliang Chen, Jürgen Kurths, Shlomo Havlin, Hans Joachim Schellnhuber, and Jingfang Fan. Regional greening as a ‘positive’ tipping phenomenon, 2023.
- [13] Flavio C. D. Moraes, Ana Lia Leonel, Pedro H. C. Torres, Pedro R. Jacobi, and Sandra Momm. Climate change and social sciences: a bibliometric analysis, 2020.
- [14] D. Pinto-Ramos, M. G. Clerc, A. Makhoute, and M. Tlidi. Vegetation clustering and self-organization in inhomogeneous environments, 2024.
- [15] Smriti Janaswamy and Jeremy Blackburn. Exploring climate change discourse: Measurements and analysis of reddit data, 2024.
- [16] Kashif Abbass, Muhammad Zeeshan Qasim, Huaming Song, Muntasir Murshed, Haider Mahmood, and Ijaz Younis. A review of the global climate change impacts, adaptation, and sustainable mitigation measures. *Environmental science and pollution research*, 29(28):42539–42559, 2022.
- [17] Mitchell Krock, Julie Bessac, Michael L. Stein, and Adam H. Monahan. Nonstationary seasonal model for daily mean temperature distribution bridging bulk and tails, 2021.
- [18] Ashlee Cunsolo and Neville R Ellis. Ecological grief as a mental health response to climate change-related loss. *Nature Climate Change*, 8(4):275–281, 2018.
- [19] Lucas Benney and Anca Radulescu. An energy balance model of carbon’s effect on climate change, 2015.
- [20] Jorge Castillo-Mateo, Alan E. Gelfand, Zeus Gracia-Tabuenca, Jesús Asín, and Ana C. Cebrián. Spatio-temporal modeling for record-breaking temperature events in Spain, 2024.



- 
- [21] Jorge Castillo-Mateo, Miguel Lafuente, Jesús Asín, Ana C. Cebrián, Alan E. Gelfand, and Jesús Abaurrea. Spatial modeling of day-within-year temperature time series: an examination of daily maximum temperatures in aragón, spain, 2022.
- [22] Tao Li, Lixing Wang, Zihan Qiu, Philippe Ciais, Taochun Sun, Matthew W. Jones, Robbie M. Andrew, Glen P. Peters, Piyu ke, Xiaoting Huang, Robert B. Jackson, and Zhu Liu. Reconstructing global daily co2 emissions via machine learning, 2024.
- [23] Giuliano Vitali, Sergei Rogosin, and Guido Baldoni. Climate change and grain production fluctuations, 2019.
- [24] Michael Bertolacci, Andrew Zammit-Mangion, Andrew Schuh, Beata Bukosa, Jenny Fisher, Yi Cao, Aleya Kaushik, and Noel Cressie. Inferring changes to the global carbon cycle with wombat v2.0, a hierarchical flux-inversion framework, 2022.
- [25] Bora Akgun, Zeynep Isvan, Levent Tuter, and Mehmet Levent Kurnaz. Climate prediction through statistical methods, 2008.
- [26] Jianyu Chen, Jianyu Hu, Wei Wei, and Jinqiao Duan. A data-driven approach for discovering the most probable transition pathway for a stochastic carbon cycle system, 2022.
- [27] Jessie Lydia Henshaw. The systemic growth constants of climate change: From its origin in 1780 to its major post-wwii acceleration, 2019.
- [28] Marta Magnani, Rubén Díaz-Sierra, Luke Sweeney, Antonello Provenzale, and Mara Baudena. Fire responses shape plant communities in a minimal model for fire ecosystems across the world, 2023.
- [29] Samuel Asante Gyamerah, Philip Ngare, and Dennis Ikpe. Regime-switching temperature dynamics model for weather derivatives, 2018.
- [30] Nicola Scafetta. Solar and planetary oscillation control on climate change: hind-cast, forecast and a comparison with the cmip5 gcms, 2013.
- [31] Jyrki Kauppinen and Pekka Malmi. No experimental evidence for the significant anthropogenic climate change, 2019.
- [32] Timothy M. Merlis and Tapio Schneider. Changes in zonal surface temperature gradients and walker circulations in a wide range of climates, 2011.
- [33] Huang Huang, Dorit Hammerling, Bo Li, and Richard Smith. Combining interdependent climate model outputs in cmip5: A spatial bayesian approach, 2020.
- [34] Manmeet Singh, Chirag Dhara, Adarsh Kumar, Sukhpal Singh Gill, and Steve Uhlig. Quantum artificial intelligence for the science of climate change, 2021.
- [35] Casey Berger, Agustin Di Paolo, Tracey Forrest, Stuart Hadfield, Nicolas Sawaya, Michał Stęchły, and Karl Thibault. Quantum technologies for climate change: Preliminary assessment, 2021.
- [36] Valerio Lucarini and Mickaël Chekroun. Theoretical tools for understanding the climate crisis from hasselmann’s program and beyond, 2023.
- [37] Natalie J. Briscoe, Shane D. Morris, Paul D. Mathewson, Lauren B. Buckley, Marko Jusup, Ofir Levy, Ilya M. D. Maclean, Sylvain Pincebourde, Eric A. Riddell, Jessica A. Roberts, Rafael Schouten, Michael W. Sears, and Michael R. Kearney. Mechanistic forecasts of species responses to climate change: the promise of biophysical ecology, 2022.
- [38] Sylvain Gretchko, Jessa Marley, and Rebecca C. Tyson. The effects of climate change on predator-prey dynamics, 2018.
- [39] Tamsin L. Edwards, Michel Crucifix, and Sandy P. Harrison. Using the past to constrain the future: how the palaeorecord can improve estimates of global warming, 2012.

- 
- [40] Adam Giammarese, Jacob Brown, and Nishant Malik. Reconfiguration of amazon's connectivity in the climate system, 2023.
- [41] Hyun Choi, Ali Sadeghian, Sergio Marconi, Ethan White, and Daisy Zhe Wang. Measuring impact of climate change on tree species: analysis of jsdm on fia data, 2019.
- [42] Hideyasu Shimadzu, Maria Dornelas, and Anne E. Magurran. Measuring temporal turnover in ecological communities, 2016.
- [43] Lionel Roques, Alain Roques, Henri Berestycki, and André Kretzschmar. A population facing climate change: joint influences of allee effects and environmental boundary geometry, 2009.
- [44] R. Juhász and B. Oborny. Percolation theory suggests some general features in range margins across environmental gradients, 2020.
- [45] Henri Berestycki, Romain Ducasse, and Luca Rossi. Influence of a road on a population in an ecological niche facing climate change, 2019.
- [46] Matthew Davidow, Cory Merow, Judy Che-Castaldo, Toryn Schafer, Marie-Christine Duker, Derek Corcoran, and David Matteson. Clustering future scenarios based on predicted range maps, 2022.
- [47] Jingfang Fan, Jun Meng, Yosef Ashkenazy, Shlomo Havlin, and Hans Joachim Schellnhuber. Climate network percolation reveals the expansion and weakening of the tropical component under global warming, 2018.
- [48] Meihua Zhou, Nan Wan, Tianlong Zheng, Hanwen Xu, Li Yang, and Tingting Wang. New insights into global warming: End-to-end visual analysis and prediction of temperature variations, 2024.
- [49] Greta T Pecl, Miguel B Araújo, Johann D Bell, Julia Blanchard, Timothy C Bonebrake, I-Ching Chen, Timothy D Clark, Robert K Colwell, Finn Danielsen, Birgitta Evengård, et al. Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, 355(6332):eaai9214, 2017.
- [50] Sergey A. Vakulenko, Ivan Sudakov, Sergei V. Petrovskii, and Dmitry V. Lukichev. Stability of a planetary climate system with the biosphere competing for resources, 2020.
- [51] Mauro Gobbi. Global warning: challenges, threats and opportunities for ground beetles (coleoptera: Carabidae) in high altitude habitats, 2020.
- [52] Lynn Wahab, Ezzat Chebaro, Jad Ismail, Amir Nasrelddine, and Ali El-Zein. A predictive model for geographic distributions of mangroves, 2021.
- [53] Chirag Dhara. Constraining global changes in temperature and precipitation from observable changes in surface radiative heating, 2020.
- [54] J. M. Finkel and J. I. Katz. Changing u.s. extreme temperature statistics, 2017.
- [55] James Hansen, Makiko Sato, Pushker Kharecha, Karina von Schuckmann, David J Beerling, Junji Cao, Shaun Marcott, Valerie Masson-Delmotte, Michael J Prather, Eelco J Rohling, Jeremy Shakun, and Pete Smith. Young people's burden: Requirement of negative co2 emissions, 2017.
- [56] Junze Zhang, Kerry Zhang, Mary Zhang, Jonathan H. Jiang, Philip E. Rosen, and Kristen A. Fahy. Avoiding the "great filter": An assessment of climate change solutions and combinations for effective implementation, 2022.
- [57] Simone Vincenzi and Michele Bellingeri. Consequences of catastrophic disturbances on population persistence and adaptations, 2011.
- [58] Zhu Liu, Zhu Deng, Philippe Ciais, Jianguang Tan, Biqing Zhu, Steven J. Davis, Robbie Andrew, Olivier Boucher, Simon Ben Arous, Pep Canadel, Xinyu Dou, Pierre Friedlingstein, Pierre Gentine, Rui Guo, Chaopeng Hong, Robert B. Jackson, Daniel M. Kammen, Piyu Ke, Corinne Le Quere, Crippa Monica, Greet Janssens-Maenhout, Glen Peters, Katsumasa Tanaka, Yilong Wang, Bo Zheng, Haiwang Zhong, Taochun Sun, and Hans Joachim Schellnhuber. Global daily co<sub>2</sub> emissions for the year 2020, 2021.

- 
- [59] Nicola Scafetta. Global temperatures and sunspot numbers. are they related? yes, but non linearly. a reply to gil-alana et al. (2014), 2014.
- [60] Adriano Mazzarella and Nicola Scafetta. Evidences for a quasi 60-year north atlantic oscillation since 1700 and its meaning for global climate change, 2012.
- [61] Andreas Morr and Niklas Boers. Detection of approaching critical transitions in natural systems driven by red noise, 2024.
- [62] David Dunn and James P. Crutchfield. Entomogenic climate change, 2008.
- [63] Sandra Skendžić, Monika Zovko, Ivana Pajač Živković, Vinko Lešić, and Darija Lemić. The impact of climate change on agricultural insect pests. *Insects*, 12(5):440, 2021.
- [64] Clément Le Priol, Joy M. Monteiro, and Freddy Bouchet. Using rare event algorithms to understand the statistics and dynamics of extreme heatwave seasons in south asia, 2024.
- [65] Zhiang Xie, Dongwei Chen, and Puxi Li. Rediscover climate change during global warming slowdown via wasserstein stability analysis, 2023.
- [66] Frederik Van Daele, Olivier Honnay, Steven Janssens, and Hanne De Kort. Habitat fragmentation affects climate adaptation in a forest herb, 2023.
- [67] Jun Chen, Jordy O Rodriguez Rincon, Gloria DeGrandi-Hoffman, Jennifer Fewell, Jon Harrison, and Yun Kang. Impacts of seasonality and parasitism on honey bee population dynamics, 2023.
- [68] Fernando M. Aragón, Francisco Oteiza, and Juan Pablo Rud. Climate change and agriculture: Subsistence farmers’ response to extreme heat, 2019.
- [69] Sesa Singha Roy, Reid Tingley, and Alan Dorin. Image background assessment as a novel technique for insect microhabitat identification, 2023.
- [70] M. Tuomi, T. Thum, H. Järvinen, S. Fronzek, B. Berg, M. Harmon, J. A. Trofymow, S. Sevanto, and J. Liski. Leaf litter decomposition – estimates of global variability based on yasso07 model, 2009.
- [71] Tyler Poppenwimer, Itay Mayrose, and Niv DeMalach. Revising the global biogeography of annual and perennial plants, 2023.
- [72] L. Mark W. Leggett and David. A. Ball. Forecasts of the trend in global-mean temperature to 2100 arising from the scenarios of first-difference co2 and peak fossil fuel, 2020.
- [73] Claire L. Harris, Neil Brummitt, Christina A. Cobbold, and Richard Reeve. Dynamic virtual ecosystems as a tool for detecting large-scale responses of biodiversity to environmental and land-use change, 2023.
- [74] Anthony J. Webster. Lightning strikes and attribution of climatic change, 2015.
- [75] Kgomo S. Morupisi and Chris J. Budd. An analysis of the periodically forced pp04 climate model, using the theory of non-smooth dynamical systems, 2020.
- [76] Sebastian Sippel, Jakob Zscheischler, Martin Heimann, Holger Lange, Miguel D. Mahecha, Geert Jan van Oldenborgh, Friederike E. L. Otto, and Markus Reichstein. Have precipitation extremes and annual totals been increasing in the world’s dry regions over the last 60 years?, 2016.
- [77] Angie Michaiel and Amy Bernard. Neurobiology and changing ecosystems: toward understanding the impact of anthropogenic influences on neurons and circuits, 2022.
- [78] R. Juhász and I. A. Kovács. Population boundary across an environmental gradient: Effects of quenched disorder, 2020.
- [79] Donald C. Morton. An astronomer’s view of climate change, 2014.
- [80] Gerald E. Marsh. Climate change: The sun’s role, 2007.

- 
- [81] Ranjini Swaminathan, Jacob Schewe, Jeremy Walton, Klaus Zimmermann, Colin Jones, Richard A. Betts, Chantelle Burton, Chris D. Jones, Matthias Mengel, Christopher P. O. Reyer, Andrew G. Turner, and Katja Weigel. Regional impacts poorly constrained by climate sensitivity, 2024.
- [82] Samer Fawzy, Ahmed I Osman, John Doran, and David W Rooney. Strategies for mitigation of climate change: a review. *Environmental Chemistry Letters*, 18:2069–2094, 2020.
- [83] B G Sidharth and B S Lakshmi. A model for long term climate change, 2003.
- [84] R. C. T. Rainey. The earths long-term climate changes and ice ages: a derivation of milankovitch cycles from first principles, 2021.
- [85] F. M. Sabatini, R. B. de Andrade, Y. Paillet, P. Odor, C. Bouget, T. Campagnaro, F. Gosselin, P. Janssen, W. Mattioli, J. Nascimbene, T. Sitzia, T. Kuemmerle, and S. Burrascano. Trade-offs between carbon stocks and biodiversity in european temperate forests, 2019.

---

**Disclaimer:**

SurveyX is an AI-powered system designed to automate the generation of surveys. While it aims to produce high-quality, coherent, and comprehensive surveys with accurate citations, the final output is derived from the AI's synthesis of pre-processed materials, which may contain limitations or inaccuracies. As such, the generated content should not be used for academic publication or formal submissions and must be independently reviewed and verified. The developers of SurveyX do not assume responsibility for any errors or consequences arising from the use of the generated surveys.

www.SurveyX.cn