
Forest Fire Early Warning and Monitoring Technology: A Survey

www.surveyx.cn

Abstract

Forest fire early warning and monitoring technologies, incorporating remote sensing and fire detection systems, are pivotal in addressing the increasing frequency and intensity of wildfires exacerbated by climate change. This survey explores the integration of satellite imagery, IoT, and advanced analytical frameworks to enhance the detection, monitoring, and management of wildfires. The paper highlights the role of remote sensing in providing real-time data for strategic planning, emphasizing the importance of machine learning and AI in improving predictive capabilities. Key advancements include the development of innovative frameworks like Multiple Instance Multi-Resolution Fusion (MIMRF) and the integration of deep learning models, which enhance the accuracy and timeliness of fire detection. However, challenges such as data quality, integration, and technological limitations persist, necessitating further research into adaptive models and robust evaluation frameworks. The survey suggests that future directions should focus on improving data fusion techniques, enhancing model robustness, and leveraging emerging technologies to overcome existing constraints. By advancing these technologies, wildfire management strategies can be significantly improved, offering more effective interventions to safeguard ecosystems and communities from the devastating impacts of forest fires.

1 Introduction

1.1 Importance of Forest Fire Early Warning and Monitoring

Early warning and monitoring systems are crucial for managing forest fires, especially as climate change and extreme weather conditions increase their frequency and intensity. These systems play a vital role in mitigating the severe environmental, economic, and social impacts of wildfires, which are significant natural hazards globally [1]. The heterogeneous effects of wildfires, shaped by geographical, climatic, and land cover variables, demand robust and adaptable monitoring frameworks [1].

In Australia, for instance, traditional fire management approaches have proven insufficient against the extensive damage caused by bushfires, highlighting the need for innovative predictive models and enhanced monitoring capabilities [2]. The historical absence of effective early warning systems has led to recurring fire events, underscoring the necessity for advancements in this field [3].

The integration of advanced remote sensing technologies and digital tools, including digital twins, supports timely interventions that can prevent significant damage [4]. These technologies enable rapid data processing and informed decision-making, which are essential for effective disaster response [5]. As artificial intelligence increasingly influences remote sensing, ensuring the security and reliability of these systems becomes paramount [6].




Figure 1: chapter structure

1.2 Role of Remote Sensing and Fire Detection Systems

Remote sensing and fire detection systems are integral to modern wildfire management, providing comprehensive capabilities for monitoring, predicting, and mitigating forest fire impacts. The combination of remote sensing technologies with advanced data analytics, such as machine learning, has significantly improved fire prediction and severity assessment. High-resolution satellite imagery and UAV data facilitate real-time monitoring and precise mapping of burned areas, enhancing timely interventions and strategic planning initiatives. Utilizing cutting-edge technologies and algorithms, including the U-Net model and the Sen2Fire dataset, these systems enhance wildfire detection and burned-area mapping accuracy, ultimately aiding in the mitigation of wildfires' environmental, economic, and social consequences [7, 8].

A core function of remote sensing in wildfire management is modeling vegetation recovery and quantifying post-wildfire changes. Techniques such as Functional Data Analysis (FDA) model recovery as a continuous function over time, considering pre-wildfire conditions. Additionally, integrating remote sensing data with IoT-based automated systems improves the detection of fire, humidity, and gas levels, supporting proactive fire management strategies [3].

While artificial intelligence enhances remote sensing applications, it also introduces security challenges, including adversarial and backdoor attacks that threaten reliability in critical contexts [6]. Addressing these vulnerabilities is essential for maintaining the integrity of AI applications in wildfire management.

Moreover, explainable artificial intelligence (XAI) techniques are increasingly utilized to enhance the transparency of decision-making processes in remote sensing image classification. These techniques are vital for selecting appropriate methods to interpret model outputs, thereby improving the interpretability and trustworthiness of remote sensing data analyses [9].

1.3 Structure of the Survey

This survey provides a comprehensive overview of forest fire early warning and monitoring technologies, emphasizing remote sensing and fire detection systems. It begins with an introduction that highlights the significance of these technologies in mitigating wildfire impacts, followed by a discussion on the critical role of remote sensing and fire detection systems in wildfire management.

The second section, "Background and Core Concepts," addresses foundational concepts and terminologies essential for understanding forest fire monitoring technologies, explaining their integration to enhance wildfire management.

The third section, "Remote Sensing Technologies in Fire Detection," explores various remote sensing technologies used in fire detection, discussing their advantages and limitations. This section is further divided into subsections covering emerging remote sensing technologies and innovative applications of satellite imagery.

The fourth section, "Fire Detection Systems," examines different fire detection systems employed in early warning and monitoring, highlighting the technologies and methodologies used for effective wildfire detection and monitoring. It includes subsections on automated monitoring systems and the application of machine learning and deep learning approaches.

Section five, "Integration of Technologies for Wildfire Management," discusses the integration of remote sensing and fire detection systems to enhance wildfire management strategies. It provides an overview of recent advancements in remote sensing, focusing on data fusion and analysis frameworks, particularly in integrating diverse data sources such as social media and remote sensing imagery. This section includes case studies of successful implementations and a meta-analysis of image fusion techniques, alongside a systematic review of explainable AI in remote sensing, emphasizing emerging methodologies and ethical considerations in the field [10, 11, 12, 13, 14].

The sixth section, "Challenges and Future Directions," identifies current challenges in forest fire early warning and monitoring technology, discussing potential advancements to enhance effectiveness. This section addresses data quality and integration challenges, technological limitations and scalability, future research directions in remote sensing, and strategies for improving model robustness and evaluation.

The conclusion synthesizes principal findings related to the integration of advanced technologies in forest fire management, emphasizing the critical role of generative modeling and remote sensing in accurately characterizing mid-story fuels and predicting fire behavior. It highlights the need for improved data collection methods, such as terrestrial LiDAR, to enhance fire outcome prediction reliability. Additionally, it suggests promising avenues for future research, including the development of UAV-based swarm coordination models for real-time fire prediction and targeted firefighting efforts in high-risk areas, reinforcing the importance of technological innovation in mitigating forest fire impacts [2, 15, 16, 14, 17]. The following sections are organized as shown in Figure 1.

2 Background and Core Concepts

2.1 Core Concepts in Fire Monitoring

Effective wildfire management hinges on accurate fire monitoring, which involves detecting, assessing, and predicting fire behavior and impacts. The Leaf Area Index (LAI) is crucial for ecological modeling and climate change research, quantifying leaf area per unit ground area [18]. Accurate mapping of burned areas is essential for evaluating fire damage and planning recovery, particularly as climate change exacerbates wildfire frequency and severity [1]. Remote sensing technologies provide extensive data on land cover and fire dynamics, though processing large datasets from satellites like Sentinel-2 remains challenging [19]. The Sen2Fire benchmark addresses wildfire detection using satellite imagery, simulating classification of image patches as fire or non-fire [8]. Incorporating deep learning into remote sensing enhances data analysis, especially in managing complex datasets [20], yet traditional object detection struggles with imaging variations, necessitating adaptive approaches [21].

Modeling wildland fire spread through mathematical simulations is vital for effective management [22]. Characterizing mid-story fuels, often obscured by canopies, challenges reliable fire outcome predictions [16]. Frameworks like RasterForge facilitate raster data manipulation and analysis, essential for wildfire management and remote sensing applications [23]. Earth Observation (EO) systems are critical for disaster surveillance and resource management, providing essential data for fire monitoring [5].

Detecting land cover change points using MODIS time series data is complicated by missing data and multivariate structures, requiring Bayesian change point models [24]. Deep learning solutions

enhance remote sensing image classification, improving accuracy in distinguishing forest from non-forest areas [25]. The scarcity of high-quality labeled training data for semantic segmentation in satellite images impacts model prediction accuracy [26], and adapting classification models to new datasets with differing conditions remains challenging [27].

Effective multimodal remote sensing data fusion enhances land cover classification, supported by benchmarks improving diverse data integration [28]. Temporal change retrieval in land cover is hindered by pixel-level computations, overlooking object-level characteristics [29].

2.2 Integration of Technologies

Integrating advanced technologies in wildfire management enhances fire detection, monitoring, and mitigation. Traditional methods relying on low-resolution imagery and rule-based algorithms face limitations in detecting small fires, leading to high errors [1]. Frameworks like the deep residual cycle GAN integrate heterogeneous spatio-spectral-temporal observations through forward fusion and backward degeneration feedback [14]. Foundation models, alongside traditional and deep learning approaches, have advanced burn scar and flood mapping, enhancing understanding of wildfire impacts across remote sensing domains [1]. Machine learning integration with remote sensing data bolsters predictive capabilities and response strategies, providing a robust framework for fire monitoring technology [5].

The Multiple Instance Multi-Resolution Fusion (MIMRF) framework improves fire detection accuracy by fusing outputs from multiple sensors and learning from imprecisely labeled data, enhancing system timeliness and reliability [1]. The improved A* algorithm determines optimal drone escape routes, considering dynamic fire and smoke locations, enhancing drone operation safety in fire-prone areas [1]. A 7-band Back-Side Illuminated (BSI) CCD-in-CMOS sensor for simultaneous multispectral TDI capture enhances imaging capabilities, providing high-resolution data essential for accurate fire monitoring and management [1].

A decision-level fusion strategy based on uncertainty quantification improves weight assignment in classification tasks, enhancing remote sensing scene analysis credibility [27]. The EFDR-CS method uses conditional simulation to generate estimates from incomplete data, facilitating robust inference in remote sensing applications [1]. Integrating ground-based and satellite observations with particle dispersion modeling offers a comprehensive understanding of wildfire smoke transport and atmospheric effects [1].

The FDA-WR method captures continuous vegetation recovery post-wildfire through regression models accounting for temporal changes, providing insights into ecosystem recovery dynamics [22]. Integrated technologies collectively enhance wildfire management strategies, enabling precise and timely interventions. The complexity of models and unique properties of remote sensing data necessitate effective interpretative methods, as highlighted by research on safety-critical applications. The proposed spatiotemporal bilateral filter improves image quality by normalizing spectral differences across time-series images, leading to better classification outcomes [1]. Integrating mobile phone records, fire detection data, and night lights facilitates analysis of human activity and economic patterns [1]. Utilizing multi-spectral satellite images from Sentinel-2 and Landsat 8 underscores integrated technologies' potential in enhancing wildfire management [8].

The SatViT-V2 model architecture optimizes pretraining and patch sizes, improving performance metrics across various biomes [1]. This development exemplifies the ongoing evolution of remote sensing technologies in enhancing wildfire management. Integrating multispectral satellite imagery from diverse sources improves spatio-temporal coverage and measurement accuracy, facilitating comprehensive wildfire monitoring [1].

The integration of advanced technologies in fire detection has become increasingly vital in recent years. As illustrated in Figure 2, the hierarchical structure of remote sensing technologies plays a crucial role in this domain. This figure categorizes emerging technologies and innovative uses of satellite imagery, emphasizing key technological advancements. Additionally, it showcases the applications of machine learning and enhanced image analysis techniques, which collectively contribute to improved wildfire detection and management. The visual representation not only clarifies the relationships between different technologies but also underscores the significance of these advancements in addressing the challenges posed by wildfires.




Figure 2: This figure illustrates the hierarchical structure of remote sensing technologies in fire detection, categorizing emerging technologies and innovative uses of satellite imagery. It highlights key technological advancements, machine learning applications, and enhanced image analysis techniques that collectively contribute to improved wildfire detection and management.

3 Remote Sensing Technologies in Fire Detection

3.1 Emerging Remote Sensing Technologies

Advancements in remote sensing technologies have significantly enhanced wildfire detection and monitoring, addressing the limitations of traditional methods. The integration of multi-source data streams, notably from Sentinel-1 and Sentinel-2 satellites, illustrates innovative approaches by providing multi-temporal data crucial for estimating the Leaf Area Index (LAI) and mitigating cloud cover challenges, thereby improving vegetation monitoring accuracy in fire-prone regions [18]. The Sen2Fire dataset, containing extensive image patches from Sentinel-2 and Sentinel-5P, is vital for developing and benchmarking fire detection algorithms [8].

The OBSUM framework captures significant temporal changes using object-level information, achieving superior fusion accuracy for mapping fire-affected zones and assessing ecological impacts [29]. Additionally, the GSC method leverages time-series satellite data to analyze vegetation health indices before and after wildfires, offering insights into the heterogeneous effects of wildfires on ecosystems [1].

Emerging technologies also include lightweight geospatial indexing algorithms like GeoHash, QuadTree, and Orthogonal List on HDFS, optimizing indexing performance without complex setups for efficient remote sensing data processing [30]. The MISR model enhances multispectral satellite imagery quality, outperforming traditional single-image super-resolution methods [31].

TorchGeo, a Python library, integrates geospatial data into the PyTorch ecosystem, providing tools for multispectral imagery analysis and supporting advanced remote sensing model development [32]. The HRSemI2I model improves segmentation performance by adapting Landsat 8 images to the style of Sentinel-2 images, showcasing the potential of image-to-image translation in remote sensing data analysis [27].

On-Air Deep Learning Integrated Semantic Inference Models (ODL-ISIM) optimize data transmission in Earth Observation networks using deep learning and semantic processing, enabling real-time monitoring and automated responses to environmental changes [5]. Furthermore, the Markov random field approximation to thin plate spline offers a novel approach to modeling discrepancies in proxy data, advancing fire detection technologies [33].

As illustrated in Figure 3, the hierarchical structure of emerging remote sensing technologies focuses on the integration of satellite data, advancements in geospatial indexing, and the application of advanced analysis techniques. The key technologies and methodologies are categorized into three main areas, highlighting their contributions to wildfire detection and management. Technological advancements in remote sensing and machine learning mark a transformative evolution in wildfire detection and management. Innovations such as generative modeling for mid-story fuel characterization enhance fire behavior prediction accuracy using terrestrial LiDAR data. The integration of remote sensing with machine learning techniques, such as the XGBoost model, has achieved 86.13




Figure 3: This figure illustrates the hierarchical structure of emerging remote sensing technologies, focusing on the integration of satellite data, advancements in geospatial indexing, and the application of advanced analysis techniques. The key technologies and methodologies are categorized into three main areas, highlighting their contributions to wildfire detection and management.

3.2 Innovative Use of Satellite Imagery

The innovative application of satellite imagery in wildfire detection and monitoring has advanced through sophisticated models and frameworks that enhance data analysis accuracy and efficiency. Machine learning-based entropy approximations, such as SvdEn2D, PermEn2D, SampEn2D, and NNetEn2D, significantly reduce computation time, enabling rapid satellite data processing for fire monitoring [35].

Remote sensing image classification has improved with specific configurations of feature points (FPs), which generally outperform anchor points (APs), resulting in higher classification accuracies in fire-affected regions [36]. Graph-based fusion techniques for change detection, applied to NIR and

RED band images from Landsat satellites, enhance precision while reducing false alarms and missed detections [12].

The Sen2Chain toolbox exemplifies an open-source approach for automating the download, preprocessing, and analysis of Sentinel-2 images, generating invaluable time-series data for environmental monitoring applications, including wildfire detection [19]. The RL-AOD method employs deep reinforcement learning to adaptively fine-tune image attributes, improving object detection performance in remote sensing imagery [21].

Innovative models like S2FL demonstrate superior classification accuracy across benchmark datasets, underscoring the potential of multimodal remote sensing data integration in wildfire monitoring [28]. The Sen2Fire dataset supports the development of fire detection algorithms by providing a challenging benchmark with optimal band combinations, particularly in the SWIR composite strategy [8].

The TorchGeo library facilitates geospatial data integration into the PyTorch ecosystem, enhancing the analysis and interpretation of complex datasets [32]. The HRSemI2I method employs style transfer to align the spectral characteristics of Landsat 8 images with those of Sentinel-2 images, ensuring semantic consistency and improving domain adaptation [27].

The innovative applications of satellite imagery, particularly through advanced computational techniques like the U-Net model and Visual ChatGPT, highlight a growing trend in wildfire detection, monitoring, and management. These approaches leverage cutting-edge remote sensing technology and artificial intelligence to automate burned-area mapping, significantly enhancing real-time fire monitoring capabilities. Optimized models can achieve high accuracy in mapping burned areas, with the AllSizes model yielding a Dice Coefficient of 0.93. Remote sensing plays a crucial role in assessing post-wildfire vegetation recovery, revealing that recovery dynamics are influenced by various pre-wildfire conditions. The integration of natural language processing with visual computation, exemplified in Visual ChatGPT, opens new avenues for image analysis in remote sensing, promising improved interpretation and extraction of critical information from satellite images. These advancements collectively present promising solutions to the complex challenges posed by wildfires, enhancing our ability to mitigate their environmental, economic, and social impacts [7, 37, 34].

4 Fire Detection Systems

4.1 Automated Monitoring Systems

Method Name	Technological Components	Data Processing Methods	Modeling Techniques
LSTM-DT[4]	Remote Sensing	Feature Extraction	Lstm-based Forecasting
DSS[38]	Multi-spectral Sensors	Sparse Feature Vector	Dynamic Semantic Segmentation
ODL-ISIM[5]	Onboard Sensors	Semantic Extraction	Deep Learning
GMMF[16]	Remote Sensing	Gaussian Process Modeling	Generative Model
BCPM[24]	Modis Time Series	Expectation-Maximization Algorithm	Bayesian Change Point
MIX-RS[30]	Satellite Technologies	Hadoop Mapreduce	Bayesian Change Point

Table 1: Comparison of various automated monitoring systems deployed in wildfire management, detailing their technological components, data processing methods, and modeling techniques. This table highlights the integration of remote sensing technologies and machine learning algorithms to enhance detection and response efficiency.

Automated monitoring systems are pivotal in modern wildfire management, enhancing early detection and response efficiency through advanced remote sensing and machine learning algorithms. Multispectral CCD-in-CMOS sensors are integral, offering reduced power consumption and compact design for effective Earth observation [39]. These systems adapt to dynamic conditions, crucial for real-time updates and strategic route planning, particularly when employing digital twin technologies and LSTM-based methods to forecast future remote sensing images [4].

As illustrated in Figure 4, the hierarchical structure of these automated monitoring systems encompasses key technological components, data processing methods, and modeling and analysis techniques. Edge computing and heterogeneous fire propagation models enhance reliability by accounting for diverse dynamics across various landscapes [17]. Content-based image retrieval methods like DSS improve computational efficiency, facilitating the analysis of large satellite datasets [38]. The ODL-

ISIM model streamlines data processing and transmission, supporting timely decision-making [5], while tools like RasterForge enhance raster data handling for researchers [23].

These systems effectively generate representative mid-story fuel layouts using non-homogeneous Boolean processes and Gaussian process modeling, crucial for understanding wildfire behavior [16]. The Bayesian change point model enhances detection by utilizing data structure and incorporating prior land cover knowledge [24]. The MIX-RS method optimizes query response times by constructing multiple geospatial indices in parallel, improving automated monitoring performance [30]. The U-Net architecture, tested on the Sen2Fire dataset, demonstrates the potential of advanced models to enhance wildfire detection [8].

Table 1 provides a comprehensive comparison of different automated monitoring systems used in wildfire management, focusing on their technological components, data processing methods, and modeling techniques.




Figure 4: This figure illustrates the hierarchical structure of automated monitoring systems in wildfire management, highlighting key technological components, data processing methods, and modeling and analysis techniques.

4.2 Machine Learning and Deep Learning Approaches

Method Name	Model Types	Data Utilization	Performance Challenges
AGNI[40]	Neural Network	Remote Sensing Data	Missing Data
XGBoost[2]	Xgboost	Satellite Imagery	Resource-constrained Devices
OTB-TF[41]	Fully Convolutional Model	Remote Sensing Images	Memory Footprint
ML_Ent[35]	Gradient Boosting	Sentinel-2 Images	Longer Time Series
PISM[42]	-	Satellite Data	Computational Efficiency
LAI-DL[18]	U-nets	Satellite Data	Resource-constrained Devices

Table 2: Overview of machine learning and deep learning methodologies employed in wildfire detection and monitoring. The table highlights various models, their data utilization strategies, and the specific performance challenges they encounter. Notable methods include neural networks, gradient boosting, and U-nets, each addressing different aspects of fire detection and prediction.

Machine learning and deep learning methodologies have significantly improved fire detection systems, enhancing their ability to accurately identify and monitor wildfires. The Agni model, for instance, leverages machine learning to outperform traditional feature-based models in fire detection [40]. The XGBoost model effectively integrates diverse datasets to model fire severity [2]. Deep learning frameworks like CNNs and RNNs process remote sensing images efficiently, extracting complex patterns and temporal dynamics for improved detection accuracy [41]. Table 2 provides a comprehensive comparison of different machine learning and deep learning approaches used in wildfire detection, detailing their model types, data sources, and associated performance challenges.

Machine learning regression techniques enhance the speed and efficiency of entropy calculations in remote sensing applications [35]. Physics-informed statistical models with multi-source satellite data model aerosol optical depth propagation, providing insights into fire behavior and atmospheric impacts [42]. Deep learning models like ResNet and DenseNet have improved classification accuracy and visual quality in remote sensing data analysis [25].

Advancements in model compression and efficiency have facilitated the deployment of high-performing models on resource-constrained devices, enabling real-time fire detection in challenging environments [43]. Deep neural network architectures, such as U-nets, enhance prediction accuracy, particularly in estimating the Leaf Area Index from satellite data [18]. However, challenges remain in aligning self-supervised learning tasks with pixel-level classification requirements, highlighting the complexity of training large foundation models without sufficient data [44]. Addressing these challenges is crucial for the ongoing development of machine learning and deep learning approaches in fire detection systems, ensuring their efficacy and reliability across diverse contexts.

5 Integration of Technologies for Wildfire Management

5.1 Advancements in Remote Sensing and Fire Detection Integration

Recent advancements in integrating remote sensing with fire detection technologies have substantially improved wildfire management. The synergy of deep learning with semantic communication optimizes Earth Observation (EO) data management, enhancing wildfire monitoring efficiency [5]. This integration streamlines data processing, enabling prompt responses to fire incidents.

Markov Random Field (MRF) models enhance the integration of remote sensing data by capturing spatial variability, crucial for accurate wildfire-prone landscape modeling [33]. Physics-informed statistical models, which blend physical principles with statistical methods, manage data heterogeneity and increase prediction accuracy, particularly in forecasting atmospheric aerosols from wildfires [33, 11, 45, 42, 14]. These models, along with Gaussian Process regression and latent force models, address errors in multi-source datasets, enhancing robustness and interpreting complex dynamics. Transfer learning further aids model adaptation across environments.

Tools like Sen2Chain and advanced computational methods, such as entropy approximation, improve access to remote sensing data and enhance real-time management capacity. The integration of AI with remote sensing, exemplified by the U-Net model for burned-area mapping and the Sen2Fire dataset, optimizes monitoring capabilities, enabling timely responses to wildfires [7, 16, 8].

Future research should enhance sensor accuracy, system resilience, and explore IoT applications for environmental monitoring. Utilizing diverse data sources, including social media and remote sensing imagery, will refine geo-information extraction. Advanced machine learning, particularly deep learning, is vital for addressing large-scale environmental challenges. Collaborative methodology and dataset sharing will foster innovative monitoring systems, improving fire detection and response mechanisms, thereby advancing environmental protection and management [3, 6, 34, 14, 20].


5.2 Frameworks for Data Fusion and Analysis

Data fusion frameworks are essential for enhancing wildfire management systems' accuracy and efficiency. By integrating data from multiple sources, these frameworks provide comprehensive insights into fire dynamics, improving decision-making. The Multiple Instance Multi-Resolution Fusion (MIMRF) framework exemplifies this by integrating sensor outputs to enhance detection reliability, effectively handling imprecise labels [1].

Machine learning models combined with remote sensing data bolster predictive capabilities, supporting robust monitoring strategies [5]. The OBSUM framework, utilizing object-level information, achieves superior fusion accuracy in mapping fire-affected areas and ecological changes [29]. Graph-based fusion techniques for change detection improve precision, reducing false alarms and missed detections, as demonstrated with Landsat data [12].

Deep learning models integrated with semantic communication principles enhance data fusion frameworks, facilitating real-time wildfire responses [5]. MRF specifications improve data fusion processes, enabling precise fire dynamics monitoring [33].

Figure 5 illustrates the transformative potential of data fusion frameworks in wildfire management. The first subfigure shows satellite image comparisons, highlighting multi-resolution and multi-modal sensor fusion's ability to provide detailed visual data for wildfire risk assessment. The second subfigure contrasts network architectures, demonstrating neural networks' capability for large-scale data analysis essential for effective management [46, 11].



(a) Comparison of Two Satellite Images of a Residential Area[46] (b) Comparison of Two Network Architectures[11]

Figure 5: Examples of Frameworks for Data Fusion and Analysis

5.3 Case Studies and Successful Implementations

The successful integration of remote sensing and fire detection technologies has provided valuable insights into wildfire management strategies. The Sen2Fire benchmark dataset has advanced model development and evaluation, enhancing performance across diverse biomes through improved architectures like SatViT-V2 [8, 1].

The MIMRF framework improves detection accuracy by fusing sensor outputs and learning from imprecisely labeled data, offering timely alerts and responses to forest fires [1]. The OBSUM framework, leveraging object-level information, accurately maps fire-affected areas and ecological changes, demonstrating its potential in remote sensing applications [29].

On-Air Deep Learning Integrated Semantic Inference Models (ODL-ISIM) optimize data transmission in EO networks, facilitating real-time monitoring and automated environmental responses [5]. These case studies highlight the impact of integrated technologies in advancing wildfire management strategies.

Advanced data integration and machine learning techniques have significantly improved fire detection and monitoring systems' accuracy, timeliness, and reliability. Combining remote sensing data with machine learning, such as Landsat imagery and spectral indices, has led to predictive models with accuracy levels above 86

6 Challenges and Future Directions

Understanding the complexities of wildfire management requires addressing challenges related to data quality and integration, which are crucial for the effectiveness of fire monitoring systems. This section delves into the limitations posed by incomplete datasets and the integration of diverse data sources, which are pivotal in overcoming obstacles that hinder the advancement of these technologies.

6.1 Data Quality and Integration Challenges

Data quality and integration are critical challenges that impact the development of effective fire monitoring systems. The reliability of models, such as Agni, is contingent on comprehensive input data, with issues like missing satellite data and cloud cover affecting predictions and ecological assessments, including the Leaf Area Index (LAI) [40]. As climate change continues to alter vegetation patterns and fire regimes, the dependence on historical data becomes problematic, necessitating adaptive models that can accommodate these environmental changes [2].

The integration of diverse data sources, including social media and traditional remote sensing, adds complexity. The sheer volume of social media data, coupled with licensing restrictions, complicates the merging of these datasets with conventional remote sensing data [14]. Spatially correlated errors in proxy data can obscure true signals, further complicating integration efforts [33]. Additionally, the lack of comprehensive evaluation metrics for satellite video object detection and tracking hampers algorithm performance assessment [47].

Active learning strategies rely on the quality of initial labeled data, which influences the efficacy of semi-supervised learning methods [26]. Moreover, inefficient processing of large Earth Observation (EO) datasets can delay decision-making, underscoring the need for improved data handling solutions [5]. Addressing these challenges is crucial for enhancing the reliability of fire monitoring systems, which depend on accurate datasets for applications ranging from environmental monitoring to early warning systems for humanitarian crises [15, 8, 16]. Future research should focus on developing adaptive models and comprehensive evaluation frameworks that can respond to dynamic environmental conditions and data sources.

6.2 Technological Limitations and Scalability

The integration of advanced technologies in wildfire management systems encounters significant challenges concerning technological limitations and scalability. Sensor malfunctions and the need for stable internet connectivity for data transmission pose substantial issues in remote areas affected by wildfires [3]. Weak supervision in existing studies often limits model accuracy and robustness, impacting the reliability of fire detection [48].

Methods like graph-based fusion techniques are sensitive to sample selection, which can hinder the detection of minor changes in dynamic environments [12]. Calibration challenges arise when observational data is scarce or of poor quality, affecting generative modeling approaches [16]. The assumption of stable land-cover types during data fusion may not hold, affecting the accuracy of models like OBSUM [29]. The MIX-RS framework, although effective, incurs higher memory consumption, presenting challenges in resource-constrained environments [30].

Continuous updates are necessary to maintain compatibility with evolving data formats and user needs, posing ongoing challenges for fire detection technologies [23]. The inability to detect disturbances not well represented in training data complicates model scalability [24]. While frameworks like TorchGeo address many geospatial data processing challenges, they may still struggle with extremely large datasets or specific edge cases critical for large-scale wildfire management [32]. Addressing these technological limitations and scalability issues is essential for enhancing wildfire management systems, necessitating continued research and development.

6.3 Future Research Directions in Remote Sensing

Future research in remote sensing technologies for wildfire monitoring is set to advance through several innovative avenues. Developing methodologies for improved data integration, addressing ethical concerns, and exploring emerging technologies in geo-information harvesting are key areas of focus [14]. Fully multi-spectral Multi-Image Super-Resolution (MISR) approaches and enhancing model robustness against artifacts from different imaging instruments are crucial for further investigation [31].

Enhancing model performance through additional data sources and improved handling of varying cloud cover conditions is vital for advancing remote sensing capabilities [18]. Effective input strategies and the integration of diverse data sources can significantly improve wildfire detection, as ongoing research demonstrates [8]. Hybrid models that leverage the strengths of various approaches, alongside improved data collection methods and new mathematical frameworks, are essential for better predictions in wildfire spread modeling [22].

Advancements in data augmentation techniques for geospatial imagery, refining model architectures to suit remote sensing characteristics, and expanding benchmark datasets will contribute to progress in this field [32]. Improving label maps and detection methods for challenging environments like wetlands is another critical research direction [27]. Optimizing the integration of large language models (LLMs) in Earth Observation contexts is crucial for enhancing data labeling and training efficiency, ultimately improving fire monitoring and response strategies [5].

Enhancing model identifiability and exploring alternative modeling strategies are vital for advancing remote sensing technologies. Investigating potential research areas focused on improving the accuracy and reliability of remote sensing models will contribute to innovative solutions for the complex challenges of wildfire management through improved data integration, machine learning, and model optimization [33].

6.4 Improving Model Robustness and Evaluation

Benchmark	Size	Domain	Task Format	Metric
AIDER[49]	8,540	Emergency Response	Image Classification	F1 Score
Prithvi[50]	1,000,000	Remote Sensing	Image Retrieval	mean Average Precision
SatViT-V2[51]	1,300,000	Land Cover Classification	Pixel Classification	Accuracy, Calibration
Prithvi[52]	944	Remote Sensing	Object Detection	mAP50
VISO[47]	1,646,038	Moving Object Detection	Multi-object Tracking	mAP, F1-score
FPS[36]	1,000,000	Remote Sensing	Classification	Overall Accuracy, Kappa Coefficient
CCBM[44]	5,567	Climate	Image Classification	F1, mIoU
DL-Forest[25]	2,400,000	Forest Classification	Binary Classification	ACC, IoU

Table 3: This table presents a comprehensive overview of various benchmarks utilized in the evaluation of remote sensing and image classification tasks. It details the size, domain, task format, and performance metrics for each benchmark, providing insights into their applicability and scope within the context of model robustness and evaluation in fire detection systems.

Enhancing the robustness and evaluation of models used in fire detection is critical for improving the reliability and accuracy of wildfire management systems. Developing pansharpening validation approaches that address existing drawbacks and ensure consistency across different validation scales is a promising research avenue [53]. This includes refining methods to effectively combine high-resolution panchromatic and multispectral data, thereby improving the spatial resolution and accuracy of remote sensing imagery used in fire detection.

To bolster model robustness, it is essential to develop adaptive algorithms that can dynamically respond to varying environmental conditions and data quality issues, such as cloud cover and sensor noise. These algorithms should incorporate advanced data fusion techniques to integrate diverse data sources, providing a comprehensive understanding of fire dynamics. Enhancing the interpretability of machine learning models through explainable AI techniques fosters trust and transparency in model outputs, supporting informed decision-making in wildfire management. Recent systematic reviews underscore the importance of understanding decision-making mechanisms in these models, revealing key trends and challenges in applying explainable AI within remote sensing. Quantitative analyses of various explainable AI methods can help identify the most effective approaches for elucidating model behavior, thereby facilitating better-informed strategies in managing wildfire risks and optimizing resource allocation [9, 13].

Establishing standardized benchmarks and metrics for evaluating fire detection models is essential. These metrics should assess model accuracy, precision, and recall while accounting for variability in fire scenarios and datasets, as demonstrated by studies utilizing generative modeling for mid-story fuel characterization, the Sen2Fire dataset for optimizing wildfire detection, and machine learning techniques for predicting bushfire severity across diverse ecosystems [2, 15, 16, 8]. Developing comprehensive datasets that capture the complexity of real-world fire events will provide a robust foundation for model training and validation. Table 3 enumerates key benchmarks that serve as standardized datasets for assessing the performance of models in remote sensing and image classification, highlighting their relevance to improving model robustness in wildfire management systems.

Integrating hybrid modeling approaches that leverage the strengths of diverse machine learning and statistical methods, such as generative modeling for fuel characterization and advanced predictive modeling using remote sensing data, can enhance the robustness and resilience of fire detection systems. By calibrating these models with high-fidelity data, including terrestrial LiDAR observations and spectral indices from satellite imagery, more accurate predictions of fire severity can be achieved, ultimately leading to improved fire management strategies and mitigation efforts [2, 16]. Leveraging the complementary capabilities of different models can improve predictive accuracy and reduce the likelihood of false positives and negatives.

7 Conclusion

The exploration of advanced technologies in forest fire early warning and monitoring has highlighted the transformative impact of integrating remote sensing and fire detection systems in wildfire management. These technologies provide crucial, real-time data that enhance decision-making processes, significantly contributing to effective fire prevention and response strategies. The deployment of

IoT-based systems has notably improved environmental monitoring capabilities, offering prompt alerts that are instrumental in forest fire prevention efforts. Progress in remote sensing, particularly through the use of satellite imagery and machine learning models, has refined the precision of wildfire detection and monitoring, presenting innovative solutions to the complexities associated with these natural hazards.

The integration of data fusion frameworks and advanced analytical techniques has further optimized the performance and accuracy of fire detection systems. This examination emphasizes the importance of overcoming challenges related to data quality, integration, and technological constraints to improve system efficacy. Future research should focus on developing adaptive models capable of responding to fluctuating environmental conditions and exploring the potential of emerging remote sensing technologies in wildfire management. Additionally, enhancing model robustness and evaluation through standardized benchmarks and hybrid modeling approaches will result in more dependable fire detection systems. Leveraging these technological advancements will significantly enhance forest fire management strategies, ultimately safeguarding ecosystems and communities from the destructive consequences of wildfires.

References

- [1] Feliu Serra-Burriel, Pedro Delicado, Andrew T. Prata, and Fernando M. Cucchiatti. Estimating heterogeneous wildfire effects using synthetic controls and satellite remote sensing, 2021.
- [2] Shouthiri Partheepan, Farzad Sanati, and Jahan Hassan. Bushfire severity modelling and future trend prediction across australia: Integrating remote sensing and machine learning, 2024.
- [3] Poltak Sihombing, Herriyance Herriyance, and M Rizki Syaputra. Smart system to prevent forest fire based on internet of things. *Internetworking Indonesia Journal*, 13(1):45–50, 2021.
- [4] Xuetao Jiang, Meiyu Jiang, YuChun Gou, Qian Li, and Qingguo Zhou. Forestry digital twin with machine learning in landsat 7 data, 2022.
- [5] Hong fu Chou, Vu Nguyen Ha, Prabhu Thiruvassagam, Thanh-Dung Le, Geoffrey Eappen, Ti Ti Nguyen, Luis M. Garces-Socarras, Jorge L. Gonzalez-Rios, Juan Carlos Merlano-Duncan, and Symeon Chatzinotas. On-air deep learning integrated semantic inference models for enhanced earth observation satellite networks, 2024.
- [6] Yonghao Xu, Tao Bai, Weikang Yu, Shizhen Chang, Peter M. Atkinson, and Pedram Ghamisi. Ai security for geoscience and remote sensing: Challenges and future trends, 2023.
- [7] Ian Mancilla-Wulff, Jaime Carrasco, Cristobal Pais, Alejandro Miranda, and Andres Weintraub. Two scalable approaches for burned-area mapping using u-net and landsat imagery, 2023.
- [8] Yonghao Xu, Amanda Berg, and Leif Haglund. Sen2fire: A challenging benchmark dataset for wildfire detection using sentinel data, 2024.
- [9] Akshatha Mohan and Joshua Peeples. Quantitative analysis of primary attribution explainable artificial intelligence methods for remote sensing image classification, 2023.
- [10] Hessah Albanwan, Rongjun Qin, and Yang Tang. Image fusion in remote sensing: An overview and meta analysis, 2024.
- [11] Luigi Mascolo, Marco Quartulli, Pietro Guccione, Giovanni Nico, and Igor G. Olaizola. Distributed mining of large scale remote sensing image archives on public computing infrastructures, 2015.
- [12] David Alejandro Jimenez Sierra, Hernán Darío Benítez Restrepo, Hernán Darío Vargas Cardonay, and Jocelyn Chanussot. Graph-based fusion for change detection in multi-spectral images, 2020.
- [13] Adrian Höhl, Ivica Obadic, Miguel Ángel Fernández Torres, Hiba Najjar, Dario Oliveira, Zeynep Akata, Andreas Dengel, and Xiao Xiang Zhu. Opening the black-box: A systematic review on explainable ai in remote sensing, 2024.
- [14] Xiao Xiang Zhu, Yuanyuan Wang, Mrinalini Kochupillai, Martin Werner, Matthias Häberle, Eike Jens Hoffmann, Hannes Taubenböck, Devis Tuia, Alex Levering, Nathan Jacobs, Anna Kruspe, and Karam Abdulahad. Geo-information harvesting from social media data, 2022.
- [15] David Pastor-Escuredo, Thierry Savy, and Miguel A. Luengo-Oroz. Can fires, night lights, and mobile phones reveal behavioral fingerprints useful for development?, 2015.
- [16] Grant Hutchings, James Gattiker, and Braden Scherting. Wildland fire mid-story: A generative modeling approach for representative fuels, 2023.
- [17] Laurent Hébert-Dufresne, Adam F. A. Pellegrini, Uttam Bhat, Sidney Redner, Stephen W. Pacala, and Andrew M. Berdahl. Edge fires drive the shape and stability of tropical forests, 2018.
- [18] Clement Wang, Antoine Debouchage, Valentin Goldité, Aurélien Wery, and Jules Salzinger. Leveraging multi-temporal sentinel 1 and 2 satellite data for leaf area index estimation with deep learning, 2024.

-
- [19] Christophe Revillion, Pascal Mouquet, Jeremy Commins, Juliette Miranville, Charlotte Wolff, Thomas Germain, Sylvaine Jegou, Lucas Longour, Florian Girond, Didier Bouche, Rodolphe Devillers, Gwenaëlle Pennober, and Vincent Herbreteau. Sen2chain: An open-source toolbox for processing sentinel-2 satellite images and producing time-series of spectral indices, 2024.
- [20] Xiao Xiang Zhu, Devis Tuia, Lichao Mou, Gui-Song Xia, Liangpei Zhang, Feng Xu, and Friedrich Fraundorfer. Deep learning in remote sensing: a review, 2017.
- [21] Nuo Xu, Chunlei Huo, Jiacheng Guo, Yiwei Liu, Jian Wang, and Chunhong Pan. Adaptive remote sensing image attribute learning for active object detection, 2021.
- [22] A. L. Sullivan. A review of wildland fire spread modelling, 1990-present 3: Mathematical analogues and simulation models, 2007.
- [23] Afonso Oliveira, Nuno Fachada, and João P. Matos-Carvalho. Raster forge: Interactive raster manipulation library and gui for python, 2024.
- [24] Hunter Glanz, Xiaoman Huang, Minhui Zheng, and Luis E. Carvalho. A bayesian change point model for detecting land cover changes in modis time series, 2015.
- [25] Antonio Mazza and Francescopaolo Sica. Deep learning solutions for tandem-x-based forest classification, 2019.
- [26] Shasvat Desai and Debasmita Ghose. Active learning for improved semi-supervised semantic segmentation in satellite images, 2021.
- [27] M. Sokolov, J. L. Storie, C. J. Henry, C. D. Storie, J. Cameron, R. S. Ødegård, V. Zubinaite, and S. Stikbakke. Semantically-consistent landsat 8 image to sentinel-2 image translation for alpine areas, 2022.
- [28] Danfeng Hong, Jingliang Hu, Jing Yao, Jocelyn Chanussot, and Xiao Xiang Zhu. Multimodal remote sensing benchmark datasets for land cover classification with a shared and specific feature learning model, 2021.
- [29] Houcai Guo, Dingqi Ye, and Lorenzo Bruzzone. Obsum: An object-based spatial unmixing model for spatiotemporal fusion of remote sensing images, 2023.
- [30] Jiashu Wu, Jingpan Xiong, Hao Dai, Yang Wang, and Chengzhong Xu. Mix-rs: A multi-indexing system based on hdfs for remote sensing data storage, 2022.
- [31] Muhammed Razzak, Gonzalo Mateo-Garcia, Luis Gómez-Chova, Yarin Gal, and Freddie Kalaitzis. Multi-spectral multi-image super-resolution of sentinel-2 with radiometric consistency losses and its effect on building delineation, 2021.
- [32] Adam J. Stewart, Caleb Robinson, Isaac A. Corley, Anthony Ortiz, Juan M. Lavista Ferres, and Arindam Banerjee. Torchgeo: Deep learning with geospatial data, 2022.
- [33] Christopher J. Paciorek. Combining spatial information sources while accounting for systematic errors in proxies, 2011.
- [34] Lucas Prado Osco, Eduardo Lopes de Lemos, Wesley Nunes Gonçalves, Ana Paula Marques Ramos, and José Marcato Junior. The potential of visual chatgpt for remote sensing, 2023.
- [35] Andrei Velichko, Maksim Belyaev, Matthias P. Wagner, and Alireza Taravat. Entropy approximation by machine learning regression: Application for irregularity evaluation of images in remote sensing, 2022.
- [36] Minh-Tan Pham, Erchan Aptoula, and Sébastien Lefèvre. Classification of remote sensing images using attribute profiles and feature profiles from different trees: a comparative study, 2018.
- [37] Feliu Serra-Burriel, Pedro Delicado, and Fernando M. Cucchietti. Wildfires vegetation recovery through satellite remote sensing and functional data analysis, 2021.

-
- [38] Suraj Kothawade, Kunjan Mhaske, Sahil Sharma, and Furkhan Shaikh. Content based image retrieval from awifs images repository of IRS Resourcesat-2 satellite based on water bodies and burnt areas, 2018.
- [39] Swaraj Bandhu Mahato, Steven Thijs, Jonas Bentell, Linkun Wu, Klaas Tack, Pierre Boulenc, Dorian Lasnet, Renaud Van Langendonck, and Piet De Moor. Multispectral ccd-in-cmos time delay integration imager for high resolution earth observation, 2021.
- [40] Suwei Yang, Massimo Lupascu, and Kuldeep S. Meel. Predicting forest fire using remote sensing data and machine learning, 2021.
- [41] Rémi Cresson. A framework for remote sensing images processing using deep learning technique, 2018.
- [42] Guanzhou Wei, Venkat Krishnan, Yu Xie, Manajit Sengupta, Yingchen Zhang, Haitao Liao, and Xiao Liu. Physics-informed statistical modeling for wildfire aerosols process using multi-source geostationary satellite remote-sensing data streams, 2022.
- [43] Yassine Himeur, Nour Aburaed, Omar Elharrouss, Iraklis Varlamis, Shadi Atalla, Wathiq Mansoor, and Hussain Al Ahmad. Applications of knowledge distillation in remote sensing: A survey, 2024.
- [44] Alexandre Lacoste, Evan David Sherwin, Hannah Kerner, Hamed Alemohammad, Björn Lütjens, Jeremy Irvin, David Dao, Alex Chang, Mehmet Gunturkun, Alexandre Drouin, Pau Rodriguez, and David Vazquez. Toward foundation models for earth monitoring: Proposal for a climate change benchmark, 2021.
- [45] Gustau Camps-Valls, Luca Martino, Daniel H. Svendsen, Manuel Campos-Taberner, Jordi Muñoz-Marí, Valero Laparra, David Luengo, and Francisco Javier García-Haro. Physics-aware gaussian processes in remote sensing, 2020.
- [46] Xiaoxiao Du and Alina Zare. Multi-resolution multi-modal sensor fusion for remote sensing data with label uncertainty, 2019.
- [47] Qian Yin, Qingyong Hu, Hao Liu, Feng Zhang, Yingqian Wang, Zaiping Lin, Wei An, and Yulan Guo. Detecting and tracking small and dense moving objects in satellite videos: A benchmark, 2021.
- [48] Jun Yue, Leyuan Fang, Pedram Ghamisi, Weiyang Xie, Jun Li, Jocelyn Chanussot, and Antonio J Plaza. Optical remote sensing image understanding with weak supervision: Concepts, methods, and perspectives, 2022.
- [49] Christos Kyrkou and Theodoris Theodoridis. Emergencynet: Efficient aerial image classification for drone-based emergency monitoring using atrous convolutional feature fusion, 2021.
- [50] Benedikt Blumenstiel, Viktoria Moor, Romeo Kienzler, and Thomas Brunschweiler. Multi-spectral remote sensing image retrieval using geospatial foundation models, 2024.
- [51] Anthony Fuller, Koreen Millard, and James R. Green. Transfer learning with pretrained remote sensing transformers, 2022.
- [52] Chia-Yu Hsu, Wenwen Li, and Sizhe Wang. Geospatial foundation models for image analysis: evaluating and enhancing nasa-ibm prithvi’s domain adaptability, 2024.
- [53] Gintautas Palubinskas. On the validation of pansharpening methods, 2021.

Disclaimer:

SurveyX is an AI-powered system designed to automate the generation of surveys. While it aims to produce high-quality, coherent, and comprehensive surveys with accurate citations, the final output is derived from the AI's synthesis of pre-processed materials, which may contain limitations or inaccuracies. As such, the generated content should not be used for academic publication or formal submissions and must be independently reviewed and verified. The developers of SurveyX do not assume responsibility for any errors or consequences arising from the use of the generated surveys.

www.SurveyX.cn