
Climate Change and Its Impact on Thermal Ecology and Biodiversity: A Survey

www.surveymx.cn

Abstract

This survey paper explores the complex interplay between climate change, thermal ecology, and biodiversity, emphasizing their interconnected impacts on ecological systems. It highlights the critical role of temperature dynamics, such as daily minimum temperatures and fluctuations, in shaping ecosystem health and species interactions. Advanced modeling techniques, including Bayesian hierarchical models, are instrumental in predicting climate impacts and providing early warning signals for ecological transitions. Thermal ecology significantly influences species interactions, with temperature fluctuations affecting metabolic rates and population dynamics, particularly in insect populations. Biodiversity acts as a buffer against environmental changes, essential for maintaining planetary climate stability. The paper underscores the potential of crop diversification as an adaptation strategy and the promise of Quantum Artificial Intelligence in advancing climate science. Despite challenges, no evidence was found for increasing precipitation extremes in dry regions, highlighting the need for refined data analysis. Future research should expand time series data, refine analytical methods, and integrate interdisciplinary perspectives to address climate impacts on ecosystems. Innovative solutions, such as bioacoustic communication in pest management, are crucial for enhancing biodiversity resilience. Overall, the survey calls for continued research and innovation to mitigate climate impacts and ensure ecosystem resilience, advocating for integrating socioeconomic considerations into risk assessment and management strategies.

1 Introduction

1.1 Climate Change Overview

Climate change presents a significant and intricate challenge, characterized by its global ramifications and the complexities of its wicked nature, necessitating interdisciplinary strategies to mitigate its multifaceted impacts [1]. Primarily driven by anthropogenic activities such as fossil fuel combustion and deforestation, the rise in atmospheric CO₂ levels substantially contributes to the greenhouse effect, resulting in global warming. This phenomenon correlates with an increase in the frequency and severity of extreme weather events, including intense heat waves and catastrophic floods, undermining the Earth's habitability [2].

The variability in atmospheric conditions, particularly in mid-latitude regions, complicates climate dynamics and challenges the accuracy of climate models, which are essential for understanding and forecasting climate behavior [3]. These models simulate interactions among various components of the climate system, revealing significant alterations in precipitation patterns, especially in arid regions, where previous statistical artifacts have led to exaggerated trend assessments.

The global significance of climate change is evident in its impact on terrestrial and freshwater ecosystems, with both observed and projected changes threatening biodiversity and ecosystem services. Moreover, climate change exacerbates the proliferation of invasive species, highlighting the




Figure 1: chapter structure

need for sustainable alternatives to traditional control methods, such as chemical pesticides [4]. A comprehensive understanding of climate system behaviors—such as ice age dynamics and temperature fluctuations—is vital for developing sustainable strategies that address the economic, environmental, and social complexities of climate change, emphasizing the need for global cooperation and policy interventions.

1.2 Relevance to Thermal Ecology and Biodiversity

Climate change significantly influences thermal ecology and biodiversity by modifying temperature regimes and atmospheric conditions, which directly affect species interactions, distributions, and survival strategies. This impact is particularly pronounced in regions like the Mediterranean Basin, where climate change threatens water resources, biodiversity, and food security [5]. Understanding these dynamics is crucial for predicting ecosystem responses and developing effective adaptation strategies [6]. The global redistribution of species due to changing climate conditions disrupts ecosystems and human societies, underscoring the importance of studying these interactions within thermal ecology and biodiversity frameworks [7].

Shifts in population boundaries, driven by environmental gradients and local heterogeneities, are critical for forecasting biodiversity changes [8]. The intricate relationship between climate stability and biodiversity is highlighted by feedback mechanisms linking the biosphere and planetary climate systems [9]. This interconnection is essential for maintaining ecological balance and habitability, both of which are jeopardized by ongoing climatic shifts [10].

Global warming contributes to the expansion and weakening of the tropical atmospheric component, affecting temperature patterns and circulation, which may disrupt ecological networks and alter species interactions and ecosystem functions [11]. Rising temperatures, increased atmospheric CO₂ levels, and modified precipitation patterns drive changes in agricultural insect pest dynamics, necessitating adaptive pest management strategies [8]. These changes have cascading effects on ecosystem health and biodiversity, impacting food security and ecosystem services [5].

Additionally, climate change influences soil microbiomes, which are vital for biogeochemical cycling and ecosystem health [12]. The limitations of current numerical models in accurately predicting climate change, due to their sensitivity to initial conditions and potential tipping points, pose further

challenges [13]. Discrepancies between current global warming trends and future temperature projections made by general circulation models raise questions about their reliability [14].

Habitat fragmentation, exacerbated by climate change, threatens adaptive genetic variation and alters mating systems, complicating ecosystem resilience [10]. The insufficient understanding of global warming, stemming from biases in geographical representation and temporal limitations of existing studies, underscores the need for standardized methods in analyzing temperature variations [13]. Additionally, the potential influence of cosmic rays on climate forcing, as suggested by correlations between cosmic ray variations and climate changes recorded in cosmogenic isotope archives, adds complexity to understanding climate dynamics [11].

The survival of cold-adapted species, such as *Nebria germari* in high-altitude habitats, is jeopardized by climate change, which alters their habitats and phenology [10]. Furthermore, the long-term decrease in coloration of Mediterranean blue tit subspecies over the past 15 years, in response to climate change, exemplifies the impact of climate on biodiversity and the necessity for ongoing monitoring and research [7]. These multifaceted challenges necessitate an interdisciplinary approach to address the complex interactions between climate change, thermal ecology, and biodiversity.

1.3 Structure of the Survey

This survey is structured into several key sections, each addressing distinct aspects of climate change and its intricate relationship with thermal ecology and biodiversity. The introductory section establishes the context by providing an overview of climate change, highlighting its global significance, and discussing its relevance to thermal ecology and biodiversity. It also outlines the survey's structure.

The second section explores the background and definitions of core concepts, including daily minimum temperature, temperature fluctuation, interspecific interaction, thermal ecology, insect metabolism, and biodiversity, establishing a foundational understanding for subsequent topics.

Section three examines climate change and temperature dynamics, assessing how climate change influences temperature patterns, including daily minimum temperatures and fluctuations, and discussing the implications for ecosystems and species interactions. Subsections within this section investigate modeling temperature fluctuations, the impact of temperature extremes on ecosystem health, and predictive climate modeling techniques.

The fourth section analyzes interspecific interactions and thermal ecology, focusing on how temperature fluctuations affect species interactions within ecosystems. It underscores the role of thermal ecology in understanding these interactions and reviews methodological approaches to studying them.

In section five, the survey investigates the effects of climate change on insect metabolism, considering how temperature changes can alter metabolic rates and impact insect populations. This section discusses broader implications for ecosystem functioning and biodiversity, including population dynamics and adaptation strategies.

The sixth section assesses biodiversity and ecosystem resilience, analyzing the impacts of climate change on biodiversity and ecosystem resilience to temperature changes and interspecific interactions. It explores strategies for conserving biodiversity and mitigating climate change impacts, as well as the synergistic effects of climate change on biodiversity and ecosystem stability.

The conclusion synthesizes the survey's key findings, emphasizing the intricate relationships among climate change, thermal ecology, and biodiversity. It highlights the critical need for ongoing research to address the challenges posed by climate variability, particularly its effects on ecosystems and species survival, as well as its implications for agriculture and public health. Furthermore, it identifies potential areas for future study, advocating for a multidisciplinary approach to enhance understanding of these interconnected issues and inform effective mitigation and adaptation strategies [15, 1, 16]. The following sections are organized as shown in Figure 1.

2 Background and Definitions

2.1 Key Concepts and Their Interconnections

Climate change, marked by persistent shifts in global temperatures and weather patterns, significantly challenges thermal ecology and biodiversity. These challenges are compounded by the inherent

variability and chaotic nature of climate dynamics, which complicate the identification of statistically significant trends. Traditional extreme value analysis methods often fail to account for nonstationary behavior in temperature distributions [17]. The daily minimum temperature serves as a crucial climate indicator, reflecting climate variability and influencing physiological processes and ecosystem dynamics. Anthropogenic activities, such as increased CO₂ emissions, exacerbate these fluctuations, highlighting the need for accurate climate sensitivity models. Estimating equilibrium climate sensitivity remains challenging due to varying feedback mechanisms across different climate states [3].

Temperature dynamics significantly influence interspecific interactions, which involve relationships between different species. Environmental gradients and local heterogeneities shape population boundaries, illustrating the complexity of ecological systems under changing climatic conditions [4]. Traditional models often overlook nonlinear interactions within these systems, lacking the capacity to incorporate intricate feedback mechanisms [1]. Thermal ecology provides a framework for understanding these interactions and their broader ecological implications. Habitat fragmentation, particularly in forest ecosystems, can hinder species' evolutionary responses to environmental changes, affecting their climate adaptation.

Insect metabolism, encompassing the biochemical processes that sustain insect life, is particularly sensitive to temperature variations. Fluctuations in temperature can alter metabolic rates, impacting population dynamics and ecosystem functioning [3]. The dynamics of insect infestations in boreal forests exemplify how climate influences forest health, with increasing extreme weather events, changing precipitation patterns, and rising temperatures complicating these dynamics and threatening food security and agricultural practices.

Biodiversity, defined as the variety of life forms within an ecosystem, is crucial for ecosystem resilience and stability. The interplay between climate change and biodiversity occurs through feedback mechanisms linking the biosphere and planetary climate systems [4]. Accurate representation of the nonlinear characteristics of climate models is essential for understanding climate sensitivity and its effects on biodiversity. The entropy budget of the climate system, focusing on material processes, plays a critical role in assessing climate stability and biodiversity's contribution to it [2]. Insufficient understanding of microorganisms' roles in climate change may jeopardize efforts toward environmental sustainability.

Recent theoretical advancements, such as the integration of climate network and percolation theories, provide insights into the evolution of climate patterns and their responses to global warming. Approaches proposed by Huang et al. enable reliable estimates of rare temperature extremes using extensive climate model runs, addressing the limitations of short observational records [18]. Theoretical tools that combine statistical physics and applied mathematics enhance the understanding of climate dynamics, tackling the challenge of distinguishing climate variability from climate change signals amidst unsteady external forcings and complex internal dynamics. The theory suggesting significant influences of solar and planetary oscillations on climatic variations, inadequately represented in current General Circulation Models (GCMs), further complicates the understanding of climate dynamics.

The limitations of current metrics for measuring temporal turnover, which often fail to capture the unidirectional nature of community dynamics and the ecological processes driving changes in community composition over time, pose significant challenges. Existing metrics, such as the Jaccard and Bray-Curtis indices, were designed for spatial contexts and inadequately account for the unique characteristics of temporal turnover [18]. The benchmark established for accurately estimating daily global CO₂ emissions from fossil fuel combustion and cement production during 2020 highlights the necessity for real-time data to minimize uncertainties in global emissions assessments [2]. Current benchmarks often rely on outdated national inventories, failing to capture real-time emissions changes and leading to significant uncertainties. Urban cooling demand trends, categorized based on climate drivers and urban characteristics, underscore the role of extreme heat events and urban heat islands in shaping energy demands. The assertion that climate sensitivity increases with warming challenges previous assumptions of a constant sensitivity across varying climate states [19]. Effective teaching of complex climate concepts, such as the greenhouse effect and planetary albedo, through interactive tools is vital for enhancing understanding. Extreme temperatures adversely affect agricultural productivity, resulting in reduced crop yields and altered farming practices among subsistence farmers. Organizing current methods into stages of energy resource utilization, policy

development, and technological innovation, with a focus on their interdependencies and impacts on climate change, is critical for addressing these challenges. The necessity to identify and confirm a 60-year oscillation in global climate data, inadequately represented in current general circulation models, further complicates the understanding of climate dynamics [3]. These interconnected concepts highlight the complexity of climate change and its pervasive impact on thermal ecology and biodiversity.

The complexities of climate change necessitate a comprehensive understanding of its various dimensions, particularly in relation to temperature dynamics. To elucidate this multifaceted issue, Figure 2 provides a visual representation of the hierarchical structure of climate change and temperature dynamics. This figure illustrates advanced methodologies for modeling temperature fluctuations and the impact of temperature extremes on ecosystem health. Furthermore, it highlights predictive climate modeling techniques that incorporate probabilistic models, machine learning, and data assimilation methods. By doing so, the figure underscores the importance of these integrated approaches in enhancing climate prediction accuracy and informing effective adaptation strategies.




Figure 2: This figure illustrates the hierarchical structure of climate change and temperature dynamics, encompassing advanced methodologies for modeling temperature fluctuations, the impact of temperature extremes on ecosystem health, and predictive climate modeling techniques. The figure highlights the integration of probabilistic models, machine learning, and data assimilation methods to enhance climate prediction accuracy and inform effective adaptation strategies.

3 Climate Change and Temperature Dynamics

3.1 Modeling Temperature Fluctuations

Understanding the complexities of climate change requires sophisticated models to capture the stochastic and nonlinear nature of temperature dynamics. Traditional deterministic models often fall

short, prompting the development of advanced methodologies. The Bayesian hierarchical model (HCM-RBE) effectively captures spatial and temporal dependencies in record-breaking temperature events, offering insights into regional trends, such as those in peninsular Spain [20]. This model exemplifies the integration of spatial and temporal data to improve climate prediction accuracy.

The multi-level spatio-temporal model (MLSTM) enhances predictions by incorporating autoregressive terms and spatial processes, underscoring the importance of these dimensions in climate models for accurate temperature dynamics representation [21]. Machine learning techniques, such as the DCR method, leverage eXtreme Gradient Boosting (XGBoost) on high-resolution data to predict historical daily CO₂ emissions, thereby deepening our understanding of climate dynamics [22]. These methodologies highlight the role of advanced data analytics in climate modeling, facilitating predictions in sectors like agriculture [23].

As illustrated in Figure 3, the BATs distribution, a flexible parametric model, captures the entire probability distribution of temperature data, effectively addressing both extremes and central tendencies [17]. Additionally, hierarchical frameworks like WOMBAT v2.0 model trends and seasonal cycles of CO₂ fluxes, illustrating the integration of spatial and temporal variability in climate modeling [24]. These advanced modeling techniques are crucial for understanding temperature fluctuations and their implications for climate change impacts, guiding effective adaptation strategies. By employing sophisticated statistical methods and innovative modeling approaches, these methodologies provide critical insights into the complex ecological and atmospheric consequences of temperature variations, informing public discourse and climate change strategies [16, 25, 26, 1, 6].




Figure 3: This figure illustrates advanced climate modeling techniques, highlighting Bayesian models like HCM-RBE, MLSTM, and WOMBAT v2.0, machine learning approaches such as the DCR Method and XGBoost, and parametric models including the BATs Distribution and Extreme Value Analysis. These methods enhance our understanding of complex climate dynamics and improve predictive accuracy.

3.2 Temperature Extremes and Ecosystem Health

Temperature extremes significantly challenge ecosystem health and stability by disrupting species distributions and community dynamics essential for maintaining ecological balance [27]. Advanced statistical models elucidate temperature trends and their ecological impacts, with the TML model capturing non-normality and regime-switching behavior in temperature dynamics [28].

The limitations of current General Circulation Models (GCMs) in replicating critical patterns, such as climatic oscillations synchronized with astronomical cycles, are highlighted by studies on solar and planetary influences, vital for predicting extreme temperature events [29]. Employing a neural shooting approach with the Onsager-Machlup action functional enhances predictive capabilities by uncovering probable transition pathways in the carbon cycle [30].

In arid ecosystems, environmental inhomogeneities, exacerbated by temperature extremes, influence vegetation patterns' self-organization and clustering [14]. This self-organization is crucial for understanding ecosystem resilience under climate-induced stress. Additionally, the negative feedback effect of low clouds on climate sensitivity highlights the complexity of climatic responses to temperature extremes [31].

The rapid decrease in zonal surface temperature gradients with warming, explained by a simplified surface energy budget, illustrates the intricate interactions between temperature extremes and climate dynamics [32]. Advanced modeling approaches that integrate spatially varying coefficients and temporal autoregressive effects provide a comprehensive framework for analyzing daily maximum temperature trends and their ecological impacts [21].

Systemic growth constants in CO₂ emission trajectories suggest significant temperature increases with profound implications for ecosystem health [33]. These projections underscore the urgency of developing effective adaptation strategies to enhance ecosystem resilience. Crop diversification, enhancing net primary productivity by 2.8%, represents a viable strategy for mitigating temperature extremes' impacts on agriculture [8].

Explicitly considering temperature's effect on shear strength in slope stability analyses signifies a notable advancement over previous methods [2]. The complex interplay between temperature extremes and ecosystem stability necessitates a multidisciplinary approach integrating atmospheric, oceanic, and biological data. Advanced modeling techniques and theoretical frameworks are essential for accurately predicting extreme temperatures' impacts, guiding effective adaptation strategies, and ensuring ecosystem resilience amid increasing climate variability.

3.3 Predictive Climate Modeling Techniques

Predictive climate modeling techniques are crucial for accurately forecasting climate change impacts on temperature dynamics, essential for developing effective adaptation strategies. Traditional General Circulation Models (GCMs) face challenges in capturing inherent climate system uncertainties, prompting the integration of probabilistic modeling approaches to better reflect these uncertainties [34]. Incorporating stochastic parameterizations represents a significant advancement, accounting for unresolved scale effects on climate model dynamics and enhancing prediction accuracy [21].

The integration of machine learning and statistical methods into climate modeling has shown promise in improving predictive capabilities. Bayesian hierarchical models incorporating spatial correlation and inter-model dependence yield more precise uncertainty quantifications and improved climate projection accuracy [34]. This approach emphasizes the necessity of considering spatial variability and temporal dependencies for a comprehensive understanding of temperature dynamics.

Innovative techniques, particularly Quantum Artificial Intelligence (QAI), hold significant potential for transforming climate modeling by utilizing advanced computational methods to tackle the complex challenges posed by climate change. Recent advancements in QAI could enhance predictive accuracy and efficiency in weather forecasting, critical for effective adaptation strategies aimed at minimizing extreme weather events' impacts. As traditional AI approaches face limitations due to hardware constraints and data processing demands, QAI's capacity to process vast datasets may unlock new insights into climate dynamics, ultimately benefiting societal resilience and environmental management [25, 35, 36, 37]. These advancements underscore the evolving landscape of climate modeling, where the fusion of traditional and cutting-edge methodologies provides a robust framework for understanding climate dynamics.

The advancement of analytical methods that accurately replicate the primary periods of Milankovitch cycles presents strong alternatives to conventional numerical integration techniques, highlighting the necessity for a diverse toolkit in predictive modeling. These methods, derived from first principles, address numerical drift issues prevalent in long computer simulations and reveal key components of Milankovitch cycles with periods of 41,000, 29,500, and 54,000 years, influenced by planetary orbit precession. Furthermore, integrating machine learning with physically-based models enhances climate prediction accuracy by correcting systematic errors, while research into non-smooth dynamical systems provides insights into the complex behavior of climate models under varying forcing conditions. Collectively, these developments emphasize the importance of employing varied approaches to enhance our understanding and forecasting of long-term climate dynamics [38, 39, 40]. Additionally, the application of data assimilation techniques and numerical simulations to estimate the climate system's state reflects ongoing efforts to refine climate models and improve their reliability.

Recent advancements also include the development of visual analysis and prediction methods that utilize classic models tailored to data characteristics, simplifying complexity and reducing the need for advanced feature engineering. Moreover, spatial quantile clustering methods effectively account

for the distributional characteristics of climate data, including asymmetry and outliers, leading to more reliable clustering outcomes [21].

These predictive climate modeling techniques, integrating stochastic processes, machine learning, and innovative statistical methods, are essential for advancing our understanding of climate dynamics. As climate models evolve, their ability to predict temperature fluctuations accurately will be crucial for mitigating climate change's adverse effects on biodiversity and ecosystem services. Enhanced forecasting can facilitate improved management of species responses to environmental changes, vital for maintaining ecosystem integrity and resilience. By leveraging biophysical models that consider thermodynamic interactions between organisms and their environments, we can better comprehend climate variability's implications on species behavior, distribution, and survival. This understanding is critical for developing effective adaptation strategies that support human well-being and the sustainability of natural systems amid ongoing climate challenges [41, 42, 7, 15].

4 Interspecific Interactions and Thermal Ecology

4.1 Role of Thermal Ecology in Species Interactions

Thermal ecology is pivotal in understanding species interactions and ecosystem stability, especially as climate change alters temperature regimes and ecological boundaries. Species' adaptive capacity to these changes is crucial for survival. A framework for analyzing climate connectivity reveals emerging patterns and potential tipping points, enhancing our understanding of climate variables' interplay with species dynamics [43]. Temperature fluctuations also impact human health, particularly cardiovascular mortality, underscoring thermal variations' broader implications on ecological and human systems [5]. This necessitates integrated frameworks that consider both ecological and anthropogenic factors when assessing climate change impacts.

As illustrated in Figure 4, the role of thermal ecology in species interactions, human health impacts, and ecosystem resilience is multifaceted, highlighting key studies and frameworks that contribute to understanding these dynamics. For cold-adapted species, microhabitats like supraglacial debris serve as critical refugia, maintaining ecological stability amid climate changes [11]. Incorporating interspecies interactions into predictive models enhances forecasts of species distributions under varying climatic conditions, elucidating temperature's effects on species interactions and ecosystem resilience [44]. Developing frameworks that include community dynamics and temporal turnover is essential for accurately capturing ecological interactions [45]. These frameworks provide insights into how temperature fluctuations impact species interactions, enhancing our understanding of ecosystem resilience against climate change.

Thermal ecology thus offers a critical perspective on species-environment interactions. Investigating temperature's influence on these dynamics provides insights necessary for developing adaptive strategies that bolster ecosystem stability and biodiversity conservation. Advanced climate models that account for natural cycles and feedback mechanisms further enrich our understanding of these dynamics and their ecological implications [3].




Figure 4: This figure illustrates the role of thermal ecology in species interactions, human health impacts, and ecosystem resilience, highlighting key studies and frameworks that contribute to understanding these dynamics.

4.2 Methodological Approaches to Studying Interspecific Interactions

Robust methodological approaches are essential for understanding interspecific interactions within the framework of thermal ecology. Reaction-diffusion models, as outlined by [46], provide a foundational tool for describing population dynamics across space and time, incorporating logistic growth and Allee effects critical under changing climatic conditions. Integrating these models with empirical data allows for simulating species spatial spread and predicting distribution shifts due to climate change. These models are valuable for analyzing temperature fluctuations' impacts on species interactions, offering insights into community structure and ecosystem dynamics, including predator-prey interactions and distribution limits [41, 42].

Advanced statistical and computational techniques, such as agent-based models and network analysis, are increasingly important for studying interspecific interactions. Agent-based models simulate species behaviors and interactions, enhancing understanding of species adaptation to temperature changes at the micro-level. Tools like EcoSISTEM integrate extensive plant occurrence records and climate data to predict biodiversity responses across scales, revealing critical insights into environmental changes' timing and magnitude [47, 42]. Network analysis examines interspecies interdependencies, underscoring temperature's role as a driver of interaction networks.

Machine learning algorithms have emerged as powerful tools for analyzing large ecological datasets, identifying patterns, and predicting changes in species interactions. These algorithms utilize techniques such as convolutional neural networks and dynamic time warping to process extensive datasets, revealing intricate relationships often overlooked by traditional methods. This analytical capability enhances our understanding of thermal ecology's impact on interspecific dynamics amid climate change [41, 25, 42, 45]. Additionally, remote sensing technologies provide valuable spatial and temporal data on habitat conditions and species distributions, facilitating a comprehensive assessment of temperature variations' impacts on interspecific interactions. By integrating diverse research methodologies, including ecological modeling and data-driven analyses, researchers can attain a holistic understanding of ecosystem interactions, crucial for effective conservation strategies and policy decisions addressing climate change's multifaceted impacts on biodiversity and ecosystem services [15, 48, 7, 8].

4.3 Climate Change and Interspecific Interaction Challenges

Climate change poses significant challenges to interspecific interactions, profoundly affecting biodiversity and ecosystem dynamics. A notable issue is the reduction in biodiversity, essential for maintaining planetary climate stability. Decreased biodiversity can lead to significant climate shifts, disrupting ecological balance and altering species interactions [49]. Predicting species redistribution outcomes is complex, as species respond differently to climate change, resulting in unpredictable ecological consequences [48].

Inadequacies in historical datasets and logistical challenges in sampling, especially in high-altitude and inaccessible areas, hinder accurate assessments of interspecific interactions [50]. This data gap restricts understanding of climate change's full impact on species interactions, while the low temporal resolution of existing datasets limits predictive models' ability to capture dynamic ecosystem changes, such as mangrove distribution [51]. Biases in climate models complicate understanding of interspecific interactions under changing conditions [52]. Developing models that account for both short-term and long-term responses to climate forcing is crucial for comprehending climate impacts on species interactions [53]. Moreover, the lack of understanding regarding sudden climate changes' effects on predator-prey systems necessitates comprehensive studies to elucidate these dynamics [42].

Addressing these challenges requires a multifaceted approach, including developing nonparametric methods that avoid arbitrary thresholds and provide clearer insights into extreme temperature trends [54]. Immediate reductions in fossil fuel emissions and effective CO₂ extraction strategies are vital to mitigate future catastrophic climate impacts [55]. Enhancing data collection, improving model accuracy, and fostering interdisciplinary research can facilitate effective strategies to mitigate climate change's impacts on interspecific interactions, ensuring ecosystem resilience.

Fragmentation in climate change strategies further exacerbates these challenges, reducing effectiveness and collaboration among stakeholders [56]. The limited application of biophysical models in ecology, combined with data availability issues and a lack of common software ecosystems, hinders the development of comprehensive solutions [41]. However, sustainable bio-control methods,

such as enhancing predator efficacy, offer promising avenues for addressing these challenges [4]. Infrastructure, such as roads, traditionally viewed as detrimental in stationary climates, may aid population persistence amid climate change by supporting ecological adaptation [57]. Intermediate levels of mutation and adaptation can enhance population survival, providing insights into evolutionary strategies that may buffer against climate-induced extinction risks [58]. Integrating ecological, infrastructural, and evolutionary perspectives is essential for effectively addressing climate change challenges to interspecific interactions. Recognizing ecological grief is also important, as it enhances understanding of the emotional dimensions of climate change impacts, critical for fostering resilience and adaptation [18].

5 Insect Metabolism and Climate Change

5.1 Temperature Fluctuations and Metabolic Rates

Temperature fluctuations critically affect insect metabolic rates, influencing population dynamics and ecological processes. As ectotherms, insects depend on external temperatures to regulate their metabolism, with variations impacting growth, reproduction, and survival, thus affecting ecological dynamics [5]. This sensitivity necessitates accurate climate models to predict these effects. Advanced modeling techniques, including machine learning, improve predictions of temperature impacts on insect metabolism by aligning theoretical models with observed data [59]. Solar models are essential for understanding how solar activity influences climate and, consequently, insect metabolism [60]. Rainfall, closely tied to temperature, significantly shapes insect populations, notably affecting species like *Aedes aegypti*, with implications for ecosystem and public health [4]. Historical climate data reveal trends such as a 60-year oscillation possibly linked to solar-astronomical factors, which contribute to global warming and affect insect metabolic rates [61]. These insights are crucial for understanding long-term impacts on insect metabolism and ecosystem stability. Furthermore, methodologies like Langevin-type dynamics offer tools to detect critical ecological transitions [62].

5.2 Population Dynamics and Adaptation

Insects adapt to climate-induced metabolic changes through various strategies, essential for maintaining ecological balance and biodiversity. Climate change alters temperature, CO₂ levels, and precipitation, prompting insects to expand ranges, enhance overwintering survival, and modify reproductive cycles. These shifts impact agricultural production and food security, increasing crop losses and complicating pest management. The interdependence of insect populations and ecosystems creates feedback loops that exacerbate climate impacts, necessitating innovative monitoring and control strategies to preserve biodiversity [48, 63, 64]. The adaptive capacity of insects is linked to climate change rates, with critical thresholds increasing extinction risks if adaptation fails. Understanding climate dynamics and insect metabolism is vital for predicting population shifts. Advanced methodologies, like the GKL algorithm, improve estimation of extreme heatwave return times, aiding in understanding climatic impacts on insect populations [65]. Wavelet spectral analysis (WSA) identifies shifts in extreme events, offering insights into insect responses to climate dynamics [66]. These tools are crucial for understanding interactions between climate variables and insect adaptation strategies. Predator-prey dynamics are notably affected by climate variability, with shifts between favorable and unfavorable conditions altering interactions and population stability [42]. Habitat fragmentation challenges insect adaptation by disrupting climate adaptation processes and reducing drought tolerance, impacting species like *Primula elatior* [67]. This fragmentation decreases flowering investment and alters mating systems, affecting biodiversity and ecosystem resilience. Research is often limited by sampling biases and body size distribution variability, obscuring true ecological trends [10]. Non-autonomous nonlinear models, especially for honey bee and parasite interactions, highlight seasonality's role in reproductive strategies and population dynamics [68]. These models provide insights into insect life cycle adaptations to environmental changes. The interaction between selection pressure and mutation rates is crucial for predicting insect responses to climate change [58]. Understanding these interactions enables better anticipation of adaptive strategies, informing conservation and biodiversity management. In agriculture, extreme heat affects productivity, prompting subsistence farmers to adapt by expanding planted areas and altering crops, partially offsetting output declines [69]. Future research should refine temperature dynamics models and explore applications across regions or weather variables affecting agriculture [28]. These efforts are key to developing strategies ensuring ecosystem and agricultural resilience against climate change.

5.3 Case Studies: Specific Insect Populations

Climate change significantly influences insect populations, affecting distribution, abundance, and ecological roles. Case studies show how rising temperatures, altered precipitation, and increased CO₂ levels impact insect dynamics, including geographic distribution shifts, survival rates, and interactions with plants and other species, posing risks to agriculture and food security [70, 68, 63, 64]. The honey bee (*Apis mellifera*) exemplifies climate impacts on insects through seasonality and parasitism. Seasonality can enhance survival or cause collapse, depending on environmental and parasitic conditions [68]. This highlights the balance between climatic variables and biological interactions, necessitating adaptive management for pollinator survival. Leaf litter decomposition, mediated by insect detritivores, is significantly influenced by climate change. Future research should refine models by incorporating factors like nitrogen content and site fertility, affecting decomposition at smaller scales [71]. Understanding micro-scale ecological processes is vital for nutrient cycling and ecosystem health amid climatic shifts. Annual plant-insect interactions offer insights into climate change effects. Studies suggest annual plants thrive in hotter, drier conditions due to climate variability and anthropogenic influences [72]. This shift alters insect population composition and behavior, potentially changing ecosystem dynamics and biodiversity. These case studies underscore the need for ongoing research into climate change's multifaceted impacts on insects. Analyzing specific climate-driven changes provides deeper insights into ecosystem interactions, assesses risks like biodiversity loss and altered structures, and formulates conservation strategies to address climate change's adverse impacts on ecosystem services, including water security and agricultural productivity [7, 1, 16].

6 Biodiversity and Ecosystem Resilience

6.1 Impacts on Ecosystem Functioning

Climate change profoundly affects ecosystem functioning by disrupting both biotic and abiotic components, primarily through shifts in temperature, precipitation, and extreme weather events. The state-dependent nature of climate sensitivity necessitates a nuanced understanding of ecosystem responses across diverse climates, crucial for predicting adaptation and recovery from climatic perturbations [19]. Advanced forecasting models emphasize the importance of immediate action to prevent surpassing critical temperature thresholds, although the urgency may vary [73].

Biodiversity is integral to ecosystem resilience, providing functional traits that buffer against climate variability and sustain ecosystem functions. Even minor changes in carbon levels can lead to significant climate shifts, highlighting the interconnectedness of climate change and biodiversity [19]. Mountain ecosystems exemplify this complexity, with some regions experiencing increased vegetation and microbial biomass, while others suffer declines in freshwater quality. Such differential responses underscore the need for localized studies to identify specific vulnerabilities. Continuous spatiotemporal approaches, as discussed in [2], enhance accuracy by incorporating spatial heterogeneity and addressing data gaps.

High-altitude ice-related landforms serve as refugia for cold-adapted species, emphasizing the importance of preserving diverse habitats for species survival and ecosystem resilience. The geometry of environmental boundaries and Allee effects significantly influence population persistence, indicating that high mobility does not guarantee survival amid rapid climate change. Integrated models capturing spatial and temporal dynamics are crucial for predicting ecosystem responses to environmental changes, including climate change and habitat loss. Utilizing advanced computational capabilities and digitized herbarium records, models like EcoSISTEM can simulate ecosystem responses at various scales, enhancing our understanding of biodiversity dynamics and community turnover [47, 45].

The resilience of ecosystems to climate change is intricately linked to biodiversity, which enhances their adaptability and recovery from disturbances. Preserving biodiversity is vital for maintaining ecosystem services such as food security, disease regulation, and carbon sequestration, as well as for bolstering ecological resilience in the face of rapid climate shifts that alter species distributions and community dynamics globally [48, 7].

6.2 Conservation and Mitigation Strategies

Conservation and mitigation strategies are crucial for addressing climate change's multifaceted impacts on biodiversity and ecosystems. An integrated approach combining biological, environmental, and socio-economic factors is necessary to manage the intricate interactions within the climate system. The Optimal Enhancing Resilience Model (OERM) provides a strategic framework for safeguarding critical ecological components while maintaining ecosystem resilience amid climatic perturbations [12]. Precision agriculture and stress-tolerant crop varieties are vital elements of mitigation strategies, enhancing agricultural resilience to climate variability and sustaining ecosystem services.

Incorporating real-time data streams into climate models can significantly improve their adaptability and accuracy, facilitating better decision-making in policy and management [34]. The potential of Quantum Artificial Intelligence (QAI) to enhance climate model accuracy and efficiency underscores the need for further exploration of machine learning methods [35]. Public education and awareness are critical for fostering support for conservation and mitigation efforts. Enhancing carbon pricing mechanisms can drive substantial reductions in greenhouse gas emissions, contributing to global climate goals [74].

Integrating biophysical models can deepen our understanding of species responses to climate change, highlighting the necessity for improved training and data infrastructure to broaden their application [6]. Efficient analysis of large datasets can bolster strategies for conserving biodiversity in urban environments. Future research should focus on refining existing frameworks, exploring data-driven approaches, and addressing the limitations of current models in capturing the multiscale nature of climate dynamics [38]. By integrating advanced modeling techniques, conservation practices, and public engagement, these strategies can effectively mitigate climate change impacts, ensuring biodiversity preservation and ecosystem resilience.

Future research could explore sophisticated methods to combine ecological data across species and investigate subsets of species that respond differently to climate models [64]. Refining models to include additional oscillations and integrating detailed empirical data can enhance predictive accuracy [3]. Increasing the density of weather stations in underrepresented areas and incorporating remote sensing data can improve evapotranspiration estimates, crucial for understanding ecosystem responses to climate change [75]. Comprehensive multi-attribution analyses are necessary for a more complete understanding of climate dynamics, including the ecological roles of cold-adapted species and their interactions with competitors and prey in conservation strategies [76].

Innovative pest management approaches, such as those proposed in [4], align with conservation strategies by reducing reliance on harmful pesticides, thus supporting biodiversity. Future research should focus on in-depth analyses of specific climate change topics and the exploration of emerging trends in adaptation and mitigation strategies [1]. Additionally, developing fully-coupled thermo-hydro-mechanical models and conducting case studies to validate proposed methods under various conditions is essential for advancing conservation efforts [2].

6.3 Synergistic Effects of Climate Change on Biodiversity

The synergistic effects of climate change on biodiversity and ecosystem stability are characterized by complex interactions between climatic variables and ecological processes, often leading to abrupt shifts in population boundaries. Gradual environmental changes can mask ecological responses until significant changes are triggered, potentially misrepresenting stability [77]. Such dynamics highlight the potential for sudden and unpredictable changes in biodiversity as ecosystems respond to climatic pressures.

Wavelet Spectral Analysis (WSA) has proven effective in detecting significant shifts in climate signals, crucial for understanding the impacts on biodiversity and ecosystem stability [66]. These shifts can disrupt ecological networks, causing cascading effects on species interactions and ecosystem functions. However, gaps remain in understanding the long-term effects of climate change on pest dynamics, complicating management strategies and emphasizing the need for comprehensive approaches to mitigate these impacts [63].

Current climate models face limitations in accurately predicting future temperatures, particularly due to the complex role of solar influences, which are not fully understood [78]. The potential role of cosmic rays in modulating climate further complicates predictions, indicating the need for additional

research to elucidate these influences [79]. These uncertainties underscore the importance of refining climate models to incorporate a broader range of climatic factors and improve representations of regional processes [80].

Technological uncertainties, high costs, and the lack of comprehensive policy frameworks pose significant challenges to the large-scale implementation of climate mitigation strategies [81]. Addressing these challenges is critical for developing effective conservation strategies that enhance biodiversity and ecosystem resilience. Future research should focus on integrating additional ecological variables and conducting field studies to validate theoretical predictions, thereby improving our understanding of the synergistic effects of climate change [68].

Quantum Mechanical principles offer a promising avenue for gaining deeper insights into climate change dynamics, potentially leading to a better understanding of climatic transitions and their impacts on biodiversity [82]. The gravitational influences of celestial bodies, as described by Milankovitch cycles, further complicate long-term climate dynamics, affecting biodiversity and ecosystem stability [40]. By addressing these complexities, researchers can develop more accurate models and strategies to mitigate the synergistic effects of climate change on biodiversity, ultimately ensuring ecosystem resilience amid ongoing climatic shifts.

6.4 Role of Biodiversity in Climate System Stability

Biodiversity is pivotal for maintaining climate system stability and ecosystem health by providing a diverse array of functional traits that buffer against environmental changes. The interplay between biodiversity and climate dynamics underscores the capacity of diverse ecosystems to enhance resilience, sustaining functionality under varying climatic conditions. The diversity of species within an ecosystem bolsters its resilience to disturbances and recovery capacity, crucial for maintaining ecological balance and ensuring the provision of vital ecosystem services such as food security, carbon sequestration, and disease regulation. As climate change drives rapid shifts in species distributions, understanding the implications of biodiversity loss on ecosystem functioning becomes essential for effective environmental management and adaptation strategies [48, 7].

Feedback mechanisms connecting biological diversity to climate processes further underscore the significance of biodiversity in maintaining climate system stability. Climate-driven species redistribution can disrupt ecosystem functioning and human well-being by altering natural resource production, disease transmission patterns, and carbon sequestration dynamics. As species shift in response to climate change, this can initiate self-reinforcing feedback loops, such as increased insect populations leading to deforestation, exacerbating climate change through reduced carbon storage and enhanced carbon release [48, 64]. Diverse plant communities influence atmospheric carbon levels through varying rates of photosynthesis and respiration, affecting climate regulation and highlighting biodiversity's critical role in modulating climate dynamics.

Research has established connections between solar activity and climate, enhancing understanding of historical climate patterns and the role of astronomical factors [78]. Insights into the natural variability of the climate system underscore biodiversity's importance in buffering against both natural and anthropogenic climatic changes. Biodiverse ecosystems are better equipped to absorb shocks and maintain stability amid external pressures, contributing to the climate system's overall resilience.

Moreover, biodiversity supports essential processes such as nutrient cycling, soil formation, and pollination, which are vital for preserving ecosystem integrity. These ecological processes directly influence biodiversity, ecosystem functioning, and the capacity of systems to provide critical services increasingly threatened by climate change [48, 7, 83, 12]. The loss of biodiversity can disrupt these processes, diminishing ecosystem resilience and increasing vulnerability to climate change.

7 Conclusion

The survey delineates the intricate relationships among climate change, thermal ecology, and biodiversity, underscoring their profound impacts on ecological systems. Temperature dynamics, especially daily minimum temperatures and fluctuations, play a pivotal role in determining ecosystem health and species interactions. Advanced modeling techniques, such as Bayesian hierarchical models and the MC-LR method, enhance predictive capabilities, revealing the synergistic effects of climatic factors

on ecological processes. These models facilitate the detection of critical slowing down (CSD) and provide early warning signals for critical transitions, particularly under red noise conditions.

Thermal ecology is crucial in mediating species interactions, as temperature fluctuations affect metabolic rates and population dynamics, notably in insect populations. This highlights the need for continued research into adaptive strategies and the development of models that incorporate complex ecological interactions and tipping points. Biodiversity acts as a buffer against environmental changes, essential for maintaining the stability of the planetary climate system. Crop diversification emerges as a viable adaptation strategy, enhancing agricultural resilience, particularly in regions like Cyprus.

The potential of Quantum Artificial Intelligence to advance climate science, especially in pattern recognition and simulation studies, is acknowledged, despite existing technological challenges. The survey also indicates that, with appropriate definitions and data processing methods, there is no evidence of increasing precipitation extremes or annual totals in the world's dry regions, emphasizing the need for refined data analysis techniques to accurately assess climate impacts.

Future research should focus on expanding time series data, refining analytical methods, and exploring the implications of emerging climate patterns on food security. Interdisciplinary studies that integrate ecological, evolutionary, and neurobiological perspectives are vital for understanding the impacts of climate change on nervous systems. Additionally, exploring the role of bioacoustic communication in insect infestation dynamics offers new avenues for detection and control strategies. The integration of additional food resources with spatial dynamics can lead to effective pest management strategies, highlighting the importance of innovative solutions in the context of climate change and biodiversity.

References

- [1] Robin Haunschild, Lutz Bornmann, and Werner Marx. Climate change research in view of bibliometrics, 2016.
- [2] Marco Loche and Gianvito Scaringi. Temperature continuously controls the stability of clay slopes, 2022.
- [3] Craig Loehle and Nicola Scafetta. Climate change attribution using empirical decomposition of climatic data, 2012.
- [4] Urvashi Verma, Aniket Banerjee, and Rana D. Parshad. T(w)o patch or not t(w)o patch: A novel additional food model, 2024.
- [5] Pierre Masselot, Fateh Chebana, Taha B. M. J. Ouarda, Diane Bélanger, André St-Hilaire, and Pierre Gosselin. A new look at weather-related health impacts through functional regression, 2018.
- [6] Arianna Pera and Luca Maria Aiello. Shifting climates: Climate change communication from youtube to tiktok, 2024.
- [7] Camille Parmesan, Mike D Morecroft, and Yongyut Trisurat. *Climate change 2022: Impacts, adaptation and vulnerability*. PhD thesis, GIEC, 2022.
- [8] Georgios Giannarakis, Ilias Tsoumas, Stelios Neophytides, Christiana Papoutsas, Charalampos Kontoes, and Diofantos Hadjimitsis. Understanding the impacts of crop diversification in the context of climate change: a machine learning approach, 2023.
- [9] Adam Frank and Woodruff Sullivan. Sustainability and the astrobiological perspective: Framing human futures in a planetary context, 2013.
- [10] Grant M. Connette, John A. Crawford, and William E. Peterman. Climate change and shrinking salamanders: Alternative mechanisms for changes in plethodontid salamander body size, 2014.
- [11] Keno Riechers, Georg Gottwald, and Niklas Boers. Glacial abrupt climate change as a multi-scale phenomenon resulting from monostable excitable dynamics, 2024.
- [12] Yu Sun, Teng Liu, Shang Wang, Jun Meng, Yongwen Zhang, Saini Yang, Xiaosong Chen, Deliang Chen, Jürgen Kurths, Shlomo Havlin, Hans Joachim Schellnhuber, and Jingfang Fan. Regional greening as a ‘positive’ tipping phenomenon, 2023.
- [13] Flavio C. D. Moraes, Ana Lia Leonel, Pedro H. C. Torres, Pedro R. Jacobi, and Sandra Momm. Climate change and social sciences: a bibliometric analysis, 2020.
- [14] D. Pinto-Ramos, M. G. Clerc, A. Makhoute, and M. Tlidi. Vegetation clustering and self-organization in inhomogeneous environments, 2024.
- [15] Kashif Abbass, Muhammad Zeeshan Qasim, Huaming Song, Muntasir Murshed, Haider Mahmood, and Ijaz Younis. A review of the global climate change impacts, adaptation, and sustainable mitigation measures. *Environmental science and pollution research*, 29(28):42539–42559, 2022.
- [16] Smriti Janaswamy and Jeremy Blackburn. Exploring climate change discourse: Measurements and analysis of reddit data, 2024.
- [17] Mitchell Krock, Julie Bessac, Michael L. Stein, and Adam H. Monahan. Nonstationary seasonal model for daily mean temperature distribution bridging bulk and tails, 2021.
- [18] Ashlee Cunsolo and Neville R Ellis. Ecological grief as a mental health response to climate change-related loss. *Nature Climate Change*, 8(4):275–281, 2018.
- [19] Lucas Benney and Anca Radulescu. An energy balance model of carbon’s effect on climate change, 2015.
- [20] Jorge Castillo-Mateo, Alan E. Gelfand, Zeus Gracia-Tabuenca, Jesús Asín, and Ana C. Cebrián. Spatio-temporal modeling for record-breaking temperature events in Spain, 2024.

-
- [21] Jorge Castillo-Mateo, Miguel Lafuente, Jesús Asín, Ana C. Cebrián, Alan E. Gelfand, and Jesús Abaurrea. Spatial modeling of day-within-year temperature time series: an examination of daily maximum temperatures in aragón, spain, 2022.
- [22] Tao Li, Lixing Wang, Zihan Qiu, Philippe Ciais, Taochun Sun, Matthew W. Jones, Robbie M. Andrew, Glen P. Peters, Piyu ke, Xiaoting Huang, Robert B. Jackson, and Zhu Liu. Reconstructing global daily co2 emissions via machine learning, 2024.
- [23] Giuliano Vitali, Sergei Rogosin, and Guido Baldoni. Climate change and grain production fluctuations, 2019.
- [24] Michael Bertolacci, Andrew Zammit-Mangion, Andrew Schuh, Beata Bukosa, Jenny Fisher, Yi Cao, Aleya Kaushik, and Noel Cressie. Inferring changes to the global carbon cycle with wombat v2.0, a hierarchical flux-inversion framework, 2022.
- [25] Meihua Zhou, Nan Wan, Tianlong Zheng, Hanwen Xu, Li Yang, and Tingting Wang. New insights into global warming: End-to-end visual analysis and prediction of temperature variations, 2024.
- [26] Bora Akgun, Zeynep Isvan, Levent Tuter, and Mehmet Levent Kurnaz. Climate prediction through statistical methods, 2008.
- [27] Marta Magnani, Rubén Díaz-Sierra, Luke Sweeney, Antonello Provenzale, and Mara Baudena. Fire responses shape plant communities in a minimal model for fire ecosystems across the world, 2023.
- [28] Samuel Asante Gyamerah, Philip Ngare, and Dennis Ikpe. Regime-switching temperature dynamics model for weather derivatives, 2018.
- [29] Nicola Scafetta. Solar and planetary oscillation control on climate change: hind-cast, forecast and a comparison with the cmip5 gcms, 2013.
- [30] Jianyu Chen, Jianyu Hu, Wei Wei, and Jinqiao Duan. A data-driven approach for discovering the most probable transition pathway for a stochastic carbon cycle system, 2022.
- [31] Jyrki Kauppinen and Pekka Malmi. No experimental evidence for the significant anthropogenic climate change, 2019.
- [32] Timothy M. Merlis and Tapio Schneider. Changes in zonal surface temperature gradients and walker circulations in a wide range of climates, 2011.
- [33] Jessie Lydia Henshaw. The systemic growth constants of climate change: From its origin in 1780 to its major post-wwii acceleration, 2019.
- [34] Huang Huang, Dorit Hammerling, Bo Li, and Richard Smith. Combining interdependent climate model outputs in cmip5: A spatial bayesian approach, 2020.
- [35] Manmeet Singh, Chirag Dhara, Adarsh Kumar, Sukhpal Singh Gill, and Steve Uhlig. Quantum artificial intelligence for the science of climate change, 2021.
- [36] Kerstin Hötte and Su Jung Jee. Knowledge for a warmer world: a patent analysis of climate change adaptation technologies, 2022.
- [37] Casey Berger, Agustin Di Paolo, Tracey Forrest, Stuart Hadfield, Nicolas Sawaya, Michał Stęchły, and Karl Thibault. Quantum technologies for climate change: Preliminary assessment, 2021.
- [38] Kgomotso S. Morupisi and Chris J. Budd. An analysis of the periodically forced pp04 climate model, using the theory of non-smooth dynamical systems, 2020.
- [39] Peter A. G. Watson. Applying machine learning to improve simulations of a chaotic dynamical system using empirical error correction, 2019.
- [40] R. C. T. Rainey. The earths long-term climate changes and ice ages: a derivation of milankovitch cycles from first principles, 2021.

-
- [41] Natalie J. Briscoe, Shane D. Morris, Paul D. Mathewson, Lauren B. Buckley, Marko Jusup, Ofir Levy, Ilya M. D. Maclean, Sylvain Pincebourde, Eric A. Riddell, Jessica A. Roberts, Rafael Schouten, Michael W. Sears, and Michael R. Kearney. Mechanistic forecasts of species responses to climate change: the promise of biophysical ecology, 2022.
- [42] Sylvain Gretchko, Jessa Marley, and Rebecca C. Tyson. The effects of climate change on predator-prey dynamics, 2018.
- [43] Adam Giammarese, Jacob Brown, and Nishant Malik. Reconfiguration of amazon's connectivity in the climate system, 2023.
- [44] Hyun Choi, Ali Sadeghian, Sergio Marconi, Ethan White, and Daisy Zhe Wang. Measuring impact of climate change on tree species: analysis of jsdm on fia data, 2019.
- [45] Hideyasu Shimadzu, Maria Dornelas, and Anne E. Magurran. Measuring temporal turnover in ecological communities, 2016.
- [46] Lionel Roques, Alain Roques, Henri Berestycki, and André Kretzschmar. A population facing climate change: joint influences of allee effects and environmental boundary geometry, 2009.
- [47] Claire L. Harris, Neil Brummitt, Christina A. Cobbold, and Richard Reeve. Dynamic virtual ecosystems as a tool for detecting large-scale responses of biodiversity to environmental and land-use change, 2023.
- [48] Greta T Pecl, Miguel B Araújo, Johann D Bell, Julia Blanchard, Timothy C Bonebrake, I-Ching Chen, Timothy D Clark, Robert K Colwell, Finn Danielsen, Birgitta Evengård, et al. Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, 355(6332):eaai9214, 2017.
- [49] Sergey A. Vakulenko, Ivan Sudakov, Sergei V. Petrovskii, and Dmitry V. Lukichev. Stability of a planetary climate system with the biosphere competing for resources, 2020.
- [50] Mauro Gobbi. Global warning: challenges, threats and opportunities for ground beetles (coleoptera: Carabidae) in high altitude habitats, 2020.
- [51] Lynn Wahab, Ezzat Chebaro, Jad Ismail, Amir Nasrelddine, and Ali El-Zein. A predictive model for geographic distributions of mangroves, 2021.
- [52] Chirag Dhara. Constraining global changes in temperature and precipitation from observable changes in surface radiative heating, 2020.
- [53] Andrew Poppick, Elisabeth J. Moyer, and Michael L. Stein. Estimating trends in the global mean temperature record, 2017.
- [54] J. M. Finkel and J. I. Katz. Changing u.s. extreme temperature statistics, 2017.
- [55] James Hansen, Makiko Sato, Pushker Kharecha, Karina von Schuckmann, David J Beerling, Junji Cao, Shaun Marcott, Valerie Masson-Delmotte, Michael J Prather, Eelco J Rohling, Jeremy Shakun, and Pete Smith. Young people's burden: Requirement of negative co2 emissions, 2017.
- [56] Junze Zhang, Kerry Zhang, Mary Zhang, Jonathan H. Jiang, Philip E. Rosen, and Kristen A. Fahy. Avoiding the "great filter": An assessment of climate change solutions and combinations for effective implementation, 2022.
- [57] Henri Berestycki, Romain Ducasse, and Luca Rossi. Influence of a road on a population in an ecological niche facing climate change, 2019.
- [58] Simone Vincenzi and Michele Bellingeri. Consequences of catastrophic disturbances on population persistence and adaptations, 2011.
- [59] Zhu Liu, Zhu Deng, Philippe Ciais, Jianguang Tan, Biqing Zhu, Steven J. Davis, Robbie Andrew, Olivier Boucher, Simon Ben Arous, Pep Canadel, Xinyu Dou, Pierre Friedlingstein, Pierre Gentine, Rui Guo, Chaopeng Hong, Robert B. Jackson, Daniel M. Kammen, Piyu Ke, Corinne Le Quere, Crippa Monica, Greet Janssens-Maenhout, Glen Peters, Katsumasa Tanaka, Yilong Wang, Bo Zheng, Haiwang Zhong, Taochun Sun, and Hans Joachim Schellnhuber. Global daily co₂ emissions for the year 2020, 2021.

-
- [60] Nicola Scafetta. Global temperatures and sunspot numbers. are they related? yes, but non linearly. a reply to gil-alana et al. (2014), 2014.
- [61] Adriano Mazzarella and Nicola Scafetta. Evidences for a quasi 60-year north atlantic oscillation since 1700 and its meaning for global climate change, 2012.
- [62] Andreas Morr and Niklas Boers. Detection of approaching critical transitions in natural systems driven by red noise, 2024.
- [63] Sandra Skendžić, Monika Zovko, Ivana Pajač Živković, Vinko Lešić, and Darija Lemić. The impact of climate change on agricultural insect pests. *Insects*, 12(5):440, 2021.
- [64] David Dunn and James P. Crutchfield. Entomogenic climate change, 2008.
- [65] Clément Le Priol, Joy M. Monteiro, and Freddy Bouchet. Using rare event algorithms to understand the statistics and dynamics of extreme heatwave seasons in south asia, 2024.
- [66] Zhiang Xie, Dongwei Chen, and Puxi Li. Rediscover climate change during global warming slowdown via wasserstein stability analysis, 2023.
- [67] Frederik Van Daele, Olivier Honnay, Steven Janssens, and Hanne De Kort. Habitat fragmentation affects climate adaptation in a forest herb, 2023.
- [68] Jun Chen, Jordy O Rodriguez Rincon, Gloria DeGrandi-Hoffman, Jennifer Fewell, Jon Harrison, and Yun Kang. Impacts of seasonality and parasitism on honey bee population dynamics, 2023.
- [69] Fernando M. Aragón, Francisco Oteiza, and Juan Pablo Rud. Climate change and agriculture: Subsistence farmers’ response to extreme heat, 2019.
- [70] Sesa Singha Roy, Reid Tingley, and Alan Dorin. Image background assessment as a novel technique for insect microhabitat identification, 2023.
- [71] M. Tuomi, T. Thum, H. Järvinen, S. Fronzek, B. Berg, M. Harmon, J. A. Trofymow, S. Sevanto, and J. Liski. Leaf litter decomposition – estimates of global variability based on yasso07 model, 2009.
- [72] Tyler Poppenwimer, Itay Mayrose, and Niv DeMalach. Revising the global biogeography of annual and perennial plants, 2023.
- [73] L. Mark W. Leggett and David. A. Ball. Forecasts of the trend in global-mean temperature to 2100 arising from the scenarios of first-difference co2 and peak fossil fuel, 2020.
- [74] Anthony J. Webster. Lightning strikes and attribution of climatic change, 2015.
- [75] Sebastian Sippel, Jakob Zscheischler, Martin Heimann, Holger Lange, Miguel D. Mahecha, Geert Jan van Oldenborgh, Friederike E. L. Otto, and Markus Reichstein. Have precipitation extremes and annual totals been increasing in the world’s dry regions over the last 60 years?, 2016.
- [76] Angie Michaiel and Amy Bernard. Neurobiology and changing ecosystems: toward understanding the impact of anthropogenic influences on neurons and circuits, 2022.
- [77] R. Juhász and I. A. Kovács. Population boundary across an environmental gradient: Effects of quenched disorder, 2020.
- [78] Donald C. Morton. An astronomer’s view of climate change, 2014.
- [79] Gerald E. Marsh. Climate change: The sun’s role, 2007.
- [80] Ranjini Swaminathan, Jacob Schewe, Jeremy Walton, Klaus Zimmermann, Colin Jones, Richard A. Betts, Chantelle Burton, Chris D. Jones, Matthias Mengel, Christopher P. O. Reyer, Andrew G. Turner, and Katja Weigel. Regional impacts poorly constrained by climate sensitivity, 2024.

-
- [81] Samer Fawzy, Ahmed I Osman, John Doran, and David W Rooney. Strategies for mitigation of climate change: a review. *Environmental Chemistry Letters*, 18:2069–2094, 2020.
- [82] B G Sidharth and B S Lakshmi. A model for long term climate change, 2003.
- [83] Ricardo Cavicchioli, William J Ripple, Kenneth N Timmis, Farooq Azam, Lars R Bakken, Matthew Baylis, Michael J Behrenfeld, Antje Boetius, Philip W Boyd, Aimée T Classen, et al. Scientists’ warning to humanity: microorganisms and climate change. *Nature Reviews Microbiology*, 17(9):569–586, 2019.

www.SurveyX.cn

Disclaimer:

SurveyX is an AI-powered system designed to automate the generation of surveys. While it aims to produce high-quality, coherent, and comprehensive surveys with accurate citations, the final output is derived from the AI's synthesis of pre-processed materials, which may contain limitations or inaccuracies. As such, the generated content should not be used for academic publication or formal submissions and must be independently reviewed and verified. The developers of SurveyX do not assume responsibility for any errors or consequences arising from the use of the generated surveys.

www.SurveyX.cn