

# Lenstra Elliptic Curve Factorization

Miles Benton and Skip Moses

MATH 317

2021



- Hendrik Lenstra Jr. received his doctorate from the University of Amsterdam in 1977.



- Hendrik Lenstra Jr. received his doctorate from the University of Amsterdam in 1977.
- Discovered Elliptic Curve Factorization (ECM) in 1987.



- Hendrik Lenstra Jr. received his doctorate from the University of Amsterdam in 1977.
- Discovered Elliptic Curve Factorization (ECM) in 1987.
- ECM is third-fastest known factoring algorithm and the best algorithm for finding divisors not exceeding 50-60 digits.



- Hendrik Lenstra Jr. received his doctorate from the University of Amsterdam in 1977.
- Discovered Elliptic Curve Factorization (ECM) in 1987.
- ECM is third-fastest known factoring algorithm and the best algorithm for finding divisors not exceeding 50-60 digits.
- The largest factor found using ECM has 83 digits.

# Group Theory behind Pollard $p - 1$

# Group Theory behind Pollard $p - 1$

**Definition** Let  $(G, +)$  be a group. If  $H \subset G$  is also a group under  $+$ , then we call  $H$  a subgroup of  $G$ .

# Group Theory behind Pollard $p - 1$

**Definition** Let  $(G, +)$  be a group. If  $H \subset G$  is also a group under  $+$ , then we call  $H$  a subgroup of  $G$ .

**Lagrange's Theorem** If  $H \subset G$  with  $|G| < \infty$ , then  $|H|$  divides  $|G|$ .



# Group Theory behind Pollard $p - 1$

**Definition** Let  $(G, +)$  be a group. If  $H \subset G$  is also a group under  $+$ , then we call  $H$  a subgroup of  $G$ .

**Lagrange's Theorem** If  $H \subset G$  with  $|G| < \infty$ , then  $|H|$  divides  $|G|$ .

- Define the relation  $a \sim b$  whenever  $a = bh$  for some  $h \in H$ .

# Group Theory behind Pollard $p - 1$

**Definition** Let  $(G, +)$  be a group. If  $H \subset G$  is also a group under  $+$ , then we call  $H$  a subgroup of  $G$ .

**Lagrange's Theorem** If  $H \subset G$  with  $|G| < \infty$ , then  $|H|$  divides  $|G|$ .

- Define the relation  $a \sim b$  whenever  $a = bh$  for some  $h \in H$ .
- Let  $S$  be an equivalence class of  $\sim$  and pick an arbitrary  $a \in S$ .

**Definition** Let  $(G, +)$  be a group. If  $H \subset G$  is also a group under  $+$ , then we call  $H$  a subgroup of  $G$ .

**Lagrange's Theorem** If  $H \subset G$  with  $|G| < \infty$ , then  $|H|$  divides  $|G|$ .

- Define the relation  $a \sim b$  whenever  $a = bh$  for some  $h \in H$ .
- Let  $S$  be an equivalence class of  $\sim$  and pick an arbitrary  $a \in S$ .
- Define  $f : S \rightarrow H$ , where  $f(b) = b^{-1}a$  and  $g : H \rightarrow S$  by  $g(h) = ah^{-1}$

**Definition** Let  $(G, +)$  be a group. If  $H \subset G$  is also a group under  $+$ , then we call  $H$  a subgroup of  $G$ .

**Lagrange's Theorem** If  $H \subset G$  with  $|G| < \infty$ , then  $|H|$  divides  $|G|$ .

- Define the relation  $a \sim b$  whenever  $a = bh$  for some  $h \in H$ .
- Let  $S$  be an equivalence class of  $\sim$  and pick an arbitrary  $a \in S$ .
- Define  $f : S \rightarrow H$ , where  $f(b) = b^{-1}a$  and  $g : H \rightarrow S$  by  $g(h) = ah^{-1}$
- Observe  $f$  and  $g$  are inverses.

$$f(g(h)) = f(ah^{-1}) = (ah^{-1})^{-1}a = h$$

$$g(f(b)) = g(b^{-1}a) = a(b^{-1}a)^{-1} = b$$

- Thus,  $|S| = |H|$  for all equivalence classes  $S$ .

# Group Theory behind Pollard $p - 1$

**Definition** Let  $(G, +)$  be a group. If  $H \subset G$  is also a group under  $+$ , then we call  $H$  a subgroup of  $G$ .

**Lagrange's Theorem** If  $H \subset G$  with  $|G| < \infty$ , then  $|H|$  divides  $|G|$ .

- Define the relation  $a \sim b$  whenever  $a = bh$  for some  $h \in H$ .
- Let  $S$  be an equivalence class of  $\sim$  and pick an arbitrary  $a \in S$ .
- Define  $f : S \rightarrow H$ , where  $f(b) = b^{-1}a$  and  $g : H \rightarrow S$  by  $g(h) = ah^{-1}$
- Observe  $f$  and  $g$  are inverses.

$$f(g(h)) = f(ah^{-1}) = (ah^{-1})^{-1}a = h$$

$$g(f(b)) = g(b^{-1}a) = a(b^{-1}a)^{-1} = b$$

- Thus,  $|S| = |H|$  for all equivalence classes  $S$ .
- Therefore,  $|G| = n|H|$ .

# A different perspective

**Lemma 2.2.5** Suppose that  $m, n \in \mathbb{N}$  and  $\gcd(m, n) = 1$ . Then the map


$$\psi : (\mathbb{Z}/mn\mathbb{Z})^* \rightarrow (\mathbb{Z}/m\mathbb{Z})^* \times (\mathbb{Z}/n\mathbb{Z})^*$$

defined by

$$\psi(c) = (c \pmod{m}, c \pmod{n})$$

is a bijection.

# Example





# Example

- Let  $B_i = \text{lcm}(1, \dots, i)$ .



## Example

- Let  $B_i = \text{lcm}(1, \dots, i)$ .

| $B_i$ | $2^{B_i} \pmod{1763}$ | $(2^i \pmod{41}, 2^i \pmod{43})$ |
|-------|-----------------------|----------------------------------|
| 1     | 2                     | (2, 2)                           |

# Example

- Let  $B_i = \text{lcm}(1, \dots, i)$ .

| $B_i$ | $2^{B_i} \pmod{1763}$ | $(2^i \pmod{41}, 2^i \pmod{43})$ |
|-------|-----------------------|----------------------------------|
| 1     | 2                     | (2, 2)                           |
| 2     | 4                     | (4, 4)                           |

# Example

- Let  $B_i = \text{lcm}(1, \dots, i)$ .

| $B_i$ | $2^{B_i} \pmod{1763}$ | $(2^i \pmod{41}, 2^i \pmod{43})$ |
|-------|-----------------------|----------------------------------|
| 1     | 2                     | (2, 2)                           |
| 2     | 4                     | (4, 4)                           |
| 6     | 570                   | (37, 11)                         |

# Example

- Let  $B_i = \text{lcm}(1, \dots, i)$ .

| $B_i$ | $2^{B_i} \pmod{1763}$ | $(2^i \pmod{41}, 2^i \pmod{43})$ |
|-------|-----------------------|----------------------------------|
| 1     | 2                     | (2, 2)                           |
| 2     | 4                     | (4, 4)                           |
| 6     | 570                   | (37, 11)                         |
| 60    | 575                   | (1, 16)                          |

- We compute  $\gcd(574, 1763) = 41$

# Set up for ECM

- Let  $E$  be an elliptic curve over  $\mathbb{Z}/N\mathbb{Z}$  of the form

$$y^2 = x^3 + ax + 1$$

such that  $4a^3 + 27 \in (\mathbb{Z}/N\mathbb{Z})^*$ . This forces non singularity and ensures  $P = (0, 1)$  is on the curve.

- Let  $E$  be an elliptic curve over  $\mathbb{Z}/N\mathbb{Z}$  of the form

$$y^2 = x^3 + ax + 1$$

such that  $4a^3 + 27 \in (\mathbb{Z}/N\mathbb{Z})^*$ . This forces non singularity and ensures  $P = (0, 1)$  is on the curve.

- Definition 6.3.1 (Power Smooth). Let  $B$  be a positive integer. If  $n$  is a positive integer with prime factorization

$$n = \prod p_i^{e_i},$$

then  $n$  is  $B$ -power smooth if  $p_i^{e_i} \leq B$  for all  $i$ .



- Let  $E$  be an elliptic curve over  $\mathbb{Z}/N\mathbb{Z}$  of the form

$$y^2 = x^3 + ax + 1$$

such that  $4a^3 + 27 \in (\mathbb{Z}/N\mathbb{Z})^*$ . This forces non singularity and ensures  $P = (0, 1)$  is on the curve.

- Definition 6.3.1 (Power Smooth). Let  $B$  be a positive integer. If  $n$  is a positive integer with prime factorization

$$n = \prod p_i^{e_i},$$

then  $n$  is  $B$ -power smooth if  $p_i^{e_i} \leq B$  for all  $i$ .

- Example  $30 = 2 \cdot 3 \cdot 5$  is  $B$  power smooth for  $B \geq 5$ , but  $150 = 2 \cdot 3 \cdot 5^2$  is not 5-power smooth.

# Motivation

- Fix  $B \in \mathbb{N}$ . Let  $p \in \mathbb{N}$  such that  $p - 1$  is not  $B$ - power smooth.

# Motivation

- Fix  $B \in \mathbb{N}$ . Let  $p \in \mathbb{N}$  such that  $p - 1$  is not  $B$ - power smooth.
- Recall, in Pollard  $p - 1$ , this would be equivalent to not having  $p - 1 \nmid m = \text{lcm}(1, 2, \dots, B)$ ; i.e.  $a^m \not\equiv 1 \pmod{p}$ .

- Fix  $B \in \mathbb{N}$ . Let  $p \in \mathbb{N}$  such that  $p - 1$  is not  $B$ -power smooth.
- Recall, in Pollard  $p - 1$ , this would be equivalent to not having  $p - 1 \nmid m = \text{lcm}(1, 2, \dots, B)$ ; i.e.  $a^m \not\equiv 1 \pmod{p}$ .
- On the interval  $[10^{15}, 10^{15} + 10000]$  15 percent of the primes  $p$  are such that  $p - 1$  is not  $10^6$ -power smooth.

# Motivation

- Fix  $B \in \mathbb{N}$ . Let  $p \in \mathbb{N}$  such that  $p - 1$  is not  $B$ -power smooth.
- Recall, in Pollard  $p - 1$ , this would be equivalent to not having  $p - 1 \nmid m = \text{lcm}(1, 2, \dots, B)$ ; i.e.  $a^m \not\equiv 1 \pmod{p}$ .
- On the interval  $[10^{15}, 10^{15} + 10000]$  15 percent of the primes  $p$  are such that  $p - 1$  is not  $10^6$ -power smooth.
- The idea of ECM is to replace modular exponentiation on  $(\mathbb{Z}/N\mathbb{Z})^*$  by repeated addition of points on  $E((\mathbb{Z}/N\mathbb{Z})^*)$

- Fix  $B \in \mathbb{N}$ . Let  $p \in \mathbb{N}$  such that  $p - 1$  is not  $B$ -power smooth.
- Recall, in Pollard  $p - 1$ , this would be equivalent to not having  $p - 1 \nmid m = \text{lcm}(1, 2, \dots, B)$ ; i.e.  $a^m \not\equiv 1 \pmod{p}$ .
- On the interval  $[10^{15}, 10^{15} + 10000]$  15 percent of the primes  $p$  are such that  $p - 1$  is not  $10^6$ -power smooth.
- The idea of ECM is to replace modular exponentiation on  $(\mathbb{Z}/N\mathbb{Z})^*$  by repeated addition of points on  $E((\mathbb{Z}/N\mathbb{Z})^*)$
- Recall, by the Hasse-Weil bound we can reduce the size of our group by  $2 \cdot \sqrt{p}$ .

Algorithm 6.3.10 (Elliptic Curve Factorization Method). Let  $N$  and  $B$  be positive integers.



Algorithm 6.3.10 (Elliptic Curve Factorization Method). Let  $N$  and  $B$  be positive integers.

1. Compute  $m = \text{lcm}(1, 2, \dots, B)$ .

# Elliptic Curve Factorization

Algorithm 6.3.10 (Elliptic Curve Factorization Method). Let  $N$  and  $B$  be positive integers.

1. Compute  $m = \text{lcm}(1, 2, \dots, B)$ .
2. Choose  $a \in \mathbb{Z}/N\mathbb{Z}$  such that  $4a^3 + 27 \in (\mathbb{Z}/N\mathbb{Z})^*$ . This forces  $P = (0, 1)$  to be a point on  $y^2 = x^3 + ax + 1$  over  $\mathbb{Z}/N\mathbb{Z}$ .

# Elliptic Curve Factorization

Algorithm 6.3.10 (Elliptic Curve Factorization Method). Let  $N$  and  $B$  be positive integers.

1. Compute  $m = \text{lcm}(1, 2, \dots, B)$ .
2. Choose  $a \in \mathbb{Z}/N\mathbb{Z}$  such that  $4a^3 + 27 \in (\mathbb{Z}/N\mathbb{Z})^*$ . This forces  $P = (0, 1)$  to be a point on  $y^2 = x^3 + ax + 1$  over  $\mathbb{Z}/N\mathbb{Z}$ .
3. Try to compute  $mP$ . If at some point we cannot compute a sum of points, then some denominator  $g$  is not coprime to  $N$ , then  $\gcd(g, N)$  is a nontrivial divisor of  $N$ .

# Analogy to Pollard $p - 1$



## Analogy to Pollard $p - 1$

**Table:** Let  $E$  be an elliptic curve, and  $m = \text{lcm}(1, 2, \dots, B)$  for some  $B$

A blank sheet of white graph paper with a light gray grid. The grid consists of small squares. A vertical margin line is present on the left side, creating a narrow column. A horizontal margin line is present at the top, creating a narrow header row. The intersection of these two lines forms a small square in the top-left corner.

# Analogy to Pollard $p - 1$

**Table:** Let  $E$  be an elliptic curve, and  $m = \text{lcm}(1, 2, \dots, B)$  for some  $B$

| Pollard $p - 1$ | ECM |
|-----------------|-----|
|                 |     |
|                 |     |
|                 |     |
|                 |     |

# Analogy to Pollard $p - 1$

**Table:** Let  $E$  be an elliptic curve, and  $m = \text{lcm}(1, 2, \dots, B)$  for some  $B$

| <b>Pollard <math>p - 1</math></b> | <b>ECM</b>                  |
|-----------------------------------|-----------------------------|
| $\mathbb{Z}/N\mathbb{Z}$          | $E(\mathbb{Z}/N\mathbb{Z})$ |
|                                   |                             |
|                                   |                             |
|                                   |                             |

# Analogy to Pollard $p - 1$

**Table:** Let  $E$  be an elliptic curve, and  $m = \text{lcm}(1, 2, \dots, B)$  for some  $B$

| <b>Pollard <math>p - 1</math></b>  | <b>ECM</b>                  |
|------------------------------------|-----------------------------|
| $\mathbb{Z}/N\mathbb{Z}$           | $E(\mathbb{Z}/N\mathbb{Z})$ |
| $g \in (\mathbb{Z}/N\mathbb{Z})^*$ | $(0, 1)$                    |
|                                    |                             |
|                                    |                             |



# Analogy to Pollard $p - 1$

**Table:** Let  $E$  be an elliptic curve, and  $m = \text{lcm}(1, 2, \dots, B)$  for some  $B$

| <b>Pollard <math>p - 1</math></b>  | <b>ECM</b>                            |
|------------------------------------|---------------------------------------|
| $\mathbb{Z}/N\mathbb{Z}$           | $E(\mathbb{Z}/N\mathbb{Z})$           |
| $g \in (\mathbb{Z}/N\mathbb{Z})^*$ | $(0, 1)$                              |
| $g^m \equiv 1 \pmod{N}$            | $mP \notin E(\mathbb{Z}/N\mathbb{Z})$ |

# Analogy to Pollard $p - 1$

**Table:** Let  $E$  be an elliptic curve, and  $m = \text{lcm}(1, 2, \dots, B)$  for some  $B$

| <b>Pollard <math>p - 1</math></b>  | <b>ECM</b>                            |
|------------------------------------|---------------------------------------|
| $\mathbb{Z}/N\mathbb{Z}$           | $E(\mathbb{Z}/N\mathbb{Z})$           |
| $g \in (\mathbb{Z}/N\mathbb{Z})^*$ | $(0, 1)$                              |
| $g^m \equiv 1 \pmod{N}$            | $mP \notin E(\mathbb{Z}/N\mathbb{Z})$ |
| $\gcd(g^m - 1, N)$                 | $\gcd(m, N)$                          |

- If Pollard  $p - 1$  fails, we have no choice but to increase  $B$ .

# Analogy to Pollard $p - 1$

**Table:** Let  $E$  be an elliptic curve, and  $m = \text{lcm}(1, 2, \dots, B)$  for some  $B$

| <b>Pollard <math>p - 1</math></b>  | <b>ECM</b>                            |
|------------------------------------|---------------------------------------|
| $\mathbb{Z}/N\mathbb{Z}$           | $E(\mathbb{Z}/N\mathbb{Z})$           |
| $g \in (\mathbb{Z}/N\mathbb{Z})^*$ | $(0, 1)$                              |
| $g^m \equiv 1 \pmod{N}$            | $mP \notin E(\mathbb{Z}/N\mathbb{Z})$ |
| $\gcd(g^m - 1, N)$                 | $\gcd(m, N)$                          |

- If Pollard  $p - 1$  fails, we have no choice but to increase  $B$ .
- However, ECM has a second option. We can choose another random elliptic curve.

# Why ECM "Works"

# Why ECM "Works"

We can consider an analogous mapping

$$g : E(\mathbb{Z}/N\mathbb{Z}) \rightarrow \prod_{p|N} E(\mathbb{Z}/p\mathbb{Z})$$

where  $p$  are prime divisors of  $N$ .

# Why ECM "Works"

We can consider an analogous mapping

$$g : E(\mathbb{Z}/N\mathbb{Z}) \rightarrow \prod_{p|N} E(\mathbb{Z}/p\mathbb{Z})$$

where  $p$  are prime divisors of  $N$ .

- Note the quotations. There is a subtlety in the difference between  $E(\mathbb{Z}/N\mathbb{Z})$  and  $\mathbb{Z}/N\mathbb{Z}$ .

# Why ECM "Works"

We can consider an analogous mapping

$$"g : E(\mathbb{Z}/N\mathbb{Z}) \rightarrow \prod_{p|N} E(\mathbb{Z}/p\mathbb{Z})"$$

where  $p$  are prime divisors of  $N$ .

- Note the quotations. There is a subtlety in the difference between  $E(\mathbb{Z}/N\mathbb{Z})$  and  $\mathbb{Z}/N\mathbb{Z}$ .
- Let  $P = (0 : 1 : 1) \in E(\mathbb{Z}/1763\mathbb{Z})$   
 $P_1 = (0 : 1 : 1) \in E(\mathbb{Z}/41\mathbb{Z})$  and  $P_2 = (0 : 1 : 1) \in E(\mathbb{Z}/43\mathbb{Z})$

# Example

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|



# Example

| $i$ | $i * P_1$ | $i * P_2$ | $i * P$ |
|-----|-----------|-----------|---------|
|     |           |           |         |

## Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$ |
|-----|---------------|---------------|---------|
| 0   | $(0 : 1 : 1)$ | $(0 : 1 : 1)$ |         |

## Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$       |
|-----|---------------|---------------|---------------|
| 0   | $(0 : 1 : 1)$ | $(0 : 1 : 1)$ | $(0 : 1 : 1)$ |

## Example

| $i$ | $i * P_1$    | $i * P_2$    | $i * P$     |
|-----|--------------|--------------|-------------|
| 0   | (0 : 1 : 1)  | (0 : 1 : 1)  | (0 : 1 : 1) |
| 1   | (1 : 39 : 1) | (1 : 41 : 1) |             |

# Example

| $i$ | $i * P_1$      | $i * P_2$      | $i * P$          |
|-----|----------------|----------------|------------------|
| 0   | $(0 : 1 : 1)$  | $(0 : 1 : 1)$  | $(0 : 1 : 1)$    |
| 1   | $(1 : 39 : 1)$ | $(1 : 41 : 1)$ | $(1 : 1761 : 1)$ |

## Example

| $i$ | $i * P_1$    | $i * P_2$    | $i * P$        |
|-----|--------------|--------------|----------------|
| 0   | (0 : 1 : 1)  | (0 : 1 : 1)  | (0 : 1 : 1)    |
| 1   | (1 : 39 : 1) | (1 : 41 : 1) | (1 : 1761 : 1) |
| 2   | (8 : 23 : 1) | (8 : 23 : 1) |                |

# Example

| $i$ | $i * P_1$      | $i * P_2$      | $i * P$          |
|-----|----------------|----------------|------------------|
| 0   | $(0 : 1 : 1)$  | $(0 : 1 : 1)$  | $(0 : 1 : 1)$    |
| 1   | $(1 : 39 : 1)$ | $(1 : 41 : 1)$ | $(1 : 1761 : 1)$ |
| 2   | $(8 : 23 : 1)$ | $(8 : 23 : 1)$ | $(8 : 23 : 1)$   |

# Example

| $i$ | $i * P_1$       | $i * P_2$       | $i * P$          |
|-----|-----------------|-----------------|------------------|
| 0   | $(0 : 1 : 1)$   | $(0 : 1 : 1)$   | $(0 : 1 : 1)$    |
| 1   | $(1 : 39 : 1)$  | $(1 : 41 : 1)$  | $(1 : 1761 : 1)$ |
| 2   | $(8 : 23 : 1)$  | $(8 : 23 : 1)$  | $(8 : 23 : 1)$   |
| 3   | $(38 : 38 : 1)$ | $(13 : 17 : 1)$ |                  |



# Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$           |
|-----|---------------|---------------|-------------------|
| 0   | (0 : 1 : 1)   | (0 : 1 : 1)   | (0 : 1 : 1)       |
| 1   | (1 : 39 : 1)  | (1 : 41 : 1)  | (1 : 1761 : 1)    |
| 2   | (8 : 23 : 1)  | (8 : 23 : 1)  | (8 : 23 : 1)      |
| 3   | (38 : 38 : 1) | (13 : 17 : 1) | (1432 : 1350 : 1) |
| 4   | (23 : 23 : 1) | (2 : 23 : 1)  |                   |

# Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$           |
|-----|---------------|---------------|-------------------|
| 0   | (0 : 1 : 1)   | (0 : 1 : 1)   | (0 : 1 : 1)       |
| 1   | (1 : 39 : 1)  | (1 : 41 : 1)  | (1 : 1761 : 1)    |
| 2   | (8 : 23 : 1)  | (8 : 23 : 1)  | (8 : 23 : 1)      |
| 3   | (38 : 38 : 1) | (13 : 17 : 1) | (1432 : 1350 : 1) |
| 4   | (23 : 23 : 1) | (2 : 23 : 1)  | (1335 : 23 : 1)   |
| 5   | (20 : 28 : 1) | (33 : 23 : 1) |                   |

# Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$           |
|-----|---------------|---------------|-------------------|
| 0   | (0 : 1 : 1)   | (0 : 1 : 1)   | (0 : 1 : 1)       |
| 1   | (1 : 39 : 1)  | (1 : 41 : 1)  | (1 : 1761 : 1)    |
| 2   | (8 : 23 : 1)  | (8 : 23 : 1)  | (8 : 23 : 1)      |
| 3   | (38 : 38 : 1) | (13 : 17 : 1) | (1432 : 1350 : 1) |
| 4   | (23 : 23 : 1) | (2 : 23 : 1)  | (1335 : 23 : 1)   |
| 5   | (20 : 28 : 1) | (33 : 23 : 1) | (635 : 1012 : 1)  |

# Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$           |
|-----|---------------|---------------|-------------------|
| 0   | (0 : 1 : 1)   | (0 : 1 : 1)   | (0 : 1 : 1)       |
| 1   | (1 : 39 : 1)  | (1 : 41 : 1)  | (1 : 1761 : 1)    |
| 2   | (8 : 23 : 1)  | (8 : 23 : 1)  | (8 : 23 : 1)      |
| 3   | (38 : 38 : 1) | (13 : 17 : 1) | (1432 : 1350 : 1) |
| 4   | (23 : 23 : 1) | (2 : 23 : 1)  | (1335 : 23 : 1)   |
| 5   | (20 : 28 : 1) | (33 : 23 : 1) | (635 : 1012 : 1)  |
| 6   | (26 : 9 : 1)  | (20 : 0 : 1)  |                   |

# Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$           |
|-----|---------------|---------------|-------------------|
| 0   | (0 : 1 : 1)   | (0 : 1 : 1)   | (0 : 1 : 1)       |
| 1   | (1 : 39 : 1)  | (1 : 41 : 1)  | (1 : 1761 : 1)    |
| 2   | (8 : 23 : 1)  | (8 : 23 : 1)  | (8 : 23 : 1)      |
| 3   | (38 : 38 : 1) | (13 : 17 : 1) | (1432 : 1350 : 1) |
| 4   | (23 : 23 : 1) | (2 : 23 : 1)  | (1335 : 23 : 1)   |
| 5   | (20 : 28 : 1) | (33 : 23 : 1) | (635 : 1012 : 1)  |
| 6   | (26 : 9 : 1)  | (20 : 0 : 1)  | (149 : 1075 : 1)  |

# Example

| $i$ | $i * P_1$     | $i * P_2$     | $i * P$                      |
|-----|---------------|---------------|------------------------------|
| 0   | (0 : 1 : 1)   | (0 : 1 : 1)   | (0 : 1 : 1)                  |
| 1   | (1 : 39 : 1)  | (1 : 41 : 1)  | (1 : 1761 : 1)               |
| 2   | (8 : 23 : 1)  | (8 : 23 : 1)  | (8 : 23 : 1)                 |
| 3   | (38 : 38 : 1) | (13 : 17 : 1) | (1432 : 1350 : 1)            |
| 4   | (23 : 23 : 1) | (2 : 23 : 1)  | (1335 : 23 : 1)              |
| 5   | (20 : 28 : 1) | (33 : 23 : 1) | (635 : 1012 : 1)             |
| 6   | (26 : 9 : 1)  | (20 : 0 : 1)  | (149 : 1075 : 1)             |
| 7   | (10 : 18 : 1) | (33 : 20 : 1) | (420 : 1740 : 1)             |
| 8   | (22 : 19 : 1) | (2 : 20 : 1)  | (432 : 880 : 1)              |
| 9   | (40 : 11 : 1) | (13 : 26 : 1) | (1475 : 585 : 1)             |
| 10  | (19 : 25 : 1) | (8 : 20 : 1)  | (1126 : 1009 : 1)            |
| 11  | (32 : 19 : 1) | (1 : 2 : 1)   | (1549 : 1249 : 1)            |
| 12  | (13 : 25 : 1) | (0 : 42 : 1)  | $\gcd(\text{denom}, N) = 43$ |
| 13  | (12 : 21 : 1) | (0 : 1 : 0)   |                              |

# Implementation

- Generate a random elliptic curve  $E \pmod{N}$  and let  $P = (0, 1)$ .
- Compute  $m = \text{lcm}(1, 2, \dots, B)$ .
- Compute  $mP$  (don't be naive!).
- If the calculation fails, you have found a non-trivial factor of  $N$ .
- Otherwise, just generate a new Elliptic curve and try again.

# Computing $\text{lcm}(1, 2, \dots, B)$

Recall,

$$\text{lcm}(1, 2, \dots, B) = \prod_{p \in P} p^r$$

where  $r = \max\{r \in \mathbb{Z} \mid p^r \leq B\}$ .


$$p^r \leq B$$

$$r \log(p) \leq \log(B)$$

$$r \leq \log_p(B)$$

$$r = \lfloor \log_p(B) \rfloor$$





$$mP = \overbrace{P + P + P \dots P}^{m \text{ times}}$$

A very bad way to compute  $mP$ .


There are many algorithms for computing general elliptic curve point multiplication efficiently, but given the very specific make-up of  $m$ , we can save time by being thoughtful here.

Consider,

$$m_n = q_1^{r_1} \cdot q_2^{r_2} \dots q_n^{r_n}$$

then

$$m_n P = q_n^{r_n} \cdot m_{n-1} P$$



# Coded Example

```
1 def ecm(n, B=10^4, trials=100):
2     R = Zmod(n)
3     primes = list(prime_range(B+1))
4
5     for _ in range(trials):
6         while True:
7             a = R.random_element()
8             if gcd(4 * Integer(a)^3 + 27, n) == 1:
9                 break
10
11         E = EllipticCurve([a, 1])
12         P = E([0,1])
13
14         try:
15             for p in primes:
16                 P = P * p^floor(math.log(B,p))
17
18         except ZeroDivisionError as e:
19             return gcd(Integer(str(e).split()[2]), n)
20
21     return -1
```

